

Rec'd PCT/PTO 25 FEB 2002

10/069282

SPECIFICATION

ANTIANDROGENIC AGENTSTECHNICAL FIELD

5 This invention relates to androstane derivatives having various substituents in 7- or 11-position, substances that act as antagonist against but not as agonist for the androgen receptor, and pharmaceuticals that contain said androstane derivatives and said substances.

10

BACKGROUND ART

 It has become known to date that prostate cancer, prostatomegaly, male pattern alopecia, sexual prematurity, acne vulgaris, seborrhea and hirsutism are closely
15 associated with the male hormone, androgen. For example, it is known that prostate cancer and prostatomegaly are rare in castrated men and patients with gonad dysfunction.

 Already used antiandrogenic agents, or agonists for the androgen receptor, include, for example, cyproterone
20 acetate, chlormadinone acetate, flutamide and bicaltamide. Cyproterone acetate is known to suppress the progress of acne and the onset of baldness in the teens. Cyproterone acetate is also used in women for treatment of masculinization and alopecia. Flutamide and bicaltamide
25 are used as therapeutics for prostatomegaly.

 These antiandrogenic agents have exhibited marked efficacy in many cases including drug therapy of prostate cancer and comprise an important part of the effective

therapeutics. However, one of the problems with these antiandrogenic agents is that even if they exhibit marked efficacy, recurrence is common in almost all cases after the lapse of two to five years; in other words, they are
5 known to induce androgen tolerance.

It was recently reported that hydroxyflutamide, the active essence of flutamide, elevated the transcriptional activity of the androgen receptor at a concentration of 10 mol/L. Plasma levels of hydroxyflutamide in prostate
10 cancer patients under flutamide treatment are several mol/L which, according to the report, is the level at which the agonist action is manifested (see J. Biol. Chem., vol. 270, 19998-20003, 1995). It was also reported that a two-week
15 continuous administration of cyproterone acetate and chlormadinone acetate to castrated rats increased the prostate weight (Folia endocrinol., vol. 66, 597-606, 1990). As for flutamide and bicartamide, cases of side effects such as hepatotoxicity have also been reported.

20 Speaking of the so-called pure antagonists which are substances that act as antagonist against but not as agonist for a nuclear receptor, namely, substances that can completely inhibit the action of the receptor, they have been known for the estrogen receptor (see, for example,
25 WO98/25916, European Patent Publication No. 0138504, USP 4,659,516 and Cancer Res., 1991, 51, 3867). The molecular structures of the hormone-binding domains of nuclear receptors are being unravelled by X-ray crystallography and

the like for RXR (retinoid-X receptor), RAR (retinoic acid receptor) and the like (see, for example, Nature, vol. 375, 377-382, 1995).

W097/49709 discloses androgen receptor modifiers that
5 are nonsteroidal four-ring compounds.

Steroid compounds having an aminocarbonylalkyl group in 7-position or an aminocarbonylalkynyl group in 17-position are known by being described in W091/00732. These are androgen synthesis inhibitors and/or substances that
10 act as antanosit against the androgen receptor and steroid compounds are disclosed that have a freely selectable double bond in 1(2) position, 4(5) position, 6(7) position, 9(10) position and/or 11(12) position in the general formula and the only specific compounds that are disclosed
15 have a double bond in 4(5) position. One of the compounds that are mentioned as the most preferred is EM-101 which is a steroid compound having a 10-(N-butyl-N-methylaminocarbonyl)decyl group in 7 α -position and a hydroxyl group in 17 β -position. However, these compounds
20 have problems such as inadequate antagonist action against the androgen receptor, strong toxicity, etc.

As a steroid compound having an aromatic ring or an alkyloxy group in 11-position, RU486 is described in W095/17192 and known as an agent for dealing with multi-
25 drug tolerance.

DISCLOSURE OF INVENTION

An object of the this invention is to provide

androstane derivatives having various substituents in 7- or 11-position, pharmaceutically acceptable salts thereof, or prodrugs of the derivatives or their salts.


Another object of this invention is to provide
5 substances that act as antagonist against but not as agonist for the androgen receptor, pharmaceutically acceptable salts thereof, or prodrugs of the substances or their salts.

Still another object of this invention is to provide
10 pharmaceuticals that contain said androstane derivatives and pharmaceuticals that contain said substances.


With a view to attaining these objects, the present inventor hypothesized that one of the causes of side effects, such as androgen tolerance and the increase in
15 prostate weight, that are developed by heretofore known antagonists against the androgen receptor is the proliferation of androgen-responsive cells (e.g. prostate cells) due to the agonist action possessed by said antagonists, and anticipated that finding a pure antagonist
20 against the androgen receptor, namely, an antagonist that does not act as agonist for the androgen receptor, would lead to the finding of antiandrogenic agents that do not show any side effects such as the development of androgen tolerance and hepatotoxicity after prolonged administration;
25 the inventor then undertook the designing of said antagonist. To begin with, the androgen receptor was modelled from existing nuclear receptors such as RXR and RAR by the homology technique using software packages such

as Homology (from MSI) and Look (from MAG). Second, it was found that if a pure antagonist against the androgen receptor was designed by using testosterone and/or dihydrotestosterone as a ligand and, with the resulting
 5 model of a complex between said ligand and the androgen receptor being utilized, by introducing into suitable positions those side chains which had suitable lengths and functional groups to form the interaction with the receptor, substances or compounds could be designed that could be
 10 anticipated to act as pure antagonist against the androgen receptor and/or antiandrogenic agents that had lesser side effects such as lower hepatotoxicity; the present invention has been accomplished on the basis of this finding.

According to a first aspect of this invention, there
 15 are provided compounds represented by the general formula (I), pharmaceutically acceptable salts thereof, or prodrugs of the compounds or their salts:



[wherein X^1 and X^2 represent independently a hydrogen atom
 20 or a group represented by the general formula (II)



R^a represents a hydrogen atom or a protective group of a hydroxyl group, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound,

represent an optionally protected $-(C=O)-$, and the dashed line in combination with the solid line represents the formation of a single bond or a double bond;

in addition, Ar represents a single bond or an aromatic hydrocarbon group, A represents a methylene group or $-O-$, R^1 represents an optionally substituted alkyl group, an optionally substituted alkenyl group or an optionally substituted alkynyl group;

provided that X^1 and X^2 are not a hydrogen atom at the same time].

According to a second aspect of this invention, there are provided substances that act as antagonist against but not as agonist for the androgen receptor, pharmaceutically acceptable salts thereof, or prodrugs of the substances or their salts.

According to a third aspect of this invention, there are provided pharmaceuticals that contain compounds represented by the general formula (I), as well as pharmaceuticals that contain substances that act as antagonist against but not as agonist for the androgen receptor.

BEST MODE FOR CARRYING OUT THE INVENTION

In this specification, straight-chained or branched alkyl groups having 1 - 3 carbon atoms include methyl group, ethyl group, n-propyl group and i-propyl group.

Straight-chained or branched alkyl groups having 1 - 6 carbon atoms include, for example, methyl group, ethyl

group, n-propyl group, i-propyl group, n-butyl group, s-butyl group, i-butyl group, t-butyl group, n-pentyl group, 3-methylbutyl group, 2-methylbutyl group, 1-methylbutyl group, 1-ethylpropyl group and n-hexyl group.


5 In this specification, ω position means the terminal position of a divalent group which is other than 1-position. For example, in hexane-1,6-diyl group, ω position is 6-position.

10 In this specification, the single bond means that the group of interest does not exist but that the groups adjacent both sides of said group directly form a single bond. For example, to say Ar is a single bond in the group represented by the general formula (II) shows that 7-position and/or 11-position of the steroid ring in the
15 compound represented by the general formula (I) and A directly form a single bond.

20 In this specification, to say that the dashed line in combination with the solid line represents the formation of a single bond or a double bond means, for example, that the bond between 4-position and 5-position of the steroid ring denoted by the dashed line is a single bond or a double bond. This is also true with compound (2) in process A to be described later and it is meant that the bond between 5-position and 6-position of the steroid ring denoted by the
25 dashed line is a single bond or a double bond.

 In the definition of the compounds represented by the general formula (I), X^1 and X^2 represent independently a hydrogen atom or a group represented by the general formula

(II)



(wherein, in addition, Ar represents a single bond or an aromatic hydrocarbon group, A represents a methylene group or -O-, R¹ represents an optionally substituted alkyl group, an optionally substituted alkenyl group or an optionally substituted alkynyl group); preferred are the case where X¹ is -Ar-A-R¹ (wherein Ar, A and R¹ have the same meanings as defined above) and X² is a hydrogen atom, and the case where X¹ is a hydrogen atom and X² is -Ar-A-R¹ (wherein Ar, A and R¹ have the same meanings as defined above). Further preferred are compounds in which the steric configuration of X¹ in 11-position of the steroid ring is β configuration and those in which the steric configuration of X² in 7-position is α configuration. Note that X¹ and X² are not a hydrogen atom at the same time.

While R^a represents a hydrogen atom or a protective group of a hydroxyl group, a hydrogen atom is preferred. Protective groups of a hydroxyl group include acyl groups such as formyl group, acetyl group, propionyl group, butyryl group, isobutyryl group, valeryl group, isovaleryl group, pivaloyl group, caproyl group, trifluoroacetyl group and benzoyl group, alkoxycarbonyl groups such as methoxycarbonyl group, ethoxycarbonyl group, propoxycarbonyl group, isopropoxycarbonyl group, allyloxycarbonyl group, benzyloxycarbonyl group and phenoxycarbonyl group, substituted silyl groups such as trimethylsilyl group, triethylsilyl group,

triisopropylsilyl group, dimethylisopropylsilyl group,
 diethylisopropylsilyl group, dimethyltexylsilyl group, t-
 butyldimethylsilyl group, t-butyldiphenylsilyl group,
 tribenzylsilyl group, tri-p-xylylsilyl group,
 5 triphenylsilyl group, diphenylmethylsilyl group and t-
 butylmethoxyphenylsilyl group, substituted methyl groups
 such as methoxymethyl group, methoxyethoxymethyl group,
 methylthiomethyl group, t-butylthiomethyl group, β -
 trichloroethyloxymethyl group, trimethylsilylethoxymethyl
 10 group, p-methoxybenzyloxymethyl group and p-
 chlorobenzyloxymethyl group, 2-oxacycloalkyl groups such as
 tetrahydrofurallyl and tetrahydropyranyl, and aralkyl
 groups such as benzyl group. Among these, substituted
 silyl groups such as trimethylsilyl group, triethylsilyl
 15 group, triisopropylsilyl group, dimethylisopropylsilyl
 group, diethylisopropylsilyl group, dimethyltexylsilyl
 group, t-butyldimethylsilyl group, t-butyldiphenylsilyl
 group, tribenzylsilyl group, tri-p-xylylsilyl group,
 triphenylsilyl group, diphenylmethylsilyl group and t-
 20 butylmethoxyphenylsilyl group, as well as substituted
 methyl groups such as methoxymethyl group,
 methoxyethoxymethyl group, methylthiomethyl group, t-
 butylthiomethyl group, β -trichloroethyloxymethyl group,
 trimethylsilylethoxymethyl group, p-methoxybenzyloxymethyl
 25 group and p-chlorobenzyloxymethyl group are preferred, and
 t-butyldimethylsilyl group and methoxymethyl group are
 particularly preferred.

R^b and R^c , when taken together with the carbon atom in

3-position to which they are bound, represent an optionally protected $-(C=O)-$ and they preferably represent $-(C=O)-$. Examples of protected $-(C=O)-$ include noncyclic acetals or ketals such as dimethoxystyrene, bis(2,2,2-

5 trichloroethyloxy)methylene, dibenzylmethylene, bis(2-nitrobenzyloxy)methylene, bis(acetyloxy)methylene, bis(methylthio)methylene, bis(ethylthio)methylene, bis(propylthio)methylene, bis(butylthio)methylene, bis(phenylthio)methylene, bis(benzylthio)methylene,

10 bis(acetylthio)methylene, trimethylsilyloxymethylthiomethylene, trimethylsilyloxyethylthiomethylene, trimethylsilyloxyphenylthiomethylene, methyloxymethylthiomethylene, methyloxyphenylthiomethylene,

15 methyloxy-2-(methylthio)ethylthiomethylene, bis(methylselenenyl)methylene and bis(phenylselenenyl)methylene, and cyclic acetals or ketals such as 1,3-dioxane, 5,5-dibromo-1,3-dioxane, 5-(2-pyridyl)-1,3-dioxane, 1,3-dioxolane, 4-bromomethyl-1,3-

20 dioxolane, 4-(3-butenyl)-1,3-dioxolane, 4-phenyl-1,3-dioxolane, 4-(2-nitrophenyl)-1,3-dioxolane, 4,5-dimethoxymethyl-1,3-dioxolane, 1,5-dihydro-3H-2,4-benzodioxepin, 1,3-dithian, 1,3-dithiolan, 1,5-dihydro-3H-2,4-benzodithiepin and 1,3-oxathiolan; preferred are 1,3-

25 dioxane, 1,3-dioxolane and 1,3-dithian, etc. and particularly preferred are 1,3-dioxolane, etc.

The dashed line denotes that in combination with the solid line, it forms a single bond or a double bond; in

other words, a single bond and a double bond may be mentioned as the bond between 4-position and 5-position of the steroid ring; preferably, the formation of a single bond is meant. In the case where the dashed line forms a
 5 single bond in combination with the solid line, the hydrogen atom in 5-position of the steroid ring is preferably in α configuration.

In the group represented by the general formula (II), Ar represents a single bond or an aromatic hydrocarbon
 10 group.


If Ar is an aromatic hydrocarbon group, exemplary aromatic hydrocarbon rings include benzene ring, naphthalene ring, anthracene ring, naphthacene ring, pentacene ring, hexacene ring, phenanthrene ring,
 15 triphenylene ring, pyrene ring, chrysene ring, picene ring, perylene ring, pentaphene ring, coronene ring, heptaphene ring, pyranthrene ring and ovalene ring; the benzene ring is preferred. The aromatic hydrocarbon group as Ar means a group having one bond each in two different positions in
 20 the above-mentioned aromatic hydrocarbon rings and the p-phenylene group is preferably mentioned.

A represents a methylene group or -O- and a methylene group is preferred.

























If Ar is an aromatic hydrocarbon group, A is
 25 preferably -O-.


R^1 represents an optionally substituted alkyl group, an optionally substituted alkenyl group or an optionally substituted alkynyl group; preferably, R^1 is R^{1a}

[where R^{1a} is the general formula (III)




{wherein G represents an optionally substituted
straight-chained or branched alkylene group having 1 - 30
5 carbon atoms, an optionally substituted straight-chained or
branched alkenylene groups having 2 - 30 carbon atoms or an
optionally substituted straight-chained or branched
alkynylene group having 2 - 30 carbon atoms, E represents a
single bond or -O-, J represents a single bond, an
10 optionally substituted aromatic hydrocarbon group or an
optionally substituted heterocyclic group, Y represents a
single bond or -O-, L represents a single bond, a straight-
chained or branched alkylene group having 1 - 10 carbon
atoms, a straight-chained or branched alkenylene group
15 having 2 - 10 carbon atoms or a straight-chained or
branched alkynylene group having 2 - 10 carbon atoms, Q
represents a single bond or one group selected from among
the following formulae:





and



(where R^7 and R^8 represent independently a hydrogen atom or a straight-chained or branched lower alkyl group having 1 - 6 carbon atoms, R^9 , R^{10} and R^{11} each independently represent a hydrogen atom or a straight-chained or branched lower alkyl group having 1 - 3 carbon atoms), Z represents a

hydrogen atom, a straight-chained or branched alkyl group having 1 - 10 carbon atoms that may optionally be substituted by a halogen atom, a straight-chained or branched alkenyl group having 2 - 10 carbon atoms that may optionally be substituted by a halogen atom, a straight-chained or branched alkynyl group having 2 - 10 carbon atoms that may optionally be substituted by a halogen atom, $-O-R^d$ (where R^d represents a hydrogen atom or a protective group of a hydroxyl group), or $-COOH$), provided that when Q is Q^3 , the nitrogen atom and R^8 in Q^3 may combine with Z to form a heterocyclic group}}].

Examples of the substituent in G which G represents an optionally substituted straight-chained or branched alkylene group having 1 - 30 carbon atoms, an optionally substituted straight-chained or branched alkenylene group having 2 - 30 carbon atoms or an optionally substituted straight-chained or branched alkynylene group having 2 - 30 carbon atoms include $-(CH_2)_m-COOR^{7a}$, $-(CH_2)_p-CONR^{8a}R^{9a}$, $-NR^{8b}R^{9b}$, hydroxyl group, oxo group, etc. Here, m and p represent independently 0 or 1, R^{7a} represents a hydrogen atom or a straight-chained or branched alkyl group having 1 - 6 carbon atoms, R^{8a} , R^{9a} , R^{8b} and R^{9b} each independently represent a hydrogen atom or a straight-chained or branched alkyl group having 1 - 3 carbon atoms. The substituent is preferably absent or a hydroxyl group and its absence is particularly preferred. In the case where G is substituted, the number of substituents is from one to four, preferably

one.

If G represents an optionally substituted straight-chained or branched alkylene group having 1 - 30 carbon atoms, exemplary straight-chained or branched alkylene groups having 1 - 30 carbon atoms include straight-chained alkylene groups such as methylene group, ethane-1,2-diyl group, propane-1,3-diyl group, butane-1,4-diyl group, pentane-1,5-diyl group, hexane-1,6-diyl group, heptane-1,7-diyl group, octane-1,8-diyl group, nonane-1,9-diyl group, decane-1,10-diyl group, undecane-1,11-diyl group, dodecane-1,12-diyl group, tridecane-1,13-diyl group, tetradecane-1,14-diyl group, pentadecane-1,15-diyl group, hexadecane-1,16-diyl group, heptadecane-1,17-diyl group, octadecane-1,18-diyl group, nonadecane-1,19-diyl group, icosane-1,20-diyl group, henicosane-1,21-diyl group, docosane-1,22-diyl group, tricosane-1,23-diyl group, tetracosane-1,24-diyl group, pentacosane-1,25-diyl group, hexacosane-1,26-diyl group, heptacosane-1,27-diyl group, octacosane-1,28-diyl group, nonacosane-1,29-diyl group and triacontane-1,30-diyl group;

as well as branched alkylene groups such as 2-methylpropane-1,3-diyl group, 2-methylbutane-1,4-diyl group, 3-methylbutane-1,4-diyl group, 2,3-dimethylbutane-1,4-diyl group, 2-methylpentane-1,5-diyl group, 3-methylpentane-1,5-diyl group, 4-methylpentane-1,5-diyl group, 2,3-dimethylpentane-1,5-diyl group, 2,4-dimethylpentane-1,5-diyl group, 3,3-dimethylpentane-1,5-diyl group, 3,4-dimethylpentane-1,5-diyl group, 2,3,4-trimethylpentane-1,5-

diyl group, 3-ethylpentane-1,5-diyl group, 3-ethyl-2-
 methylpentane-1,5-diyl group, 3-ethyl-4-methylpentane-1,5-
 diyl group, 2,4-dimethyl-3-ethylpentane-1,5-diyl group, 2-
 methylhexane-1,6-diyl group, 3-methylhexane-1,6-diyl group,
 5 4-methylhexane-1,6-diyl group, 5-methylhexane-1,6-diyl
 group, 2,3-dimethylhexane-1,6-diyl group, 2,4-
 dimethylhexane-1,6-diyl group, 2,5-dimethylhexane-1,6-diyl
 group, 3,3-dimethylhexane-1,6-diyl group, 3,4-
 dimethylhexane-1,6-diyl group, 3,5-dimethylhexane-1,6-diyl
 10 group, 4,4-dimethylhexane-1,6-diyl group, 4,5-
 dimethylhexane-1,6-diyl group, 2,3,3-trimethylhexane-1,6-
 diyl group, 2,3,4-trimethylhexane-1,6-diyl group, 2,3,5-
 trimethylhexane-1,6-diyl group, 2,4,4-trimethylhexane-1,6-
 diyl group, 2,4,5-trimethylhexane-1,6-diyl group, 3,3,4-
 15 trimethylhexane-1,6-diyl group, 3,3,5-trimethylhexane-1,6-
 diyl group, 3,4,5-trimethylhexane-1,6-diyl group, 4,4,5-
 trimethylhexane-1,6-diyl group, 2,3,4,5-tetramethylhexane-
 1,6-diyl group, 3-ethylhexane-1,6-diyl group, 4-
 ethylhexane-1,6-diyl group, 3-ethyl-2-methylhexane-1,6-diyl
 20 group, 3-ethyl-4-methylhexane-1,6-diyl group, 3-ethyl-5-
 methylhexane-1,6-diyl group, 4-ethyl-2-methylhexane-1,6-
 diyl group, 4-ethyl-3-methylhexane-1,6-diyl group, 4-ethyl-
 5-methylhexane-1,6-diyl group, 2,4-dimethyl-3-ethylhexane-
 1,6-diyl group, 2,5-dimethyl-3-ethylhexane-1,6-diyl group,
 25 4,5-dimethyl-3-ethylhexane-1,6-diyl group, 2,3-dimethyl-4-
 ethylhexane-1,6-diyl group, 2,5-dimethyl-4-ethylhexane-1,6-
 diyl group, 3,5-dimethyl-4-ethylhexane-1,6-diyl group, 3,4-
 diethylhexane-1,6-diyl group;

2-methylheptane-1,7-diyl group, 3-methylheptane-1,7-diyl
 group, 4-methylheptane-1,7-diyl group, 5-methylheptane-1,7-
 diyl group, 6-methylheptane-1,7-diyl group, 2,3-
 dimethylheptane-1,7-diyl group, 2,4-dimethylheptane-1,7-
 5 diyl group, 2,5-dimethylheptane-1,7-diyl group, 2,6-
 dimethylheptane-1,7-diyl group, 3,3-dimethylheptane-1,7-
 diyl group, 3,4-dimethylheptane-1,7-diyl group, 3,5-
 dimethylheptane-1,7-diyl group, 3,6-dimethylheptane-1,7-
 diyl group, 4,4-dimethylheptane-1,7-diyl group, 4,5-
 10 dimethylheptane-1,7-diyl group, 4,6-dimethylheptane-1,7-
 diyl group, 5,5-dimethylheptane-1,7-diyl group, 5,6-
 dimethylheptane-1,7-diyl group, 2,3,3-trimethylheptane-1,7-
 diyl group, 2,3,4-trimethylheptane-1,7-diyl group, 2,3,5-
 trimethylheptane-1,7-diyl group, 2,3,6-trimethylheptane-
 15 1,7-diyl group, 2,4,4-trimethylheptane-1,7-diyl group,
 2,4,5-trimethylheptane-1,7-diyl group, 2,4,6-
 trimethylheptane-1,7-diyl group, 2,5,5-trimethylheptane-
 1,7-diyl group, 2,5,6-trimethylheptane-1,7-diyl group,
 3,3,4-trimethylheptane-1,7-diyl group, 3,3,5-
 20 trimethylheptane-1,7-diyl group, 3,3,6-trimethylheptane-
 1,7-diyl group, 3,4,4-trimethylheptane-1,7-diyl group,
 3,4,5-trimethylheptane-1,7-diyl group, 3,4,6-
 trimethylheptane-1,7-diyl group, 3,5,5-trimethylheptane-
 1,7-diyl group, 3,5,6-trimethylheptane-1,7-diyl group,
 25 4,4,5-trimethylheptane-1,7-diyl group, 4,4,6-
 trimethylheptane-1,7-diyl group, 4,5,5-trimethylheptane-
 1,7-diyl group, 4,5,6-trimethylheptane-1,7-diyl group, 3-
 ethylheptane-1,7-diyl group, 4-ethylheptane-1,7-diyl group,

- 5-ethylheptane-1,7-diyl group, 3-ethyl-2-methylheptane-1,7-diyl group, 3-ethyl-4-methylheptane-1,7-diyl group, 3-ethyl-5-methylheptane-1,7-diyl group, 3-ethyl-6-methylheptane-1,7-diyl group, 4-ethyl-2-methylheptane-1,7-diyl group, 4-ethyl-3-methylheptane-1,7-diyl group, 4-ethyl-4-methylheptane-1,7-diyl group, 4-ethyl-5-methylheptane-1,7-diyl group, 4-ethyl-6-methylheptane-1,7-diyl group, 5-ethyl-2-methylheptane-1,7-diyl group, 5-ethyl-3-methylheptane-1,7-diyl group, 5-ethyl-4-methylheptane-1,7-diyl group, 5-ethyl-5-methylheptane-1,7-diyl group, 5-ethyl-6-methylheptane-1,7-diyl group, 4-n-propylheptane-1,7-diyl group, 4-i-propylheptane-1,7-diyl group;
- 2-methyloctane-1,8-diyl group, 3-methyloctane-1,8-diyl group, 3-methyloctane-1,8-diyl group, 4-methyloctane-1,8-diyl group, 5-methyloctane-1,8-diyl group, 6-methyloctane-1,8-diyl group, 7-methyloctane-1,8-diyl group, 2,3-dimethyloctane-1,8-diyl group, 2,4-dimethyloctane-1,8-diyl group, 2,5-dimethyloctane-1,8-diyl group, 2,6-dimethyloctane-1,8-diyl group, 2,7-dimethyloctane-1,8-diyl group, 3,3-dimethyloctane-1,8-diyl group, 3,4-dimethyloctane-1,8-diyl group, 3,5-dimethyloctane-1,8-diyl group, 3,6-dimethyloctane-1,8-diyl group, 3,7-dimethyloctane-1,8-diyl group, 4,4-dimethyloctane-1,8-diyl group, 4,5-dimethyloctane-1,8-diyl group, 4,6-dimethyloctane-1,8-diyl group, 4,7-dimethyloctane-1,8-diyl group, 5,5-dimethyloctane-1,8-diyl group, 5,6-dimethyloctane-1,8-diyl group, 5,7-dimethyloctane-1,8-diyl

- group, 6,6-dimethyloctane-1,8-diyl group, 6,7-dimethyloctane-1,8-diyl group, 3-ethyloctane-1,8-diyl group, 4-ethyloctane-1,8-diyl group, 5-ethyloctane-1,8-diyl group, 6-ethyloctane-1,8-diyl group, 2-methylnonane-1,9-diyl group,
- 5 3-methylnonane-1,9-diyl group, 4-methylnonane-1,9-diyl group, 5-methylnonane-1,9-diyl group, 6-methylnonane-1,9-diyl group, 7-methylnonane-1,9-diyl group, 8-methylnonane-1,9-diyl group;
- 2-methyldecane-1,10-diyl group, 3-methyldecane-1,10-diyl group, 4-methyldecane-1,10-diyl group, 5-methyldecane-1,10-diyl group, 6-methyldecane-1,10-diyl group, 7-methyldecane-1,10-diyl group, 8-methyldecane-1,10-diyl group, 4-ethyldecane-1,10-diyl group, 5-ethyldecane-1,10-diyl group, 6-ethyldecane-1,10-diyl group, 7-ethyldecane-1,10-diyl group,
- 15 5-n-propyldecane-1,10-diyl group, 6-n-propyldecane-1,10-diyl group, 3-ethyl-2-methyldecane-1,10-diyl group, 4-ethyl-2-methyldecane-1,10-diyl group, 5-ethyl-2-methyldecane-1,10-diyl group, 6-ethyl-2-methyldecane-1,10-diyl group, 7-ethyl-2-methyldecane-1,10-diyl group, 3-ethyl-3-methyldecane-1,10-diyl group, 4-ethyl-3-methyldecane-1,10-diyl group, 5-ethyl-3-methyldecane-1,10-diyl group, 6-ethyl-3-methyldecane-1,10-diyl group, 7-ethyl-3-methyldecane-1,10-diyl group, 3-ethyl-4-methyldecane-1,10-diyl group, 4-ethyl-4-methyldecane-1,10-diyl group, 5-ethyl-4-methyldecane-1,10-diyl group, 6-ethyl-4-methyldecane-1,10-diyl group, 7-ethyl-4-methyldecane-1,10-diyl group, 3-ethyl-5-methyldecane-1,10-diyl group, 4-ethyl-5-methyldecane-1,10-diyl group, 5-
- 25

ethyl-5-methyldecane-1,10-diyl group, 6-ethyl-5-methyl-
decane-1,10-diyl group, 7-ethyl-5-methyldecane-1,10-diyl
group;

2-methylundecane-1,11-diyl group, 3-methylundecane-1,11-
5 diyl group, 4-methylundecane-1,11-diyl group, 5-
methylundecane-1,11-diyl group, 6-methylundecane-1,11-diyl
group, 7-methylundecane-1,11-diyl group, 8-methylundecane-
1,11-diyl group, 9-methylundecane-1,11-diyl group, 10-
methylundecane-1,11-diyl group, 3-ethylundecane-1,11-diyl
10 group, 4-ethylundecane-1,11-diyl group, 5-ethylundecane-
1,11-diyl group, 6-ethylundecane-1,11-diyl group, 7-
ethylundecane-1,11-diyl group, 8-ethylundecane-1,11-diyl
group, 9-ethylundecane-1,11-diyl group;

2-methyldodecane-1,12-diyl group, 3-methyldodecane-1,12-
15 diyl group, 4-methyldodecane-1,12-diyl group, 5-
methyldodecane-1,12-diyl group, 6-methyldodecane-1,12-diyl
group, 7-methyldodecane-1,12-diyl group, 8-methyldodecane-
1,12-diyl group, 9-methyldodecane-1,12-diyl group, 10-
methyldodecane-1,12-diyl group, 11-methyldodecane-1,12-diyl
20 group;

3-ethyldodecane-1,12-diyl group, 4-ethyldodecane-1,12-diyl
group, 5-ethyldodecane-1,12-diyl group, 6-ethyldodecane-
1,12-diyl group, 7-ethyldodecane-1,12-diyl group, 8-
ethyldodecane-1,12-diyl group, 9-ethyldodecane-1,12-diyl
25 group, 10-ethyldodecane-1,12-diyl group;

2-methyltridecane-1,13-diyl group, 3-methyltridecane-1,13-
diyl group, 4-methyltridecane-1,13-diyl group, 5-
methyltridecane-1,13-diyl group, 6-methyltridecane-1,13-

- diyl group, 7-methyltridecane-1,13-diyl group, 8-methyltridecane-1,13-diyl group, 9-methyltridecane-1,13-diyl group, 10-methyltridecane-1,13-diyl group, 11-methyltridecane-1,13-diyl group, 12-methyltridecane-1,13-
- 5 diyl group;
- 3-ethyltridecane-1,13-diyl group, 4-ethyltridecane-1,13-diyl group, 5-ethyltridecane-1,13-diyl group, 6-ethyltridecane-1,13-diyl group, 7-ethyltridecane-1,13-diyl group, 8-ethyltridecane-1,13-diyl group, 9-ethyltridecane-
- 10 1,13-diyl group, 10-ethyltridecane-1,13-diyl group, 11-ethyltridecane-1,13-diyl group;
- 2-methyltetradecane-1,14-diyl group, 3-methyltetradecane-1,14-diyl group, 4-methyltetradecane-1,14-diyl group, 5-methyltetradecane-1,14-diyl group, 6-methyltetradecane-
- 15 1,14-diyl group, 7-methyltetradecane-1,14-diyl group, 8-methyltetradecane-1,14-diyl group, 9-methyltetradecane-1,14-diyl group, 10-methyltetradecane-1,14-diyl group, 11-methyltetradecane-1,14-diyl group, 12-methyltetradecane-1,14-diyl group, 13-methyltetradecane-1,14-diyl group;
- 20 3-ethyltetradecane-1,14-diyl group, 4-ethyltetradecane-1,14-diyl group, 5-ethyltetradecane-1,14-diyl group, 6-ethyltetradecane-1,14-diyl group, 7-ethyltetradecane-1,14-diyl group, 8-ethyltetradecane-1,14-diyl group, 9-ethyltetradecane-1,14-diyl group, 10-ethyltetradecane-1,14-
- 25 diyl group, 11-ethyltetradecane-1,14-diyl group, 12-ethyltetradecane-1,14-diyl group;
- 2-methylpentadecane-1,15-diyl group, 3-methylpentadecane-1,15-diyl group, 4-methylpentadecane-1,15-diyl group, 5-

- methylpentadecane-1,15-diyl group, 6-methylpentadecane-
 1,15-diyl group, 7-methylpentadecane-1,15-diyl group, 8-
 methylpentadecane-1,15-diyl group, 9-methylpentadecane-
 1,15-diyl group, 10-methylpentadecane-1,15-diyl group, 11-
 5 methylpentadecane-1,15-diyl group, 12-methylpentadecane-
 1,15-diyl group, 13-methylpentadecane-1,15-diyl group, 14-
 methylpentadecane-1,15-diyl group;
 3-ethylpentadecane-1,15-diyl group, 4-ethylpentadecane-
 1,15-diyl group, 5-ethylpentadecane-1,15-diyl group, 6-
 10 ethylpentadecane-1,15-diyl group, 7-ethylpentadecane-1,15-
 diyl group, 8-ethylpentadecane-1,15-diyl group, 9-
 ethylpentadecane-1,15-diyl group, 10-ethylpentadecane-1,15-
 diyl group, 11-ethylpentadecane-1,15-diyl group, 12-
 ethylpentadecane-1,15-diyl group, 13-ethylpentadecane-1,15-
 15 diyl group;
 2-methylhexadecane-1,16-diyl group, 3-methylhexadecane-
 1,16-diyl group, 4-methylhexadecane-1,16-diyl group, 5-
 methylhexadecane-1,16-diyl group, 6-methylhexadecane-1,16-
 diyl group, 7-methylhexadecane-1,16-diyl group, 8-
 20 methylhexadecane-1,16-diyl group, 9-methylhexadecane-1,16-
 diyl group, 10-methylhexadecane-1,16-diyl group, 11-
 methylhexadecane-1,16-diyl group, 12-methylhexadecane-1,16-
 diyl group, 13-methylhexadecane-1,16-diyl group, 14-
 methylhexadecane-1,16-diyl group, 15-methylhexadecane-1,16-
 25 diyl group;
 3-ethylhexadecane-1,16-diyl group, 4-ethylhexadecane-1,16-
 diyl group, 5-ethylhexadecane-1,16-diyl group, 6-
 ethylhexadecane-1,16-diyl group, 7-ethylhexadecane-1,16-

- diyl group, 8-ethylhexadecane-1,16-diyl group, 9-ethylhexadecane-1,16-diyl group, 10-ethylhexadecane-1,16-diyl group, 11-ethylhexadecane-1,16-diyl group, 12-ethylhexadecane-1,16-diyl group, 13-ethylhexadecane-1,16-diyl group, 14-ethylhexadecane-1,16-diyl group;
- 2-methylheptadecane-1,17-diyl group, 3-methylheptadecane-1,17-diyl group, 4-methylheptadecane-1,17-diyl group, 5-methylheptadecane-1,17-diyl group, 6-methylheptadecane-1,17-diyl group, 7-methylheptadecane-1,17-diyl group, 8-methylheptadecane-1,17-diyl group, 9-methylheptadecane-1,17-diyl group, 10-methylheptadecane-1,17-diyl group, 11-methylheptadecane-1,17-diyl group, 12-methylheptadecane-1,17-diyl group, 13-methylheptadecane-1,17-diyl group, 14-methylheptadecane-1,17-diyl group, 15-methylheptadecane-1,17-diyl group, 16-methylheptadecane-1,17-diyl group;
- 3-ethylheptadecane-1,17-diyl group, 4-ethylheptadecane-1,17-diyl group, 5-ethylheptadecane-1,17-diyl group, 6-ethylheptadecane-1,17-diyl group, 7-ethylheptadecane-1,17-diyl group, 8-ethylheptadecane-1,17-diyl group, 9-ethylheptadecane-1,17-diyl group, 10-ethylheptadecane-1,17-diyl group, 11-ethylheptadecane-1,17-diyl group, 12-ethylheptadecane-1,17-diyl group, 13-ethylheptadecane-1,17-diyl group, 14-ethylheptadecane-1,17-diyl group, 15-ethylheptadecane-1,17-diyl group;
- 2-methyloctadecane-1,18-diyl group, 3-methyloctadecane-1,18-diyl group, 4-methyloctadecane-1,18-diyl group, 5-methyloctadecane-1,18-diyl group, 6-methyloctadecane-1,18-diyl group, 7-methyloctadecane-1,18-diyl group, 8-

- methyloctadecane-1,18-diyl group, 9-methyloctadecane-1,18-diyl group, 10-methyloctadecane-1,18-diyl group, 11-methyloctadecane-1,18-diyl group, 12-methyloctadecane-1,18-diyl group, 13-methyloctadecane-1,18-diyl group, 14-methyloctadecane-1,18-diyl group, 15-methyloctadecane-1,18-diyl group, 16-methyloctadecane-1,18-diyl group, 17-methyloctadecane-1,18-diyl group;
- 3-ethyloctadecane-1,18-diyl group, 4-ethyloctadecane-1,18-diyl group, 5-ethyloctadecane-1,18-diyl group, 6-ethyloctadecane-1,18-diyl group, 7-ethyloctadecane-1,18-diyl group, 8-ethyloctadecane-1,18-diyl group, 9-ethyloctadecane-1,18-diyl group, 10-ethyloctadecane-1,18-diyl group, 11-ethyloctadecane-1,18-diyl group, 12-ethyloctadecane-1,18-diyl group, 13-ethyloctadecane-1,18-diyl group, 14-ethyloctadecane-1,18-diyl group, 15-ethyloctadecane-1,18-diyl group, 16-ethyloctadecane-1,18-diyl group;
- 2-methylnonadecane-1,19-diyl group, 3-methylnonadecane-1,19-diyl group, 4-methylnonadecane-1,19-diyl group, 5-methylnonadecane-1,19-diyl group, 6-methylnonadecane-1,19-diyl group, 7-methylnonadecane-1,19-diyl group, 8-methylnonadecane-1,19-diyl group, 9-methylnonadecane-1,19-diyl group, 10-methylnonadecane-1,19-diyl group, 11-methylnonadecane-1,19-diyl group, 12-methylnonadecane-1,19-diyl group, 13-methylnonadecane-1,19-diyl group, 14-methylnonadecane-1,19-diyl group, 15-methylnonadecane-1,19-diyl group, 16-methylnonadecane-1,19-diyl group, 17-methylnonadecane-1,19-diyl group, 18-methylnonadecane-1,19-diyl group;

diyl group;

3-ethylnonadecane-1,19-diyl group, 4-ethylnonadecane-1,19-diyl group, 5-ethylnonadecane-1,19-diyl group, 6-ethylnonadecane-1,19-diyl group, 7-ethylnonadecane-1,19-diyl group, 8-ethylnonadecane-1,19-diyl group, 9-ethylnonadecane-1,19-diyl group, 10-ethylnonadecane-1,19-diyl group, 11-ethylnonadecane-1,19-diyl group, 12-ethylnonadecane-1,19-diyl group, 13-ethylnonadecane-1,19-diyl group, 14-ethylnonadecane-1,19-diyl group, 15-ethylnonadecane-1,19-diyl group, 16-ethylnonadecane-1,19-diyl group, 17-ethylnonadecane-1,19-diyl group;

2-methylicosane-1,20-diyl group, 3-methylicosane-1,20-diyl group, 4-methylicosane-1,20-diyl group, 5-methylicosane-1,20-diyl group, 6-methylicosane-1,20-diyl group, 7-methylicosane-1,20-diyl group, 8-methylicosane-1,20-diyl group, 9-methylicosane-1,20-diyl group, 10-methylicosane-1,20-diyl group, 11-methylicosane-1,20-diyl group, 12-methylicosane-1,20-diyl group, 13-methylicosane-1,20-diyl group, 14-methylicosane-1,20-diyl group, 15-methylicosane-1,20-diyl group, 16-methylicosane-1,20-diyl group, 17-methylicosane-1,20-diyl group, 18-methylicosane-1,20-diyl group, 19-methylicosane-1,20-diyl group;

3-ethylicosane-1,20-diyl group, 4-ethylicosane-1,20-diyl group, 5-ethylicosane-1,20-diyl group, 6-ethylicosane-1,20-diyl group, 7-ethylicosane-1,20-diyl group, 8-ethylicosane-1,20-diyl group, 9-ethylicosane-1,20-diyl group, 10-ethylicosane-1,20-diyl group, 11-ethylicosane-1,20-diyl group, 12-ethylicosane-1,20-diyl group, 13-ethylicosane-

- 1,20-diyl group, 14-ethylicosane-1,20-diyl group, 15-ethylicosane-1,20-diyl group, 16-ethylicosane-1,20-diyl group, 17-ethylicosane-1,20-diyl group, 18-ethylicosane-1,20-diyl group;
- 5 2-methylhenicosane-1,21-diyl group, 3-methylhenicosane-1,21-diyl group, 4-methylhenicosane-1,21-diyl group, 5-methylhenicosane-1,21-diyl group, 6-methylhenicosane-1,21-diyl group, 7-methylhenicosane-1,21-diyl group, 8-methylhenicosane-1,21-diyl group, 9-methylhenicosane-1,21-
- 10 diyl group, 10-methylhenicosane-1,21-diyl group, 11-methylhenicosane-1,21-diyl group, 12-methylhenicosane-1,21-diyl group, 13-methylhenicosane-1,21-diyl group, 14-methylhenicosane-1,21-diyl group, 15-methylhenicosane-1,21-diyl group, 16-methylhenicosane-1,21-diyl group, 17-
- 15 methylhenicosane-1,21-diyl group, 18-methylhenicosane-1,21-diyl group, 19-methylhenicosane-1,21-diyl group, 20-methylhenicosane-1,21-diyl group;
- 3-ethylhenicosane-1,21-diyl group, 4-ethylhenicosane-1,21-diyl group, 5-ethylhenicosane-1,21-diyl group, 6-
- 20 ethylhenicosane-1,21-diyl group, 7-ethylhenicosane-1,21-diyl group, 8-ethylhenicosane-1,21-diyl group, 9-ethylhenicosane-1,21-diyl group, 10-ethylhenicosane-1,21-diyl group, 11-ethylhenicosane-1,21-diyl group, 12-ethylhenicosane-1,21-diyl group, 13-ethylhenicosane-1,21-
- 25 diyl group, 14-ethylhenicosane-1,21-diyl group, 15-ethylhenicosane-1,21-diyl group, 16-ethylhenicosane-1,21-diyl group, 17-ethylhenicosane-1,21-diyl group, 18-ethylhenicosane-1,21-diyl group, 19-ethylhenicosane-1,21-

- diyl group;
- 2-methyldocosane-1,22-diyl group, 3-methyldocosane-1,22-diyl group, 4-methyldocosane-1,22-diyl group, 5-methyldocosane-1,22-diyl group, 6-methyldocosane-1,22-diyl group, 7-methyldocosane-1,22-diyl group, 8-methyldocosane-1,22-diyl group, 9-methyldocosane-1,22-diyl group, 10-methyldocosane-1,22-diyl group, 11-methyldocosane-1,22-diyl group, 12-methyldocosane-1,22-diyl group, 13-methyldocosane-1,22-diyl group, 14-methyldocosane-1,22-diyl group, 15-methyldocosane-1,22-diyl group, 16-methyldocosane-1,22-diyl group, 17-methyldocosane-1,22-diyl group, 18-methyldocosane-1,22-diyl group, 19-methyldocosane-1,22-diyl group, 20-methyldocosane-1,22-diyl group, 21-methyldocosane-1,22-diyl group;
- 15 3-ethyldocosane-1,22-diyl group, 4-ethyldocosane-1,22-diyl group, 5-ethyldocosane-1,22-diyl group, 6-ethyldocosane-1,22-diyl group, 7-ethyldocosane-1,22-diyl group, 8-ethyldocosane-1,22-diyl group, 9-ethyldocosane-1,22-diyl group, 10-ethyldocosane-1,22-diyl group, 11-ethyldocosane-1,22-diyl group, 12-ethyldocosane-1,22-diyl group, 13-ethyldocosane-1,22-diyl group, 14-ethyldocosane-1,22-diyl group, 15-ethyldocosane-1,22-diyl group, 16-ethyldocosane-1,22-diyl group, 17-ethyldocosane-1,22-diyl group, 18-ethyldocosane-1,22-diyl group, 19-ethyldocosane-1,22-diyl group, 20-ethyldocosane-1,22-diyl group;
- 25 2-methyltricosane-1,23-diyl group, 3-methyltricosane-1,23-diyl group, 4-methyltricosane-1,23-diyl group, 5-methyltricosane-1,23-diyl group, 6-methyltricosane-1,23-

- diyl group, 7-methyltricosane-1,23-diyl group, 8-methyltricosane-1,23-diyl group, 9-methyltricosane-1,23-diyl group, 10-methyltricosane-1,23-diyl group, 11-methyltricosane-1,23-diyl group, 12-methyltricosane-1,23-diyl group, 13-methyltricosane-1,23-diyl group, 14-methyltricosane-1,23-diyl group, 15-methyltricosane-1,23-diyl group, 16-methyltricosane-1,23-diyl group, 17-methyltricosane-1,23-diyl group, 18-methyltricosane-1,23-diyl group, 19-methyltricosane-1,23-diyl group, 20-methyltricosane-1,23-diyl group, 21-methyltricosane-1,23-diyl group, 22-methyltricosane-1,23-diyl group; 3-ethyltricosane-1,23-diyl group, 4-ethyltricosane-1,23-diyl group, 5-ethyltricosane-1,23-diyl group, 6-ethyltricosane-1,23-diyl group, 7-ethyltricosane-1,23-diyl group, 8-ethyltricosane-1,23-diyl group, 9-ethyltricosane-1,23-diyl group, 10-ethyltricosane-1,23-diyl group, 11-ethyltricosane-1,23-diyl group, 12-ethyltricosane-1,23-diyl group, 13-ethyltricosane-1,23-diyl group, 14-ethyltricosane-1,23-diyl group, 15-ethyltricosane-1,23-diyl group, 16-ethyltricosane-1,23-diyl group, 17-ethyltricosane-1,23-diyl group, 18-ethyltricosane-1,23-diyl group, 19-ethyltricosane-1,23-diyl group, 20-ethyltricosane-1,23-diyl group, 21-ethyltricosane-1,23-diyl group;
- 2-methyltetracosane-1,24-diyl group, 3-methyltetracosane-1,24-diyl group, 4-methyltetracosane-1,24-diyl group, 5-methyltetracosane-1,24-diyl group, 6-methyltetracosane-1,24-diyl group, 7-methyltetracosane-1,24-diyl group, 8-

methyltetracosane-1,24-diyl group, 9-methyltetracosane-
 1,24-diyl group, 10-methyltetracosane-1,24-diyl group, 11-
 methyltetracosane-1,24-diyl group, 12-methyltetracosane-
 1,24-diyl group, 13-methyltetracosane-1,24-diyl group, 14-
 5 methyltetracosane-1,24-diyl group, 15-methyltetracosane-
 1,24-diyl group, 16-methyltetracosane-1,24-diyl group, 17-
 methyltetracosane-1,24-diyl group, 18-methyltetracosane-
 1,24-diyl group, 19-methyltetracosane-1,24-diyl group, 20-
 methyltetracosane-1,24-diyl group, 21-methyltetracosane-
 10 1,24-diyl group, 22-methyltetracosane-1,24-diyl group, 23-
 methyltetracosane-1,24-diyl group;
 3-ethyltetracosane-1,24-diyl group, 4-ethyltetracosane-
 1,24-diyl group, 5-ethyltetracosane-1,24-diyl group, 6-
 ethyltetracosane-1,24-diyl group, 7-ethyltetracosane-1,24-
 15 diyl group, 8-ethyltetracosane-1,24-diyl group, 9-
 ethyltetracosane-1,24-diyl group, 10-ethyltetracosane-1,24-
 diyl group, 11-ethyltetracosane-1,24-diyl group, 12-
 ethyltetracosane-1,24-diyl group, 13-ethyltetracosane-1,24-
 diyl group, 14-ethyltetracosane-1,24-diyl group, 15-
 20 ethyltetracosane-1,24-diyl group, 16-ethyltetracosane-1,24-
 diyl group, 17-ethyltetracosane-1,24-diyl group, 18-
 ethyltetracosane-1,24-diyl group, 19-ethyltetracosane-1,24-
 diyl group, 20-ethyltetracosane-1,24-diyl group, 21-
 ethyltetracosane-1,24-diyl group, 22-ethyltetracosane-1,24-
 25 diyl group;
 2-methylpentacosane-1,25-diyl group, 3-methylpentacosane-
 1,25-diyl group, 4-methylpentacosane-1,25-diyl group, 5-
 methylpentacosane-1,25-diyl group, 6-methylpentacosane-

- 1,25-diyl group, 7-methylpentacosane-1,25-diyl group, 8-methylpentacosane-1,25-diyl group, 9-methylpentacosane-1,25-diyl group, 10-methylpentacosane-1,25-diyl group, 11-methylpentacosane-1,25-diyl group, 12-methylpentacosane-1,25-diyl group, 13-methylpentacosane-1,25-diyl group, 14-methylpentacosane-1,25-diyl group, 15-methylpentacosane-1,25-diyl group, 16-methylpentacosane-1,25-diyl group, 17-methylpentacosane-1,25-diyl group, 18-methylpentacosane-1,25-diyl group, 19-methylpentacosane-1,25-diyl group, 20-methylpentacosane-1,25-diyl group, 21-methylpentacosane-1,25-diyl group, 22-methylpentacosane-1,25-diyl group, 23-methylpentacosane-1,25-diyl group, 24-methylpentacosane-1,25-diyl group;
- 3-ethylpentacosane-1,25-diyl group, 4-ethylpentacosane-1,25-diyl group, 5-ethylpentacosane-1,25-diyl group, 6-ethylpentacosane-1,25-diyl group, 7-ethylpentacosane-1,25-diyl group, 8-ethylpentacosane-1,25-diyl group, 9-ethylpentacosane-1,25-diyl group, 10-ethylpentacosane-1,25-diyl group, 11-ethylpentacosane-1,25-diyl group, 12-ethylpentacosane-1,25-diyl group, 13-ethylpentacosane-1,25-diyl group, 14-ethylpentacosane-1,25-diyl group, 15-ethylpentacosane-1,25-diyl group, 16-ethylpentacosane-1,25-diyl group, 17-ethylpentacosane-1,25-diyl group, 18-ethylpentacosane-1,25-diyl group, 19-ethylpentacosane-1,25-diyl group, 20-ethylpentacosane-1,25-diyl group, 21-ethylpentacosane-1,25-diyl group, 22-ethylpentacosane-1,25-diyl group, 23-ethylpentacosane-1,25-diyl group;
- 2-methylhexacosane-1,26-diyl group, 3-methylhexacosane-

- 1,26-diyl group, 4-methylhexacosane-1,26-diyl group, 5-methylhexacosane-1,26-diyl group, 6-methylhexacosane-1,26-diyl group, 7-methylhexacosane-1,26-diyl group, 8-methylhexacosane-1,26-diyl group, 9-methylhexacosane-1,26-diyl group, 10-methylhexacosane-1,26-diyl group, 11-methylhexacosane-1,26-diyl group, 12-methylhexacosane-1,26-diyl group, 13-methylhexacosane-1,26-diyl group, 14-methylhexacosane-1,26-diyl group, 15-methylhexacosane-1,26-diyl group, 16-methylhexacosane-1,26-diyl group, 17-methylhexacosane-1,26-diyl group, 18-methylhexacosane-1,26-diyl group, 19-methylhexacosane-1,26-diyl group, 20-methylhexacosane-1,26-diyl group, 21-methylhexacosane-1,26-diyl group, 22-methylhexacosane-1,26-diyl group, 23-methylhexacosane-1,26-diyl group, 24-methylhexacosane-1,26-diyl group, 25-methylhexacosane-1,26-diyl group;
- 3-ethylhexacosane-1,26-diyl group, 4-ethylhexacosane-1,26-diyl group, 5-ethylhexacosane-1,26-diyl group, 6-ethylhexacosane-1,26-diyl group, 7-ethylhexacosane-1,26-diyl group, 8-ethylhexacosane-1,26-diyl group, 9-ethylhexacosane-1,26-diyl group, 10-ethylhexacosane-1,26-diyl group, 11-ethylhexacosane-1,26-diyl group, 12-ethylhexacosane-1,26-diyl group, 13-ethylhexacosane-1,26-diyl group, 14-ethylhexacosane-1,26-diyl group, 15-ethylhexacosane-1,26-diyl group, 16-ethylhexacosane-1,26-diyl group, 17-ethylhexacosane-1,26-diyl group, 18-ethylhexacosane-1,26-diyl group, 19-ethylhexacosane-1,26-diyl group, 20-ethylhexacosane-1,26-diyl group, 21-ethylhexacosane-1,26-diyl group, 22-ethylhexacosane-1,26-

- diyl group, 23-ethylhexacosane-1,26-diyl group, 24-ethylhexacosane-1,26-diyl group;
- 2-methylheptacosane-1,27-diyl group, 3-methylheptacosane-1,27-diyl group, 4-methylheptacosane-1,27-diyl group, 5-
5. methylheptacosane-1,27-diyl group, 6-methylheptacosane-1,27-diyl group, 7-methylheptacosane-1,27-diyl group, 8-methylheptacosane-1,27-diyl group, 9-methylheptacosane-1,27-diyl group, 10-methylheptacosane-1,27-diyl group, 11-methylheptacosane-1,27-diyl group, 12-methylheptacosane-
- 10 1,27-diyl group, 13-methylheptacosane-1,27-diyl group, 14-methylheptacosane-1,27-diyl group, 15-methylheptacosane-1,27-diyl group, 16-methylheptacosane-1,27-diyl group, 17-methylheptacosane-1,27-diyl group, 18-methylheptacosane-1,27-diyl group, 19-methylheptacosane-1,27-diyl group, 20-
- 15 methylheptacosane-1,27-diyl group, 21-methylheptacosane-1,27-diyl group, 22-methylheptacosane-1,27-diyl group, 23-methylheptacosane-1,27-diyl group, 24-methylheptacosane-1,27-diyl group, 25-methylheptacosane-1,27-diyl group, 26-methylheptacosane-1,27-diyl group;
- 20 3-ethylheptacosane-1,27-diyl group, 4-ethylheptacosane-1,27-diyl group, 5-ethylheptacosane-1,27-diyl group, 6-ethylheptacosane-1,27-diyl group, 7-ethylheptacosane-1,27-diyl group, 8-ethylheptacosane-1,27-diyl group, 9-ethylheptacosane-1,27-diyl group, 10-ethylheptacosane-1,27-
- 25 diyl group, 11-ethylheptacosane-1,27-diyl group, 12-ethylheptacosane-1,27-diyl group, 13-ethylheptacosane-1,27-diyl group, 14-ethylheptacosane-1,27-diyl group, 15-ethylheptacosane-1,27-diyl group, 16-ethylheptacosane-1,27-

diyl group, 17-ethylheptacosane-1,27-diyl group, 18-
 ethylheptacosane-1,27-diyl group, 19-ethylheptacosane-1,27-
 diyl group, 20-ethylheptacosane-1,27-diyl group, 21-
 ethylheptacosane-1,27-diyl group, 22-ethylheptacosane-1,27-
 5 diyl group, 23-ethylheptacosane-1,27-diyl group, 24-
 ethylheptacosane-1,27-diyl group, 25-ethylheptacosane-1,27-
 diyl group;
 2-methyloctacosane-1,28-diyl group, 3-methyloctacosane-
 1,28-diyl group, 4-methyloctacosane-1,28-diyl group, 5-
 10 methyloctacosane-1,28-diyl group, 6-methyloctacosane-1,28-
 diyl group, 7-methyloctacosane-1,28-diyl group, 8-
 methyloctacosane-1,28-diyl group, 9-methyloctacosane-1,28-
 diyl group, 10-methyloctacosane-1,28-diyl group, 11-
 methyloctacosane-1,28-diyl group, 12-methyloctacosane-1,28-
 15 diyl group, 13-methyloctacosane-1,28-diyl group, 14-
 methyloctacosane-1,28-diyl group, 15-methyloctacosane-1,28-
 diyl group, 16-methyloctacosane-1,28-diyl group, 17-
 methyloctacosane-1,28-diyl group, 18-methyloctacosane-1,28-
 diyl group, 19-methyloctacosane-1,28-diyl group, 20-
 20 methyloctacosane-1,28-diyl group, 21-methyloctacosane-1,28-
 diyl group, 22-methyloctacosane-1,28-diyl group, 23-
 methyloctacosane-1,28-diyl group, 24-methyloctacosane-1,28-
 diyl group, 25-methyloctacosane-1,28-diyl group, 26-
 methyloctacosane-1,28-diyl group, 27-methyloctacosane-1,28-
 25 diyl group;
 3-ethyloctacosane-1,28-diyl group, 4-ethyloctacosane-1,28-
 diyl group, 5-ethyloctacosane-1,28-diyl group, 6-
 ethyloctacosane-1,28-diyl group, 7-ethyloctacosane-1,28-

diyl group, 8-ethyloctacosane-1,28-diyl group, 9-
 ethyloctacosane-1,28-diyl group, 10-ethyloctacosane-1,28-
 diyl group, 11-ethyloctacosane-1,28-diyl group, 12-
 ethyloctacosane-1,28-diyl group, 13-ethyloctacosane-1,28-
 5 diyl group, 14-ethyloctacosane-1,28-diyl group, 15-
 ethyloctacosane-1,28-diyl group, 16-ethyloctacosane-1,28-
 diyl group, 17-ethyloctacosane-1,28-diyl group, 18-
 ethyloctacosane-1,28-diyl group, 19-ethyloctacosane-1,28-
 diyl group, 20-ethyloctacosane-1,28-diyl group, 21-
 10 ethyloctacosane-1,28-diyl group, 22-ethyloctacosane-1,28-
 diyl group, 23-ethyloctacosane-1,28-diyl group, 24-
 ethyloctacosane-1,28-diyl group, 25-ethyloctacosane-1,28-
 diyl group, 26-ethyloctacosane-1,28-diyl group;
 2-methylnonacosane-1,29-diyl group, 3-methylnonacosane-
 15 1,29-diyl group, 4-methylnonacosane-1,29-diyl group, 5-
 methylnonacosane-1,29-diyl group, 6-methylnonacosane-1,29-
 diyl group, 7-methylnonacosane-1,29-diyl group, 8-
 methylnonacosane-1,29-diyl group, 9-methylnonacosane-1,29-
 diyl group, 10-methylnonacosane-1,29-diyl group, 11-
 20 methylnonacosane-1,29-diyl group, 12-methylnonacosane-1,29-
 diyl group, 13-methylnonacosane-1,29-diyl group, 14-
 methylnonacosane-1,29-diyl group, 15-methylnonacosane-1,29-
 diyl group, 16-methylnonacosane-1,29-diyl group, 17-
 methylnonacosane-1,29-diyl group, 18-methylnonacosane-1,29-
 25 diyl group, 19-methylnonacosane-1,29-diyl group, 20-
 methylnonacosane-1,29-diyl group, 21-methylnonacosane-1,29-
 diyl group, 22-methylnonacosane-1,29-diyl group, 23-
 methylnonacosane-1,29-diyl group, 24-methylnonacosane-1,29-

diyl group, 25-methylnonacosane-1,29-diyl group, 26-methylnonacosane-1,29-diyl group, 27-methylnonacosane-1,29-diyl group and 28-methylnonacosane-1,29-diyl group.

If G represents an optionally substituted straight-
 5 chained or branched alkenylene group having 2 - 30 carbon atoms, exemplary straight-chained or branched alkenylene groups having 2 - 30 carbon atoms include straight-chained alkenylene groups such as ethylene-1,2-diyl group, 1-propene-1,3-diyl group, 2-propene-1,3-diyl group, 1-butene-
 10 1,4-diyl group, 2-butene-1,4-diyl group, 3-butene-1,4-diyl group, 1,3-butadiene-1,4-diyl group, 2-pentene-1,5-diyl group, 3-pentene-1,5-diyl group, 2,4-pentadiene-1,5-diyl group, 2-hexene-1,6-diyl group, 3-hexene-1,6-diyl group, 4-hexene-1,6-diyl group, 2,4-hexadiene-1,6-diyl group, 2-
 15 heptene-1,7-diyl group, 3-heptene-1,7-diyl group, 4-heptene-1,7-diyl group, 5-heptene-1,7-diyl group, 2,4-heptadiene-1,7-diyl group, 2,5-heptadiene-1,7-diyl group, 3,5-heptadiene-1,7-diyl group, 2-octene-1,8-diyl group, 3-octene-1,8-diyl group, 4-octene-1,8-diyl group, 5-octene-
 20 1,8-diyl group, 6-octene-1,8-diyl group, 2,4-octadiene-1,8-diyl group, 2,5-octadiene-1,8-diyl group, 2,6-octadiene-1,8-diyl group, 2,4,6-octatriene-1,8-diyl group, 2-nonene-1,9-diyl group, 3-nonene-1,9-diyl group, 4-nonene-1,9-diyl group, 5-nonene-1,9-diyl group, 6-nonene-1,9-diyl group, 7-
 25 nonene-1,9-diyl group, 2-decene-1,10-diyl group, 3-decene-1,10-diyl group, 4-decene-1,10-diyl group, 5-decene-1,10-diyl group, 6-decene-1,10-diyl group, 7-decene-1,10-diyl group, 8-decene-1,10-diyl group;

- 2-undecene-1,11-diyl group, 3-undecene-1,11-diyl group, 4-undecene-1,11-diyl group, 5-undecene-1,11-diyl group, 6-undecene-1,11-diyl group, 7-undecene-1,11-diyl group, 8-undecene-1,11-diyl group, 9-undecene-1,11-diyl group;
- 5 2-dodecene-1,11-diyl group, 3-dodecene-1,11-diyl group, 4-dodecene-1,11-diyl group, 5-dodecene-1,11-diyl group, 6-dodecene-1,11-diyl group, 7-dodecene-1,11-diyl group, 8-dodecene-1,11-diyl group, 9-dodecene-1,11-diyl group, 10-dodecene-1,11-diyl group;
- 10 2-tridecene-1,13-diyl group, 3-tridecene-1,13-diyl group, 4-tridecene-1,13-diyl group, 5-tridecene-1,13-diyl group, 6-tridecene-1,13-diyl group, 7-tridecene-1,13-diyl group, 8-tridecene-1,13-diyl group, 9-tridecene-1,13-diyl group, 10-tridecene-1,13-diyl group, 11-tridecene-1,13-diyl group;
- 15 2-tetradecene-1,14-diyl group, 3-tetradecene-1,14-diyl group, 4-tetradecene-1,14-diyl group, 5-tetradecene-1,14-diyl group, 6-tetradecene-1,14-diyl group, 7-tetradecene-1,14-diyl group, 8-tetradecene-1,14-diyl group, 9-tetradecene-1,14-diyl group, 10-tetradecene-1,14-diyl group,
- 20 11-tetradecene-1,14-diyl group, 12-tetradecene-1,14-diyl group;
- 2 2-pentadecene-1,15-diyl group, 3-pentadecene-1,15-diyl group, 4-pentadecene-1,15-diyl group, 5-pentadecene-1,15-diyl group, 6-pentadecene-1,15-diyl group, 7-pentadecene-1,15-diyl group, 8-pentadecene-1,15-diyl group, 9-
- 25 pentadecene-1,15-diyl group, 10-pentadecene-1,15-diyl group, 11-pentadecene-1,15-diyl group, 12-pentadecene-1,15-diyl group, 13-pentadecene-1,15-diyl group;

- 2-hexadecene-1,16-diyl group, 3-hexadecene-1,16-diyl group,
 4-hexadecene-1,16-diyl group, 5-hexadecene-1,16-diyl group,
 6-hexadecene-1,16-diyl group, 7-hexadecene-1,16-diyl group,
 8-hexadecene-1,16-diyl group, 9-hexadecene-1,16-diyl group,
 5 10-hexadecene-1,16-diyl group, 11-hexadecene-1,16-diyl
 group, 12-hexadecene-1,16-diyl group, 13-hexadecene-1,16-
 diyl group, 14-hexadecene-1,16-diyl group;
 2-heptadecene-1,17-diyl group, 3-heptadecene-1,17-diyl
 group, 4-heptadecene-1,17-diyl group, 5-heptadecene-1,17-
 10 diyl group, 6-heptadecene-1,17-diyl group, 7-heptadecene-
 1,17-diyl group, 8-heptadecene-1,17-diyl group, 9-
 heptadecene-1,17-diyl group, 10-heptadecene-1,17-diyl group,
 11-heptadecene-1,17-diyl group, 12-heptadecene-1,17-diyl
 group, 13-heptadecene-1,17-diyl group, 14-heptadecene-1,17-
 15 diyl group, 15-heptadecene-1,17-diyl group;
 2-octadecene-1,18-diyl group, 3-octadecene-1,18-diyl group,
 4-octadecene-1,18-diyl group, 5-octadecene-1,18-diyl group,
 6-octadecene-1,18-diyl group, 7-octadecene-1,18-diyl group,
 8-octadecene-1,18-diyl group, 9-octadecene-1,18-diyl group,
 20 10-octadecene-1,18-diyl group, 11-octadecene-1,18-diyl
 group, 12-octadecene-1,18-diyl group, 13-octadecene-1,18-
 diyl group, 14-octadecene-1,18-diyl group, 15-octadecene-
 1,18-diyl group, 16-octadecene-1,18-diyl group;
 2-nonadecene-1,19-diyl group, 3-nonadecene-1,19-diyl group,
 25 4-nonadecene-1,19-diyl group, 5-nonadecene-1,19-diyl group,
 6-nonadecene-1,19-diyl group, 7-nonadecene-1,19-diyl group,
 8-nonadecene-1,19-diyl group, 9-nonadecene-1,19-diyl group,
 10-nonadecene-1,19-diyl group, 11-nonadecene-1,19-diyl

- group, 12-nonadecene-1,19-diyl group, 13-nonadecene-1,19-diyl group, 14-nonadecene-1,19-diyl group, 15-nonadecene-1,19-diyl group, 16-nonadecene-1,19-diyl group, 17-nonadecene-1,19-diyl group;
- 5 2-icosene-1,20-diyl group, 3-icosene-1,20-diyl group, 4-icosene-1,20-diyl group, 5-icosene-1,20-diyl group, 6-icosene-1,20-diyl group, 7-icosene-1,20-diyl group, 8-icosene-1,20-diyl group, 9-icosene-1,20-diyl group, 10-icosene-1,20-diyl group, 11-icosene-1,20-diyl group, 12-icosene-1,20-diyl group, 13-icosene-1,20-diyl group, 14-icosene-1,20-diyl group, 15-icosene-1,20-diyl group, 16-icosene-1,20-diyl group, 17-icosene-1,20-diyl group, 18-icosene-1,20-diyl group;
- 15 2-henicosene-1,21-diyl group, 3-henicosene-1,21-diyl group, 4-henicosene-1,21-diyl group, 5-henicosene-1,21-diyl group, 6-henicosene-1,21-diyl group, 7-henicosene-1,21-diyl group, 8-henicosene-1,21-diyl group, 9-henicosene-1,21-diyl group, 10-henicosene-1,21-diyl group, 11-henicosene-1,21-diyl group, 12-henicosene-1,21-diyl group, 13-henicosene-1,21-diyl group, 14-henicosene-1,21-diyl group, 15-henicosene-1,21-diyl group, 16-henicosene-1,21-diyl group, 17-henicosene-1,21-diyl group, 18-henicosene-1,21-diyl group, 19-henicosene-1,21-diyl group;
- 20 2-docosene-1,22-diyl group, 3-docosene-1,22-diyl group, 4-docosene-1,22-diyl group, 5-docosene-1,22-diyl group, 6-docosene-1,22-diyl group, 7-docosene-1,22-diyl group, 8-docosene-1,22-diyl group, 9-docosene-1,22-diyl group, 10-docosene-1,22-diyl group, 11-docosene-1,22-diyl group, 12-

- docosene-1,22-diyl group, 13-docosene-1,22-diyl group, 14-docosene-1,22-diyl group, 15-docosene-1,22-diyl group, 16-docosene-1,22-diyl group, 17-docosene-1,22-diyl group, 18-docosene-1,22-diyl group, 19-docosene-1,22-diyl group, 20-
- 5 docosene-1,22-diyl group;
- 2-tricosene-1,23-diyl group, 3-tricosene-1,23-diyl group, 4-tricosene-1,23-diyl group, 5-tricosene-1,23-diyl group, 6-tricosene-1,23-diyl group, 7-tricosene-1,23-diyl group, 8-tricosene-1,23-diyl group, 9-tricosene-1,23-diyl group,
- 10 10-tricosene-1,23-diyl group, 11-tricosene-1,23-diyl group, 12-tricosene-1,23-diyl group, 13-tricosene-1,23-diyl group, 14-tricosene-1,23-diyl group, 15-tricosene-1,23-diyl group, 16-tricosene-1,23-diyl group, 17-tricosene-1,23-diyl group, 18-tricosene-1,23-diyl group, 19-tricosene-1,23-diyl group,
- 15 20-tricosene-1,23-diyl group, 21-tricosene-1,23-diyl group;
- 2-tetracosene-1,24-diyl group, 3-tetracosene-1,24-diyl group, 4-tetracosene-1,24-diyl group, 5-tetracosene-1,24-diyl group, 6-tetracosene-1,24-diyl group, 7-tetracosene-1,24-diyl group, 8-tetracosene-1,24-diyl group, 9-
- 20 tetracosene-1,24-diyl group, 10-tetracosene-1,24-diyl group, 11-tetracosene-1,24-diyl group, 12-tetracosene-1,24-diyl group, 13-tetracosene-1,24-diyl group, 14-tetracosene-1,24-diyl group, 15-tetracosene-1,24-diyl group, 16-tetracosene-1,24-diyl group, 17-tetracosene-1,24-diyl group, 18-
- 25 tetracosene-1,24-diyl group, 19-tetracosene-1,24-diyl group, 20-tetracosene-1,24-diyl group, 21-tetracosene-1,24-diyl group, 22-tetracosene-1,24-diyl group;
- 2-pentacosene-1,25-diyl group, 3-pentacosene-1,25-diyl

group, 4-pentacosene-1,25-diyl group, 5-pentacosene-1,25-diyl group, 6-pentacosene-1,25-diyl group, 7-pentacosene-1,25-diyl group, 8-pentacosene-1,25-diyl group, 8-pentacosene-1,25-diyl group, 9-pentacosene-1,25-diyl group, 5 10-pentacosene-1,25-diyl group, 11-pentacosene-1,25-diyl group, 12-pentacosene-1,25-diyl group, 13-pentacosene-1,25-diyl group, 14-pentacosene-1,25-diyl group, 15-pentacosene-1,25-diyl group, 16-pentacosene-1,25-diyl group, 17-pentacosene-1,25-diyl group, 18-pentacosene-1,25-diyl group, 10 19-pentacosene-1,25-diyl group, 20-pentacosene-1,25-diyl group, 21-pentacosene-1,25-diyl group, 22-pentacosene-1,25-diyl group, 23-pentacosene-1,25-diyl group;

2-hexacosene-1,26-diyl group, 3-hexacosene-1,26-diyl group, 4-hexacosene-1,26-diyl group, 5-hexacosene-1,26-diyl group, 15 6-hexacosene-1,26-diyl group, 7-hexacosene-1,26-diyl group, 8-hexacosene-1,26-diyl group, 9-hexacosene-1,26-diyl group, 10-hexacosene-1,26-diyl group, 11-hexacosene-1,26-diyl group, 12-hexacosene-1,26-diyl group, 13-hexacosene-1,26-diyl group, 14-hexacosene-1,26-diyl group, 15-hexacosene-1,26-diyl group, 16-hexacosene-1,26-diyl group, 17-hexacosene-1,26-diyl group, 18-hexacosene-1,26-diyl group, 19-hexacosene-1,26-diyl group, 20-hexacosene-1,26-diyl group, 21-hexacosene-1,26-diyl group, 22-hexacosene-1,26-diyl group, 23-hexacosene-1,26-diyl group, 24-hexacosene-1,26-diyl group; 25

2-heptacosene-1,27-diyl group, 3-heptacosene-1,27-diyl group, 4-heptacosene-1,27-diyl group, 5-heptacosene-1,27-diyl group, 6-heptacosene-1,27-diyl group, 7-heptacosene-

- 1,27-diyl group, 8-heptacosene-1,27-diyl group, 9-heptacosene-1,27-diyl group, 10-heptacosene-1,27-diyl group, 11-heptacosene-1,27-diyl group, 12-heptacosene-1,27-diyl group, 13-heptacosene-1,27-diyl group, 14-heptacosene-1,27-diyl group, 15-heptacosene-1,27-diyl group, 16-heptacosene-1,27-diyl group, 17-heptacosene-1,27-diyl group, 18-heptacosene-1,27-diyl group, 19-heptacosene-1,27-diyl group, 20-heptacosene-1,27-diyl group, 21-heptacosene-1,27-diyl group, 22-heptacosene-1,27-diyl group, 23-heptacosene-1,27-diyl group, 24-heptacosene-1,27-diyl group, 25-heptacosene-1,27-diyl group;
- 2-octacosene-1,28-diyl group, 3-octacosene-1,28-diyl group, 4-octacosene-1,28-diyl group, 5-octacosene-1,28-diyl group, 6-octacosene-1,28-diyl group, 7-octacosene-1,28-diyl group, 8-octacosene-1,28-diyl group, 9-octacosene-1,28-diyl group, 10-octacosene-1,28-diyl group, 11-octacosene-1,28-diyl group, 12-octacosene-1,28-diyl group, 13-octacosene-1,28-diyl group, 14-octacosene-1,28-diyl group, 15-octacosene-1,28-diyl group, 16-octacosene-1,28-diyl group, 17-octacosene-1,28-diyl group, 18-octacosene-1,28-diyl group, 19-octacosene-1,28-diyl group, 20-octacosene-1,28-diyl group, 21-octacosene-1,28-diyl group, 22-octacosene-1,28-diyl group, 23-octacosene-1,28-diyl group, 24-octacosene-1,28-diyl group, 25-octacosene-1,28-diyl group, 26-octacosene-1,28-diyl group;
- 2-nonacosene-1,29-diyl group, 3-nonacosene-1,29-diyl group, 4-nonacosene-1,29-diyl group, 5-nonacosene-1,29-diyl group, 6-nonacosene-1,29-diyl group, 7-nonacosene-1,29-diyl group,

8-nonacosene-1,29-diyl group, 9-nonacosene-1,29-diyl group,
 10-nonacosene-1,29-diyl group, 11-nonacosene-1,29-diyl
 group, 12-nonacosene-1,29-diyl group, 13-nonacosene-1,29-
 diyl group, 14-nonacosene-1,29-diyl group, 15-nonacosene-
 5 1,29-diyl group, 16-nonacosene-1,29-diyl group, 17-
 nonacosene-1,29-diyl group, 18-nonacosene-1,29-diyl group,
 19-nonacosene-1,29-diyl group, 20-nonacosene-1,29-diyl
 group, 21-nonacosene-1,29-diyl group, 22-nonacosene-1,29-
 diyl group, 23-nonacosene-1,29-diyl group, 24-nonacosene-
 10 1,29-diyl group, 25-nonacosene-1,29-diyl group, 26-
 nonacosene-1,29-diyl group, 27-nonacosene-1,29-diyl group;
 2-triacontene-1,30-diyl group, 3-triacontene-1,30-diyl
 group, 4-triacontene-1,30-diyl group, 5-triacontene-1,30-
 diyl group, 6-triacontene-1,30-diyl group, 7-triacontene-
 15 1,30-diyl group, 8-triacontene-1,30-diyl group, 9-
 triacontene-1,30-diyl group, 10-triacontene-1,30-diyl group,
 11-triacontene-1,30-diyl group, 12-triacontene-1,30-diyl
 group, 13-triacontene-1,30-diyl group, 14-triacontene-1,30-
 diyl group, 15-triacontene-1,30-diyl group, 16-triacontene-
 20 1,30-diyl group, 17-triacontene-1,30-diyl group, 18-
 triacontene-1,30-diyl group, 19-triacontene-1,30-diyl group,
 20-triacontene-1,30-diyl group, 21-triacontene-1,30-diyl
 group, 22-triacontene-1,30-diyl group, 23-triacontene-1,30-
 diyl group, 24-triacontene-1,30-diyl group, 25-triacontene-
 25 1,30-diyl group, 26-triacontene-1,30-diyl group, 27-
 triacontene-1,30-diyl group, and 28-triacontene-1,30-diyl
 group;
 as well as branched alkenylene groups such as 1-

methlethylene-1,2-diyl group, 2-methyl-1-propene-1,3-diyl group, 2-methyl-2-propene-1,3-diyl group, 2-methyl-1-butene-1,4-diyl group, 3-methyl-2-butene-1,4-diyl group, 2-methyl-3-butene-1,4-diyl group, 2,3-dimethyl-1,3-butadiene-
 5 1,4-diyl group, 3-ethyl-2-propene-1,5-diyl group, 4-methyl-3-propene-1,5-diyl group, 3-methyl-2,4-propadiene-1,5-diyl group, 3,4-diethyl-2-hexene-1,6-diyl group, 4-methyl-3-hexene-1,6-diyl group, 2-methyl-4-hexene-1,6-diyl group, 3,5-dimethyl-2,4-hexadiene-1,6-diyl group, 5-ethyl-3-
 10 methyl-2-heptene-1,7-diyl group, 5-methyl-3-heptene-1,7-diyl group, 4-n-propyl-4-heptene-1,7-diyl group, 3,6-dimethyl-5-heptene-1,7-diyl group, 5-ethyl-2,4-heptadiene-1,7-diyl group, 2,6-dimethyl-2,5-heptadiene-1,7-diyl group, 4-ethyl-3,5-heptadiene-1,7-diyl group, 4-ethyl-6,6-
 15 dimethyl-2-octene-1,8-diyl group, 5-n-propyl-3-octene-1,8-diyl group, 3-ethyl-4-octene-1,8-diyl group, 4-ethyl-2-methyl-6-i-propyl-5-octene-1,8-diyl group, 3,4,5-trimethyl-6-octene-1,8-diyl group, 5-ethyl-7-methyl-2,4-octadiene-1,8-diyl group, 3-methyl-2,5-octadiene-1,8-diyl group, 5-n-
 20 propyl-2,6-octadiene-1,8-diyl group, 4-methyl-2,4,6-octatriene-1,8-diyl group, 5-ethyl-2-nonene-1,9-diyl group, 3,5,6-trimethyl-3-nonene-1,9-diyl group, 2,4,5,7-tetramethyl-4-nonene-1,9-diyl group, 3,4-diethyl-5-nonene-1,9-diyl group, 4-i-propyl-6-nonene-1,9-diyl group, 3-
 25 ethyl-7-nonene-1,9-diyl group, 5-n-butyl-2-decene-1,10-diyl group, 6-i-propyl-3-decene-1,10-diyl group, 5-ethyl-4-decene-1,10-diyl group, 6,7-dimethyl-5-decene-1,10-diyl group, 4-ethyl-6-decene-1,10-diyl group, 5-methyl-7-decene-

- 1,10-diyl group, 6-ethyl-4-methyl-8-decene-1,10-diyl group;
 6-methyl-2-undecene-1,11-diyl group, 4-ethyl-3-undecene-
 1,11-diyl group, 5-methyl-4-undecene-1,11-diyl group, 7-
 ethyl-5-undecene-1,11-diyl group, 5-methyl-6-undecene-1,11-
 5 diyl group, 9-ethyl-7-undecene-1,11-diyl group, 3-methyl-8-
 undecene-1,11-diyl group, 4-ethyl-9-undecene-1,11-diyl
 group;
- 4-ethyl-2-dodecene-1,12-diyl group, 5-methyl-3-dodecene-
 1,12-diyl group, 6-ethyl-4-dodecene-1,12-diyl group, 7-
 10 methyl-5-dodecene-1,12-diyl group, 8-ethyl-6-dodecene-1,12-
 diyl group, 9-methyl-7-dodecene-1,12-diyl group, 10-ethyl-
 8-dodecene-1,12-diyl group, 2-methyl-9-dodecene-1,12-diyl
 group, 5-ethyl-10-dodecene-1,12-diyl group;
- 4,7,9-trimethyl-2-tridecene-1,13-diyl group, 10-methyl-3-
 15 tridecene-1,13-diyl group, 8-ethyl-4-tridecene-1,13-diyl
 group, 4-methyl-5-tridecene-1,13-diyl group, 5-ethyl-6-
 tridecene-1,13-diyl group, 3,6-diethyl-7-tridecene-1,13-
 diyl group, 5-methyl-8-tridecene-1,13-diyl group, 7-ethyl-
 9-tridecene-1,13-diyl group, 4-methyl-10-tridecene-1,13-
 20 diyl group, 6-ethyl-11-tridecene-1,13-diyl group;
- 7-methyl-2-tetradecene-1,14-diyl group, 8-ethyl-3-
 tetradecene-1,14-diyl group, 6-n-propyl-4-tetradecene-1,14-
 diyl group, 8-methyl-5-tetradecene-1,14-diyl group, 3-
 ethyl-6-tetradecene-1,14-diyl group, 10-methyl-7-
 25 tetradecene-1,14-diyl group, 6-i-propyl-8-tetradecene-1,14-
 diyl group, 5,7,11-trimethyl-9-tetradecene-1,14-diyl group,
 5-ethyl-10-tetradecene-1,14-diyl group, 6-methyl-11-
 tetradecene-1,14-diyl group, 4-n-butyl-12-tetradecene-1,14-

- diyl group;
- 4-methyl-2-pentadecene-1,15-diyl group, 6-ethyl-3-pentadecene-1,15-diyl group, 8-methyl-4-pentadecene-1,15-diyl group, 10-ethyl-5-pentadecene-1,15-diyl group, 4,9-
- 5 dimethyl-6-pentadecene-1,15-diyl group, 10-ethyl-7-pentadecene-1,15-diyl group, 6-methyl-8-pentadecene-1,15-diyl group, 8-n-propyl-9-pentadecene-1,15-diyl group, 5-methyl-10-pentadecene-1,15-diyl group, 4,7-diethyl-11-pentadecene-1,15-diyl group, 5-methyl-12-pentadecene-1,15-
- 10 diyl group, 8-ethyl-13-pentadecene-1,15-diyl group;
- 8-i-propyl-2-hexadecene-1,16-diyl group, 6-methyl-3-hexadecene-1,16-diyl group, 8-ethyl-4-hexadecene-1,16-diyl group, 9-methyl-5-hexadecene-1,16-diyl group, 10-ethyl-6-hexadecene-1,16-diyl group, 5-methyl-7-hexadecene-1,16-diyl
- 15 group, 5,10-dimethyl-8-hexadecene-1,16-diyl group, 5-ethyl-9-hexadecene-1,16-diyl group, 7,12-diethyl-10-hexadecene-1,16-diyl group, 5-ethyl-7-methyl-11-hexadecene-1,16-diyl group, 5-methyl-12-hexadecene-1,16-diyl group, 8-s-butyl-13-hexadecene-1,16-diyl group, 5-ethyl-14-hexadecene-1,16-
- 20 diyl group;
- 11-methyl-2-heptadecene-1,17-diyl group, 9-ethyl-3-heptadecene-1,17-diyl group, 6-i-propyl-4-heptadecene-1,17-diyl group, 8-methyl-5-heptadecene-1,17-diyl group, 4-ethyl-6-heptadecene-1,17-diyl group, 10-methyl-7-
- 25 heptadecene-1,17-diyl group, 5,11-dimethyl-8-heptadecene-1,17-diyl group, 5-ethyl-9-heptadecene-1,17-diyl group, 8-ethyl-10-heptadecene-1,17-diyl group, 7-methyl-11-heptadecene-1,17-diyl group, 5-i-propyl-12-heptadecene-

- 1,17-diyl group, 9-ethyl-13-heptadecene-1,17-diyl group, 8-methyl-14-heptadecene-1,17-diyl group, 7-s-butyl-15-heptadecene-1,17-diyl group;
- 10,15-dimethyl-2-octadecene-1,18-diyl group, 6-ethyl-3-octadecene-1,18-diyl group, 10-methyl-4-octadecene-1,18-diyl group, 11-methyl-5-octadecene-1,18-diyl group, 12-ethyl-6-octadecene-1,18-diyl group, 10-methyl-7-octadecene-1,18-diyl group, 5-methyl-8-octadecene-1,18-diyl group, 8-ethyl-9-octadecene-1,18-diyl group, 7-methyl-10-octadecene-1,18-diyl group, 9-n-butyl-11-octadecene-1,18-diyl group, 7-methyl-12-octadecene-1,18-diyl group, 9-ethyl-13-octadecene-1,18-diyl group, 10-i-propyl-14-octadecene-1,18-diyl group, 7-methyl-15-octadecene-1,18-diyl group, 10-ethyl-16-octadecene-1,18-diyl group;
- 10-methyl-2-nonadecene-1,19-diyl group, 10,12-diethyl-3-nonadecene-1,19-diyl group, 6-methyl-4-nonadecene-1,19-diyl group, 7-ethyl-5-nonadecene-1,19-diyl group, 9-n-propyl-6-nonadecene-1,19-diyl group, 10-methyl-7-nonadecene-1,19-diyl group, 12-i-propyl-8-nonadecene-1,19-diyl group, 5,15-dimethyl-9-nonadecene-1,19-diyl group, 7-ethyl-13-methyl-10-nonadecene-1,19-diyl group, 6-methyl-11-nonadecene-1,19-diyl group, 6-ethyl-12-nonadecene-1,19-diyl group, 7,15-diethyl-13-nonadecene-1,19-diyl group, 9-s-butyl-14-nonadecene-1,19-diyl group, 8-methyl-15-nonadecene-1,19-diyl group, 10-ethyl-16-nonadecene-1,19-diyl group, 10-i-propyl-17-nonadecene-1,19-diyl group;
- 8-methyl-2-icosene-1,20-diyl group, 6-ethyl-3-icosene-1,20-diyl group, 10-i-propyl-4-icosene-1,20-diyl group, 11-n-

- propyl-5-icosene-1,20-diyl group, 12-methyl-6-icosene-1,20-diyl group, 11-ethyl-7-icosene-1,20-diyl group, 13-n-propyl-8-icosene-1,20-diyl group, 8-i-propyl-9-icosene-1,20-diyl group, 8-n-propyl-10-icosene-1,20-diyl group, 7-methyl-11-icosene-1,20-diyl group, 8-ethyl-12-icosene-1,20-diyl group, 10-n-propyl-13-icosene-1,20-diyl group, 9-i-propyl-14-icosene-1,20-diyl group, 10-n-butyl-15-icosene-1,20-diyl group, 8-s-butyl-16-icosene-1,20-diyl group, 7-i-butyl-17-icosene-1,20-diyl group, 9-methyl-18-icosene-1,20-diyl group;
- 11-methyl-2-henicosene-1,21-diyl group, 12-n-butyl-3-henicosene-1,21-diyl group, 10-n-pentyl-4-henicosene-1,21-diyl group, 8-ethyl-5-henicosene-1,21-diyl group, 10-i-propyl-6-henicosene-1,21-diyl group, 5-n-propyl-7-henicosene-1,21-diyl group, 13-n-butyl-8-henicosene-1,21-diyl group, 15-s-butyl-9-henicosene-1,21-diyl group, 5-methyl-10-henicosene-1,21-diyl group, 15-ethyl-6-methyl-11-henicosene-1,21-diyl group, 8-ethyl-12-henicosene-1,21-diyl group, 7-methyl-13-henicosene-1,21-diyl group, 11-ethyl-14-henicosene-1,21-diyl group, 6-ethyl-15-henicosene-1,21-diyl group, 9-methyl-16-henicosene-1,21-diyl group, 5-ethyl-9-methyl-17-henicosene-1,21-diyl group, 10,10-dimethyl-18-henicosene-1,21-diyl group, 9-ethyl-19-henicosene-1,21-diyl group;
- 11-methyl-2-docosene-1,22-diyl group, 12-ethyl-3-docosene-1,22-diyl group, 13-i-propyl-4-docosene-1,22-diyl group, 10-n-propyl-5-docosene-1,22-diyl group, 10-n-butyl-6-docosene-1,22-diyl group, 15-s-butyl-7-docosene-1,22-diyl

group, 11-i-butyl-8-docosene-1,22-diyl group, 5,15-dimethyl-9-docosene-1,22-diyl group, 8,14-diethyl-10-docosene-1,22-diyl group, 5-methyl-11-docosene-1,22-diyl group, 7-ethyl-12-docosene-1,22-diyl group, 10-methyl-13-docosene-1,22-diyl group, 10-ethyl-14-docosene-1,22-diyl group, 9-ethyl-15-docosene-1,22-diyl group, 8-methyl-16-docosene-1,22-diyl group, 7-i-propyl-17-docosene-1,22-diyl group, 10-i-butyl-18-docosene-1,22-diyl group, 9,10-dimethyl-19-docosene-1,22-diyl group, 13-ethyl-20-docosene-1,22-diyl group;

19-methyl-2-tricosene-1,23-diyl group, 10,15-dimethyl-3-tricosene-1,23-diyl group, 3,11,16-trimethyl-4-tricosene-1,23-diyl group, 12-ethyl-5-tricosene-1,23-diyl group, 6,13-diethyl-6-tricosene-1,23-diyl group, 4,12,18-triethyl-7-tricosene-1,23-diyl group, 18-i-propyl-8-tricosene-1,23-diyl group, 14-n-propyl-9-tricosene-1,23-diyl group, 8-n-butyl-10-tricosene-1,23-diyl group, 15-s-butyl-11-tricosene-1,23-diyl group, 5-i-butyl-12-tricosene-1,23-diyl group, 7-ethyl-9-methyl-13-tricosene-1,23-diyl group, 9-methyl-14-tricosene-1,23-diyl group, 4,18-dimethyl-15-tricosene-1,23-diyl group, 3,4,11-trimethyl-16-tricosene-1,23-diyl group, 9-ethyl-17-tricosene-1,23-diyl group, 10,13-diethyl-18-tricosene-1,23-diyl group, 5,8,21-triethyl-19-tricosene-1,23-diyl group, 15-i-propyl-20-tricosene-1,23-diyl group, 17-n-propyl-21-tricosene-1,23-diyl group;

16-n-butyl-2-tetracosene-1,24-diyl group, 11-s-butyl-3-tetracosene-1,24-diyl group, 8-i-butyl-4-tetracosene-1,24-

diyl group, 18-ethyl-9-methyl-5-tetracosene-1,24-diyl group,
 13-methyl-6-tetracosene-1,24-diyl group, 4,19-dimethyl-7-
 tetracosene-1,24-diyl group, 5,10,17-triethyl-8-
 tetracosene-1,24-diyl group, 6-ethyl-9-tetracosene-1,24-
 5 diyl group, 7,16-diethyl-10-tetracosene-1,24-diyl group,
 5,9,18-triethyl-11-tetracosene-1,24-diyl group, 10-n-
 propyl-12-tetracosene-1,24-diyl group, 20-i-propyl-13-
 tetracosene-1,24-diyl group, 9-n-butyl-14-tetracosene-1,24-
 diyl group, 11-s-butyl-15-tetracosene-1,24-diyl group, 13-
 10 i-butyl-16-tetracosene-1,24-diyl group, 10-ethyl-13-methyl-
 17-tetracosene-1,24-diyl group, 6-methyl-18-tetracosene-
 1,24-diyl group, 5,7-dimethyl-19-tetracosene-1,24-diyl
 group, 4,8,13-trimethyl-20-tetracosene-1,24-diyl group, 18-
 ethyl-21-tetracosene-1,24-diyl group, 6,10-diethyl-22-
 15 tetracosene-1,24-diyl group;
 9,13,16-trimethyl-2-pentacosene-1,25-diyl group, 12-n-
 propyl-3-pentacosene-1,25-diyl group, 11-i-propyl-4-
 pentacosene-1,25-diyl group, 20-n-butyl-5-pentacosene-1,25-
 diyl group, 17-i-butyl-6-pentacosene-1,25-diyl group, 15-s-
 20 butyl-7-pentacosene-1,25-diyl group, 15-ethyl-23-methyl-8-
 pentacosene-1,25-diyl group, 11-methyl-8-pentacosene-1,25-
 diyl group, 13,17-dimethyl-9-pentacosene-1,25-diyl group,
 5,8,21-trimethyl-10-pentacosene-1,25-diyl group, 17-ethyl-
 11-pentacosene-1,25-diyl group, 8,18-dimethyl-12-
 25 pentacosene-1,25-diyl group, 10,15,18-trimethyl-13-
 pentacosene-1,25-diyl group, 4-n-propyl-14-pentacosene-
 1,25-diyl group, 20-i-propyl-15-pentacosene-1,25-diyl group,
 8-n-butyl-16-pentacosene-1,25-diyl group, 11-s-butyl-17-

- pentacosene-1,25-diyl group, 5,22-dimethyl-18-pentacosene-1,25-diyl group, 5-i-butyl-19-pentacosene-1,25-diyl group, 9-methyl-13-ethyl-20-pentacosene-1,25-diyl group, 15-methyl-21-pentacosene-1,25-diyl group, 6,13-dimethyl-22-pentacosene-1,25-diyl group, 4,8,12-trimethyl-23-pentacosene-1,25-diyl group;
- 13-ethyl-2-hexacosene-1,26-diyl group, 5,16-diethyl-3-hexacosene-1,26-diyl group, 7,11,16-trimethyl-4-hexacosene-1,26-diyl group, 12-n-propyl-5-hexacosene-1,26-diyl group, 21-i-propyl-6-hexacosene-1,26-diyl group, 6-n-butyl-7-hexacosene-1,26-diyl group, 13-s-butyl-8-hexacosene-1,26-diyl group, 19-i-butyl-9-hexacosene-1,26-diyl group, 13-ethyl-18-methyl-10-hexacosene-1,26-diyl group, 10-methyl-11-hexacosene-1,26-diyl group, 10,20-dimethyl-12-hexacosene-1,26-diyl group, 7,9,17-trimethyl-13-hexacosene-1,26-diyl group, 8-ethyl-14-hexacosene-1,26-diyl group, 5,22-diethyl-15-hexacosene-1,26-diyl group, 7,10,21-trimethyl-16-hexacosene-1,26-diyl group, 15-n-propyl-17-hexacosene-1,26-diyl group, 13-i-propyl-18-hexacosene-1,26-diyl group, 8-n-butyl-19-hexacosene-1,26-diyl group, 11-s-butyl-20-hexacosene-1,26-diyl group, 14-i-butyl-21-hexacosene-1,26-diyl group, 5-ethyl-21-methyl-22-hexacosene-1,26-diyl group, 7-methyl-23-hexacosene-1,26-diyl group, 8,14-dimethyl-24-hexacosene-1,26-diyl group;
- 7,16,24-trimethyl-2-heptacosene-1,27-diyl group, 9-ethyl-3-heptacosene-1,27-diyl group, 7,16-dimethyl-4-heptacosene-1,27-diyl group, 9,13,21-trimethyl-5-heptacosene-1,27-diyl group, 13-n-propyl-6-heptacosene-1,27-diyl group, 10-i-

- propyl-7-heptacosene-1,27-diyl group, 16-n-propyl-8-heptacosene-1,27-diyl group, 18-methyl-9-heptacosene-1,27-diyl group, 9-i-propyl-10-heptacosene-1,27-diyl group, 15-ethyl-7-methyl-11-heptacosene-1,27-diyl group, 25-methyl-
- 5 12-heptacosene-1,27-diyl group, 8,21-dimethyl-13-heptacosene-1,27-diyl group, 5,11,23-trimethyl-14-heptacosene-1,27-diyl group, 9-ethyl-15-heptacosene-1,27-diyl group, 8,20-dimethyl-16-heptacosene-1,27-diyl group, 4,8,19-trimethyl-17-heptacosene-1,27-diyl group, 7-n-
- 10 propyl-18-heptacosene-1,27-diyl group, 21-i-propyl-19-heptacosene-1,27-diyl group, 14-n-propyl-20-heptacosene-1,27-diyl group, 8-ethyl-21-heptacosene-1,27-diyl group, 11-i-propyl-22-heptacosene-1,27-diyl group, 5-ethyl-13-methyl-23-heptacosene-1,27-diyl group, 16-methyl-24-
- 15 heptacosene-1,27-diyl group, 7-ethyl-25-heptacosene-1,27-diyl group;
- 14-ethyl-2-octacosene-1,28-diyl group, 20-methyl-3-octacosene-1,28-diyl group, 7,22-dimethyl-4-octacosene-1,28-diyl group, 19-ethyl-5-octacosene-1,28-diyl group, 11-
- 20 methyl-6-octacosene-1,28-diyl group, 13,16-dimethyl-7-octacosene-1,28-diyl group, 13-ethyl-8-octacosene-1,28-diyl group, 6-methyl-9-octacosene-1,28-diyl group, 9,16-dimethyl-10-octacosene-1,28-diyl group, 7-ethyl-11-octacosene-1,28-diyl group, 16-methyl-12-octacosene-1,28-
- 25 diyl group, 6,15-dimethyl-13-octacosene-1,28-diyl group, 22-ethyl-14-octacosene-1,28-diyl group, 6-methyl-15-octacosene-1,28-diyl group, 8,11-dimethyl-16-octacosene-1,28-diyl group, 23-ethyl-17-octacosene-1,28-diyl group, 4-

methyl-18-octacosene-1,28-diyl group, 7,14-dimethyl-19-octacosene-1,28-diyl group, 13-ethyl-20-octacosene-1,28-diyl group, 8-methyl-21-octacosene-1,28-diyl group, 11,17-dimethyl-22-octacosene-1,28-diyl group, 10-ethyl-23-octacosene-1,28-diyl group, 9-methyl-24-octacosene-1,28-diyl group, 7,19-dimethyl-25-octacosene-1,28-diyl group, 12-ethyl-26-octacosene-1,28-diyl group;

15-methyl-2-nonacosene-1,29-diyl group, 14-methyl-3-nonacosene-1,29-diyl group, 12-methyl-4-nonacosene-1,29-diyl group, 13-methyl-5-nonacosene-1,29-diyl group, 11-methyl-6-nonacosene-1,29-diyl group, 10-methyl-7-nonacosene-1,29-diyl group, 25-methyl-8-nonacosene-1,29-diyl group, 24-methyl-9-nonacosene-1,29-diyl group, 23-methyl-10-nonacosene-1,29-diyl group, 22-methyl-11-nonacosene-1,29-diyl group, 21-methyl-12-nonacosene-1,29-diyl group, 20-methyl-13-nonacosene-1,29-diyl group, 19-methyl-14-nonacosene-1,29-diyl group, 18-methyl-15-nonacosene-1,29-diyl group, 27-methyl-16-nonacosene-1,29-diyl group, 26-methyl-17-nonacosene-1,29-diyl group, 25-methyl-18-nonacosene-1,29-diyl group, 24-methyl-19-nonacosene-1,29-diyl group, 23-methyl-20-nonacosene-1,29-diyl group, 20-methyl-21-nonacosene-1,29-diyl group, 19-methyl-22-nonacosene-1,29-diyl group, 18-methyl-23-nonacosene-1,29-diyl group, 17-methyl-24-nonacosene-1,29-diyl group, 16-methyl-25-nonacosene-1,29-diyl group, 6-methyl-26-nonacosene-1,29-diyl group, and 5-methyl-27-nonacosene-1,29-diyl group.

If G represents an optionally substituted straight-

chained or branched alkynylene group having 2 - 30 carbon atoms, exemplary straight-chained or branched alkynylene groups having 2 - 30 carbon atoms include straight-chained alkynylene groups such as acetylene-1,2-diyl group, 1-
 5 propyne-1,3-diyl group, 2-propyne-1,3-diyl group, 1-butyne-1,4-diyl group, 2-butyne-1,4-diyl group, 3-butyne-1,4-diyl group, 1,3-butadiyne-1,4-diyl group, 2-pentyne-1,5-diyl group, 3-pentyne-1,5-diyl group, 2,4-pentadiyne-1,5-diyl group, 2-hexyne-1,6-diyl group, 3-hexyne-1,6-diyl group, 4-
 10 hexyne-1,6-diyl group, 2,4-hexadiyne-1,6-diyl group, 2-heptyne-1,7-diyl group, 3-heptyne-1,7-diyl group, 4-heptyne-1,7-diyl group, 5-heptyne-1,7-diyl group, 2,4-heptadiyne-1,7-diyl group, 2,5-heptadiyne-1,7-diyl group, 3,5-heptadiyne-1,7-diyl group, 2-octyne-1,8-diyl group, 3-
 15 octyne-1,8-diyl group, 4-octyne-1,8-diyl group, 5-octyne-1,8-diyl group, 6-octyne-1,8-diyl group, 2,4-octadiyne-1,8-diyl group, 2,5-octadiyne-1,8-diyl group, 2,6-octadiyne-1,8-diyl group, 2,4,6-octatriyne-1,8-diyl group, 2-nonyne-1,9-diyl group, 3-nonyne-1,9-diyl group, 4-nonyne-1,9-diyl
 20 group, 5-nonyne-1,9-diyl group, 6-nonyne-1,9-diyl group, 7-nonyne-1,9-diyl group, 2-decyne-1,10-diyl group, 3-decyne-1,10-diyl group, 4-decyne-1,10-diyl group, 5-decyne-1,10-diyl group, 6-decyne-1,10-diyl group, 7-decyne-1,10-diyl group, 8-decyne-1,10-diyl group;
 25 2-undecyne-1,11-diyl group, 3-undecyne-1,11-diyl group, 4-undecyne-1,11-diyl group, 5-undecyne-1,11-diyl group, 6-undecyne-1,11-diyl group, 7-undecyne-1,11-diyl group, 8-undecyne-1,11-diyl group, 9-undecyne-1,11-diyl group;

- 2-dodecyne-1,12-diyl group, 3-dodecyne-1,12-diyl group, 4-dodecyne-1,12-diyl group, 5-dodecyne-1,12-diyl group, 6-dodecyne-1,12-diyl group, 7-dodecyne-1,12-diyl group, 8-dodecyne-1,12-diyl group, 9-dodecyne-1,12-diyl group, 10-dodecyne-1,12-diyl group;
- 2-tridecyne-1,13-diyl group, 3-tridecyne-1,13-diyl group, 4-tridecyne-1,13-diyl group, 5-tridecyne-1,13-diyl group, 6-tridecyne-1,13-diyl group, 7-tridecyne-1,13-diyl group, 8-tridecyne-1,13-diyl group, 9-tridecyne-1,13-diyl group, 10-tridecyne-1,13-diyl group, 11-tridecyne-1,13-diyl group;
- 2-tetradecyne-1,14-diyl group, 3-tetradecyne-1,14-diyl group, 4-tetradecyne-1,14-diyl group, 5-tetradecyne-1,14-diyl group, 6-tetradecyne-1,14-diyl group, 7-tetradecyne-1,14-diyl group, 8-tetradecyne-1,14-diyl group, 9-tetradecyne-1,14-diyl group, 10-tetradecyne-1,14-diyl group, 11-tetradecyne-1,14-diyl group, 12-tetradecyne-1,14-diyl group;
- 2-pentadecyne-1,15-diyl group, 3-pentadecyne-1,15-diyl group, 4-pentadecyne-1,15-diyl group, 5-pentadecyne-1,15-diyl group, 6-pentadecyne-1,15-diyl group, 7-pentadecyne-1,15-diyl group, 8-pentadecyne-1,15-diyl group, 9-pentadecyne-1,15-diyl group, 10-pentadecyne-1,15-diyl group, 11-pentadecyne-1,15-diyl group, 12-pentadecyne-1,15-diyl group, 13-pentadecyne-1,15-diyl group;
- 2-hexadecyne-1,16-diyl group, 3-hexadecyne-1,16-diyl group, 4-hexadecyne-1,16-diyl group, 5-hexadecyne-1,16-diyl group, 6-hexadecyne-1,16-diyl group, 7-hexadecyne-1,16-diyl group, 8-hexadecyne-1,16-diyl group, 9-hexadecyne-1,16-diyl group,

- 10-hexadecyne-1,16-diyl group, 11-hexadecyne-1,16-diyl group, 12-hexadecyne-1,16-diyl group, 13-hexadecyne-1,16-diyl group, 14-hexadecyne-1,16-diyl group;
- 2-heptadecyne-1,17-diyl group, 3-heptadecyne-1,17-diyl group, 4-heptadecyne-1,17-diyl group, 5-heptadecyne-1,17-diyl group, 6-heptadecyne-1,17-diyl group, 7-heptadecyne-1,17-diyl group, 8-heptadecyne-1,17-diyl group, 9-heptadecyne-1,17-diyl group, 10-heptadecyne-1,17-diyl group, 11-heptadecyne-1,17-diyl group, 12-heptadecyne-1,17-diyl group, 13-heptadecyne-1,17-diyl group, 14-heptadecyne-1,17-diyl group, 15-heptadecyne-1,17-diyl group;
- 2-octadecyne-1,18-diyl group, 3-octadecyne-1,18-diyl group, 4-octadecyne-1,18-diyl group, 5-octadecyne-1,18-diyl group, 6-octadecyne-1,18-diyl group, 7-octadecyne-1,18-diyl group, 8-octadecyne-1,18-diyl group, 9-octadecyne-1,18-diyl group, 10-octadecyne-1,18-diyl group, 11-octadecyne-1,18-diyl group, 12-octadecyne-1,18-diyl group, 13-octadecyne-1,18-diyl group, 14-octadecyne-1,18-diyl group, 15-octadecyne-1,18-diyl group, 16-octadecyne-1,18-diyl group;
- 2-nonadecyne-1,19-diyl group, 3-nonadecyne-1,19-diyl group, 4-nonadecyne-1,19-diyl group, 5-nonadecyne-1,19-diyl group, 6-nonadecyne-1,19-diyl group, 7-nonadecyne-1,19-diyl group, 8-nonadecyne-1,19-diyl group, 9-nonadecyne-1,19-diyl group, 10-nonadecyne-1,19-diyl group, 11-nonadecyne-1,19-diyl group, 12-nonadecyne-1,19-diyl group, 13-nonadecyne-1,19-diyl group, 14-nonadecyne-1,19-diyl group, 15-nonadecyne-1,19-diyl group, 16-nonadecyne-1,19-diyl group, 17-nonadecyne-1,19-diyl group;

- 2-icosyne-1,20-diyl group, 3-icosyne-1,20-diyl group, 4-icosyne-1,20-diyl group, 5-icosyne-1,20-diyl group, 6-icosyne-1,20-diyl group, 7-icosyne-1,20-diyl group, 8-icosyne-1,20-diyl group, 9-icosyne-1,20-diyl group, 10-icosyne-1,20-diyl group, 11-icosyne-1,20-diyl group, 12-icosyne-1,20-diyl group, 13-icosyne-1,20-diyl group, 14-icosyne-1,20-diyl group, 15-icosyne-1,20-diyl group, 16-icosyne-1,20-diyl group, 17-icosyne-1,20-diyl group, 18-icosyne-1,20-diyl group;
- 10 2-henicosyne-1,21-diyl group, 3-henicosyne-1,21-diyl group, 4-henicosyne-1,21-diyl group, 5-henicosyne-1,21-diyl group, 6-henicosyne-1,21-diyl group, 7-henicosyne-1,21-diyl group, 8-henicosyne-1,21-diyl group, 9-henicosyne-1,21-diyl group, 10-henicosyne-1,21-diyl group, 11-henicosyne-1,21-diyl group, 12-henicosyne-1,21-diyl group, 13-henicosyne-1,21-diyl group, 14-henicosyne-1,21-diyl group, 15-henicosyne-1,21-diyl group, 16-henicosyne-1,21-diyl group, 17-henicosyne-1,21-diyl group, 18-henicosyne-1,21-diyl group, 19-henicosyne-1,21-diyl group;
- 20 2-docosyne-1,22-diyl group, 3-docosyne-1,22-diyl group, 4-docosyne-1,22-diyl group, 5-docosyne-1,22-diyl group, 6-docosyne-1,22-diyl group, 7-docosyne-1,22-diyl group, 8-docosyne-1,22-diyl group, 9-docosyne-1,22-diyl group, 10-docosyne-1,22-diyl group, 11-docosyne-1,22-diyl group, 12-docosyne-1,22-diyl group, 13-docosyne-1,22-diyl group, 14-docosyne-1,22-diyl group, 15-docosyne-1,22-diyl group, 16-docosyne-1,22-diyl group, 17-docosyne-1,22-diyl group, 18-docosyne-1,22-diyl group, 19-docosyne-1,22-diyl group, 20-

- docosyne-1,22-diyl group;
- 2-tricosyne-1,23-diyl group, 3-tricosyne-1,23-diyl group,
 4-tricosyne-1,23-diyl group, 5-tricosyne-1,23-diyl group,
 6-tricosyne-1,23-diyl group, 7-tricosyne-1,23-diyl group,
 5 8-tricosyne-1,23-diyl group, 9-tricosyne-1,23-diyl group,
 10-tricosyne-1,23-diyl group, 11-tricosyne-1,23-diyl group,
 12-tricosyne-1,23-diyl group, 13-tricosyne-1,23-diyl group,
 14-tricosyne-1,23-diyl group, 15-tricosyne-1,23-diyl group,
 16-tricosyne-1,23-diyl group, 17-tricosyne-1,23-diyl group,
 10 18-tricosyne-1,23-diyl group, 19-tricosyne-1,23-diyl group,
 20-tricosyne-1,23-diyl group, 21-tricosyne-1,23-diyl group;
 2-tetracosyne-1,24-diyl group, 3-tetracosyne-1,24-diyl
 group, 4-tetracosyne-1,24-diyl group, 5-tetracosyne-1,24-
 diyl group, 6-tetracosyne-1,24-diyl group, 7-tetracosyne-
 15 1,24-diyl group, 8-tetracosyne-1,24-diyl group, 9-
 tetracosyne-1,24-diyl group, 10-tetracosyne-1,24-diyl group,
 11-tetracosyne-1,24-diyl group, 12-tetracosyne-1,24-diyl
 group, 13-tetracosyne-1,24-diyl group, 14-tetracosyne-1,24-
 diyl group, 15-tetracosyne-1,24-diyl group, 16-tetracosyne-
 20 1,24-diyl group, 17-tetracosyne-1,24-diyl group, 18-
 tetracosyne-1,24-diyl group, 19-tetracosyne-1,24-diyl group,
 20-tetracosyne-1,24-diyl group, 21-tetracosyne-1,24-diyl
 group, 22-tetracosyne-1,24-diyl group;
 2-pentacosyne-1,25-diyl group, 3-pentacosyne-1,25-diyl
 25 group, 4-pentacosyne-1,25-diyl group, 5-pentacosyne-1,25-
 diyl group, 6-pentacosyne-1,25-diyl group, 7-pentacosyne-
 1,25-diyl group, 8-pentacosyne-1,25-diyl group, 8-
 pentacosyne-1,25-diyl group, 9-pentacosyne-1,25-diyl group,

10-pentacosyne-1,25-diyl group, 11-pentacosyne-1,25-diyl group, 12-pentacosyne-1,25-diyl group, 13-pentacosyne-1,25-diyl group, 14-pentacosyne-1,25-diyl group, 15-pentacosyne-1,25-diyl group, 16-pentacosyne-1,25-diyl group, 17-

5 pentacosyne-1,25-diyl group, 18-pentacosyne-1,25-diyl group, 19-pentacosyne-1,25-diyl group, 20-pentacosyne-1,25-diyl group, 21-pentacosyne-1,25-diyl group, 22-pentacosyne-1,25-diyl group, 23-pentacosyne-1,25-diyl group;

2-hexacosyne-1,26-diyl group, 3-hexacosyne-1,26-diyl group,

10 4-hexacosyne-1,26-diyl group, 5-hexacosyne-1,26-diyl group, 6-hexacosyne-1,26-diyl group, 7-hexacosyne-1,26-diyl group, 8-hexacosyne-1,26-diyl group, 9-hexacosyne-1,26-diyl group, 10-hexacosyne-1,26-diyl group, 11-hexacosyne-1,26-diyl group, 12-hexacosyne-1,26-diyl group, 13-hexacosyne-1,26-

15 diyl group, 14-hexacosyne-1,26-diyl group, 15-hexacosyne-1,26-diyl group, 16-hexacosyne-1,26-diyl group, 17-hexacosyne-1,26-diyl group, 18-hexacosyne-1,26-diyl group, 19-hexacosyne-1,26-diyl group, 20-hexacosyne-1,26-diyl group, 21-hexacosyne-1,26-diyl group, 22-hexacosyne-1,26-

20 diyl group, 23-hexacosyne-1,26-diyl group, 24-hexacosyne-1,26-diyl group;

2-heptacosyne-1,27-diyl group, 3-heptacosyne-1,27-diyl group, 4-heptacosyne-1,27-diyl group, 5-heptacosyne-1,27-diyl group, 6-heptacosyne-1,27-diyl group, 7-heptacosyne-

25 1,27-diyl group, 8-heptacosyne-1,27-diyl group, 9-heptacosyne-1,27-diyl group, 10-heptacosyne-1,27-diyl group, 11-heptacosyne-1,27-diyl group, 12-heptacosyne-1,27-diyl group, 13-heptacosyne-1,27-diyl group, 14-heptacosyne-1,27-

diyl group, 15-heptacosyne-1,27-diyl group, 16-heptacosyne-
 1,27-diyl group, 17-heptacosyne-1,27-diyl group, 18-
 heptacosyne-1,27-diyl group, 19-heptacosyne-1,27-diyl group,
 20-heptacosyne-1,27-diyl group, 21-heptacosyne-1,27-diyl
 5 group, 22-heptacosyne-1,27-diyl group, 2,3-heptacosyne-
 1,27-diyl group, 24-heptacosyne-1,27-diyl group, 25-
 heptacosyne-1,27-diyl group;
 2-octacosyne-1,28-diyl group, 3-octacosyne-1,28-diyl group,
 4-octacosyne-1,28-diyl group, 5-octacosyne-1,28-diyl group,
 10 6-octacosyne-1,28-diyl group, 7-octacosyne-1,28-diyl group,
 8-octacosyne-1,28-diyl group, 9-octacosyne-1,28-diyl group,
 10-octacosyne-1,28-diyl group, 11-octacosyne-1,28-diyl
 group, 12-octacosyne-1,28-diyl group, 13-octacosyne-1,28-
 diyl group, 14-octacosyne-1,28-diyl group, 15-octacosyne-
 15 1,28-diyl group, 16-octacosyne-1,28-diyl group, 17-
 octacosyne-1,28-diyl group, 18-octacosyne-1,28-diyl group,
 19-octacosyne-1,28-diyl group, 20-octacosyne-1,28-diyl
 group, 21-octacosyne-1,28-diyl group, 22-octacosyne-1,28-
 diyl group, 23-octacosyne-1,28-diyl group, 24-octacosyne-
 20 1,28-diyl group, 25-octacosyne-1,28-diyl group, 26-
 octacosyne-1,28-diyl group;
 2-nonacosyne-1,29-diyl group, 3-nonacosyne-1,29-diyl group,
 4-nonacosyne-1,29-diyl group, 5-nonacosyne-1,29-diyl group,
 6-nonacosyne-1,29-diyl group, 7-nonacosyne-1,29-diyl group,
 25 8-nonacosyne-1,29-diyl group, 9-nonacosyne-1,29-diyl group,
 10-nonacosyne-1,29-diyl group, 11-nonacosyne-1,29-diyl
 group, 12-nonacosyne-1,29-diyl group, 13-nonacosyne-1,29-
 diyl group, 14-nonacosyne-1,29-diyl group, 15-nonacosyne-

1,29-diyl group, 16-nonacosyne-1,29-diyl group, 17-
 nonacosyne-1,29-diyl group, 18-nonacosyne-1,29-diyl group,
 19-nonacosyne-1,29-diyl group, 20-nonacosyne-1,29-diyl
 group, 21-nonacosyne-1,29-diyl group, 22-nonacosyne-1,29-
 5 diyl group, 23-nonacosyne-1,29-diyl group, 24-nonacosyne-
 1,29-diyl group, 25-nonacosyne-1,29-diyl group, 26-
 nonacosyne-1,29-diyl group, 27-nonacosyne-1,29-diyl group;
 2-triacontyne-1,30-diyl group, 3-triacontyne-1,30-diyl
 group, 4-triacontyne-1,30-diyl group, 5-triacontyne-1,30-
 10 diyl group, 6-triacontyne-1,30-diyl group, 7-triacontyne-
 1,30-diyl group, 8-triacontyne-1,30-diyl group, 9-
 triacontyne-1,30-diyl group, 10-triacontyne-1,30-diyl group,
 11-triacontyne-1,30-diyl group, 12-triacontyne-1,30-diyl
 group, 13-triacontyne-1,30-diyl group, 14-triacontyne-1,30-
 15 diyl group, 15-triacontyne-1,30-diyl group, 16-triacontyne-
 1,30-diyl group, 17-triacontyne-1,30-diyl group, 18-
 triacontyne-1,30-diyl group, 19-triacontyne-1,30-diyl group,
 20-triacontyne-1,30-diyl group, 21-triacontyne-1,30-diyl
 group, 22-triacontyne-1,30-diyl group, 23-triacontyne-1,30-
 20 diyl group, 24-triacontyne-1,30-diyl group, 25-triacontyne-
 1,30-diyl group, 26-triacontyne-1,30-diyl group, 27-
 triacontyne-1,30-diyl group, and 28-triacontyne-1,30-diyl
 group;

as well as branched alkynylene groups such as 3-methyl-1-
 25 butyne-1,4-diyl group, 2-methyl-3-butyne-1,4-diyl group, 4-
 methyl-2-pentyne-1,5-diyl group, 2-methyl-3-pentyne-1,5-
 diyl group, 4-ethyl-2-hexyne-1,6-diyl group, 5-methyl-3-
 hexyne-1,6-diyl group, 2-methyl-4-hexyne-1,6-diyl group, 5-

ethyl-6-methyl-2-heptyne-1,7-diyl group, 5-methyl-3-heptyne-1,7-diyl group, 3-n-propyl-4-heptyne-1,7-diyl group, 4,4-dimethyl-5-heptyne-1,7-diyl group, 6-methyl-2,4-heptadiyne-1,7-diyl group, 4-methyl-2,5-heptadiyne-1,7-diyl group, 2-methyl-3,5-heptadiyne-1,7-diyl group, 4-ethyl-6,6-dimethyl-2-octyne-1,8-diyl group, 5-n-propyl-3-octyne-1,8-diyl group, 3-ethyl-4-octyne-1,8-diyl group, 4-ethyl-2-methyl-5-octyne-1,8-diyl group, 3,4,5-trimethyl-6-octyne-1,8-diyl group, 7-methyl-2,4-octadiyne-1,8-diyl group, 4-methyl-2,5-octadiyne-1,8-diyl group, 5-n-propyl-2,6-octadiyne-1,8-diyl group, 5-ethyl-2-nonyne-1,9-diyl group, 5,6,7-trimethyl-3-nonyne-1,9-diyl group, 2,3,6,7-tetramethyl-4-nonyne-1,9-diyl group, 3,4-diethyl-5-nonyne-1,9-diyl group, 4-i-propyl-6-nonyne-1,9-diyl group, 3-ethyl-7-nonyne-1,9-diyl group, 5-n-butyl-2-decyne-1,10-diyl group, 6-i-propyl-3-decyne-1,10-diyl group, 7-ethyl-4-decyne-1,10-diyl group, 3,7-dimethyl-5-decyne-1,10-diyl group, 4-ethyl-6-decyne-1,10-diyl group, 5-methyl-7-decyne-1,10-diyl group, 6-ethyl-4-methyl-8-decyne-1,10-diyl group; 6-methyl-2-undecyne-1,11-diyl group, 6-ethyl-3-undecyne-1,11-diyl group, 7-methyl-4-undecyne-1,11-diyl group, 7-ethyl-5-undecyne-1,11-diyl group, 5-methyl-6-undecyne-1,11-diyl group, 9-ethyl-7-undecyne-1,11-diyl group, 3-methyl-8-undecyne-1,11-diyl group, 4-ethyl-9-undecyne-1,11-diyl group; 5-ethyl-2-dodecyne-1,12-diyl group, 6-methyl-3-dodecyne-1,12-diyl group, 8-ethyl-4-dodecyne-1,12-diyl group, 8-methyl-5-dodecyne-1,12-diyl group, 9-ethyl-6-dodecyne-1,12-

- diyl group, 6-methyl-7-dodecyne-1,12-diyl group, 10-ethyl-8-dodecyne-1,12-diyl group, 2-methyl-9-dodecyne-1,12-diyl group, 5-ethyl-10-dodecyne-1,12-diyl group, 4,7,9-trimethyl-2-tridecyne-1,13-diyl group, 10-methyl-3-tridecyne-1,13-diyl group, 8-ethyl-4-tridecyne-1,13-diyl group, 4-methyl-5-tridecyne-1,13-diyl group, 5-ethyl-6-tridecyne-1,13-diyl group, 3,6-diethyl-7-tridecyne-1,13-diyl group, 5-methyl-8-tridecyne-1,13-diyl group, 7-ethyl-9-tridecyne-1,13-diyl group, 4-methyl-10-tridecyne-1,13-diyl group, 6-ethyl-11-tridecyne-1,13-diyl group; 7-methyl-2-tetradecyne-1,14-diyl group, 8-ethyl-3-tetradecyne-1,14-diyl group, 6-n-propyl-4-tetradecyne-1,14-diyl group, 8-methyl-5-tetradecyne-1,14-diyl group, 3-ethyl-6-tetradecyne-1,14-diyl group, 10-methyl-7-tetradecyne-1,14-diyl group, 6-i-propyl-8-tetradecyne-1,14-diyl group, 5,7,11-trimethyl-9-tetradecyne-1,14-diyl group, 5-ethyl-10-tetradecyne-1,14-diyl group, 6-methyl-11-tetradecyne-1,14-diyl group, 4-n-butyl-12-tetradecyne-1,14-diyl group;
- 4-methyl-2-pentadecyne-1,15-diyl group, 6-ethyl-3-pentadecyne-1,15-diyl group, 8-methyl-4-pentadecyne-1,15-diyl group, 10-ethyl-5-pentadecyne-1,15-diyl group, 4,9-dimethyl-6-pentadecyne-1,15-diyl group, 10-ethyl-7-pentadecyne-1,15-diyl group, 6-methyl-8-pentadecyne-1,15-diyl group, 8-n-propyl-9-pentadecyne-1,15-diyl group, 5-methyl-10-pentadecyne-1,15-diyl group, 4,7-diethyl-11-pentadecyne-1,15-diyl group, 5-methyl-12-pentadecyne-1,15-diyl group, 8-ethyl-13-pentadecyne-1,15-diyl group;

- 8-i-propyl-2-hexadecyne-1,16-diyl group, 6-methyl-3-hexadecyne-1,16-diyl group, 8-ethyl-4-hexadecyne-1,16-diyl group, 9-methyl-5-hexadecyne-1,16-diyl group, 10-ethyl-6-hexadecyne-1,16-diyl group, 5-methyl-7-hexadecyne-1,16-diyl group, 5,11-dimethyl-8-hexadecyne-1,16-diyl group, 5-ethyl-9-hexadecyne-1,16-diyl group, 7,13-diethyl-10-hexadecyne-1,16-diyl group, 5-ethyl-7-methyl-11-hexadecyne-1,16-diyl group, 5-methyl-12-hexadecyne-1,16-diyl group, 8-s-butyl-13-hexadecyne-1,16-diyl group, 5-ethyl-14-hexadecyne-1,16-diyl group;
- 11-methyl-2-heptadecyne-1,17-diyl group, 9-ethyl-3-heptadecyne-1,17-diyl group, 7-i-propyl-4-heptadecyne-1,17-diyl group, 8-methyl-5-heptadecyne-1,17-diyl group, 4-ethyl-6-heptadecyne-1,17-diyl group, 10-methyl-7-heptadecyne-1,17-diyl group, 5,11-dimethyl-8-heptadecyne-1,17-diyl group, 5-ethyl-9-heptadecyne-1,17-diyl group, 8-ethyl-10-heptadecyne-1,17-diyl group, 7-methyl-11-heptadecyne-1,17-diyl group, 5-i-propyl-12-heptadecyne-1,17-diyl group, 9-ethyl-13-heptadecyne-1,17-diyl group, 8-methyl-14-heptadecyne-1,17-diyl group, 7-s-butyl-15-heptadecyne-1,17-diyl group;
- 10,15-dimethyl-2-octadecyne-1,18-diyl group, 6-ethyl-3-octadecyne-1,18-diyl group, 10-methyl-4-octadecyne-1,18-diyl group, 11-methyl-5-octadecyne-1,18-diyl group, 12-ethyl-6-octadecyne-1,18-diyl group, 10-methyl-7-octadecyne-1,18-diyl group, 5-methyl-8-octadecyne-1,18-diyl group, 7-ethyl-9-octadecyne-1,18-diyl group, 7-methyl-10-octadecyne-1,18-diyl group, 8-n-butyl-11-octadecyne-1,18-diyl group,

- 7-methyl-12-octadecyne-1,18-diyl group, 9-ethyl-13-octadecyne-1,18-diyl group, 10-i-propyl-14-octadecyne-1,18-diyl group, 7-methyl-15-octadecyne-1,18-diyl group, 10-ethyl-16-octadecyne-1,18-diyl group;
- 5 10-methyl-2-nonadecyne-1,19-diyl group, 10,12-diethyl-3-nonadecyne-1,19-diyl group, 7-methyl-4-nonadecyne-1,19-diyl group, 9-ethyl-5-nonadecyne-1,19-diyl group, 9-n-propyl-6-nonadecyne-1,19-diyl group, 10-methyl-7-nonadecyne-1,19-diyl group, 12-i-propyl-8-nonadecyne-1,19-diyl group, 5,15-
- 10 dimethyl-9-nonadecyne-1,19-diyl group, 7-ethyl-13-methyl-10-nonadecyne-1,19-diyl group, 6-methyl-11-nonadecyne-1,19-diyl group, 6-ethyl-12-nonadecyne-1,19-diyl group, 7,16-diethyl-13-nonadecyne-1,19-diyl group, 9-s-butyl-14-nonadecyne-1,19-diyl group, 8-methyl-15-nonadecyne-1,19-
- 15 diyl group, 10-ethyl-16-nonadecyne-1,19-diyl group, 10-i-propyl-17-nonadecyne-1,19-diyl group;
- 8-methyl-2-icosyne-1,20-diyl group, 6-ethyl-3-icosyne-1,20-diyl group, 10-i-propyl-4-icosyne-1,20-diyl group, 11-n-propoyl-5-icosyne-1,20-diyl group, 12-methyl-6-icosyne-
- 20 1,20-diyl group, 11-ethyl-7-icosyne-1,20-diyl group, 13-n-propyl-8-icosyne-1,20-diyl group, 6-i-propyl-9-icosyne-1,20-diyl group, 5-n-propyl-10-icosyne-1,20-diyl group, 7-methyl-11-icosyne-1,20-diyl group, 8-ethyl-12-icosyne-1,20-diyl group, 10-n-propyl-13-icosyne-1,20-diyl group, 9-i-
- 25 propyl-14-icosyne-1,20-diyl group, 10-n-butyl-15-icosyne-1,20-diyl group, 8-s-butyl-16-icosyne-1,20-diyl group, 7-i-butyl-17-icosyne-1,20-diyl group, 9-methyl-18-icosyne-1,20-diyl group;

- 11-methyl-2-henicosyne-1,21-diyl group, 12-n-butyl-3-henicosyne-1,21-diyl group, 10-n-pentyl-4-henicosyne-1,21-diyl group, 8-ethyl-5-henicosyne-1,21-diyl group, 10-i-propyl-6-henicosyne-1,21-diyl group, 5-n-propyl-7-henicosyne-1,21-diyl group, 13-n-butyl-8-henicosyne-1,21-diyl group, 15-s-butyl-9-henicosyne-1,21-diyl group, 5-methyl-10-henicosyne-1,21-diyl group, 15-ethyl-6-methyl-11-henicosyne-1,21-diyl group, 8-ethyl-12-henicosyne-1,21-diyl group, 7-methyl-13-henicosyne-1,21-diyl group, 11-ethyl-14-henicosyne-1,21-diyl group, 6-ethyl-15-henicosyne-1,21-diyl group, 9-methyl-16-henicosyne-1,21-diyl group, 5-ethyl-9-methyl-17-henicosyne-1,21-diyl group, 10,10-dimethyl-18-henicosyne-1,21-diyl group, 9-ethyl-19-henicosyne-1,21-diyl group;
- 11-methyl-2-docosyne-1,22-diyl group, 12-ethyl-3-docosyne-1,22-diyl group, 13-i-propyl-4-docosyne-1,22-diyl group, 10-n-propyl-5-docosyne-1,22-diyl group, 10-n-butyl-6-docosyne-1,22-diyl group, 15-s-butyl-7-docosyne-1,22-diyl group, 11-i-butyl-8-docosyne-1,22-diyl group, 5,15-dimethyl-9-docosyne-1,22-diyl group, 8,14-diethyl-10-docosyne-1,22-diyl group, 5-methyl-11-docosyne-1,22-diyl group, 7-ethyl-12-docosyne-1,22-diyl group, 10-methyl-13-docosyne-1,22-diyl group, 10-ethyl-14-docosyne-1,22-diyl group, 9-ethyl-15-docosyne-1,22-diyl group, 8-methyl-16-docosyne-1,22-diyl group, 7-i-propyl-17-docosyne-1,22-diyl group, 10-i-butyl-18-docosyne-1,22-diyl group, 9,10-dimethyl-19-docosyne-1,22-diyl group, 13-ethyl-20-docosyne-1,22-diyl group;

- 19-methyl-2-tricosyne-1,23-diyl group, 10,15-dimethyl-3-tricosyne-1,23-diyl group, 3,11,16-trimethyl-4-tricosyne-1,23-diyl group, 12-ethyl-5-tricosyne-1,23-diyl group, 6,13-diethyl-6-tricosyne-1,23-diyl group, 4,12,18-triethyl-
- 5 7-tricosyne-1,23-diyl group, 18-i-propyl-8-tricosyne-1,23-diyl group, 14-n-propyl-9-tricosyne-1,23-diyl group, 8-n-butyl-10-tricosyne-1,23-diyl group, 15-s-butyl-11-tricosyne-1,23-diyl group, 5-i-butyl-12-tricosyne-1,23-diyl group, 7-ethyl-9-methyl-13-tricosyne-1,23-diyl group, 9-
- 10 methyl-14-tricosyne-1,23-diyl group, 4,18-dimethyl-15-tricosyne-1,23-diyl group, 3,4,11-trimethyl-16-tricosyne-1,23-diyl group, 9-ethyl-17-tricosyne-1,23-diyl group, 10,13-diethyl-18-tricosyne-1,23-diyl group, 5,8,15-triethyl-19-tricosyne-1,23-diyl group, 15-i-propyl-20-
- 15 tricosyne-1,23-diyl group, 17-n-propyl-21-tricosyne-1,23-diyl group;
- 16-n-butyl-2-tetracosyne-1,24-diyl group, 11-s-butyl-3-tetracosyne-1,24-diyl group, 8-i-butyl-4-tetracosyne-1,24-diyl group, 18-ethyl-9-methyl-5-tetracosyne-1,24-diyl group,
- 20 13-methyl-6-tetracosyne-1,24-diyl group, 4,19-dimethyl-7-tetracosyne-1,24-diyl group, 5,11,17-triethyl-8-tetracosyne-1,24-diyl group, 6-ethyl-9-tetracosyne-1,24-diyl group, 7,16-diethyl-10-tetracosyne-1,24-diyl group, 5,9,18-triethyl-11-tetracosyne-1,24-diyl group, 10-n-
- 25 propyl-12-tetracosyne-1,24-diyl group, 20-i-propyl-13-tetracosyne-1,24-diyl group, 9-n-butyl-14-tetracosyne-1,24-diyl group, 11-s-butyl-15-tetracosyne-1,24-diyl group, 13-i-butyl-16-tetracosyne-1,24-diyl group, 10-ethyl-13-methyl-

- 17-tetracosyne-1,24-diyl group, 6-methyl-18-tetracosyne-1,24-diyl group, 5,7-dimethyl-19-tetracosyne-1,24-diyl group, 4,8,13-trimethyl-20-tetracosyne-1,24-diyl group, 18-ethyl-21-tetracosyne-1,24-diyl group, 6,10-diethyl-22-tetracosyne-1,24-diyl group;
- 5 9,13,16-trimethyl-2-pentacosyne-1,25-diyl group, 12-n-propyl-3-pentacosyne-1,25-diyl group, 11-i-propyl-4-pentacosyne-1,25-diyl group, 20-n-butyl-5-pentacosyne-1,25-diyl group, 17-i-butyl-6-pentacosyne-1,25-diyl group, 15-s-
- 10 butyl-7-pentacosyne-1,25-diyl group, 15-ethyl-23-methyl-8-pentacosyne-1,25-diyl group, 11-methyl-8-pentacosyne-1,25-diyl group, 13,17-dimethyl-9-pentacosyne-1,25-diyl group, 5,8,21-trimethyl-10-pentacosyne-1,25-diyl group, 17-ethyl-11-pentacosyne-1,25-diyl group, 8,18-diethyl-12-
- 15 pentacosyne-1,25-diyl group, 10,15,18-trimethyl-13-pentacosyne-1,25-diyl group, 4-n-propyl-14-pentacosyne-1,25-diyl group, 20-i-propyl-15-pentacosyne-1,25-diyl group, 8-n-butyl-16-pentacosyne-1,25-diyl group, 11-s-butyl-17-pentacosyne-1,25-diyl group, 5,22-dimethyl-18-pentacosyne-
- 20 1,25-diyl group, 5-i-butyl-19-pentacosyne-1,25-diyl group, 9-methyl-13-ethyl-20-pentacosyne-1,25-diyl group, 15-methyl-21-pentacosyne-1,25-diyl group, 6,13-dimethyl-22-pentacosyne-1,25-diyl group, 4,8,12-trimethyl-23-pentacosyne-1,25-diyl group;
- 25 13-ethyl-2-hexacosyne-1,26-diyl group, 5,16-diethyl-3-hexacosyne-1,26-diyl group, 7,11,16-trimethyl-4-hexacosyne-1,26-diyl group, 12-n-propyl-5-hexacosyne-1,26-diyl group, 21-i-propyl-6-hexacosyne-1,26-diyl group, 6-n-butyl-7-

- hexacosyne-1,26-diyl group, 13-s-butyl-8-hexacosyne-1,26-diyl group, 19-i-butyl-9-hexacosyne-1,26-diyl group, 13-ethyl-18-methyl-10-hexacosyne-1,26-diyl group, 10-methyl-11-hexacosyne-1,26-diyl group, 10,20-dimethyl-12-
- 5 hexacosyne-1,26-diyl group, 7,9,17-trimethyl-13-hexacosyne-1,26-diyl group, 8-ethyl-14-hexacosyne-1,26-diyl group, 5,22-diethyl-15-hexacosyne-1,26-diyl group, 7,10,21-trimethyl-16-hexacosyne-1,26-diyl group, 15-n-propyl-17-hexacosyne-1,26-diyl group, 13-i-propyl-18-hexacosyne-1,26-
- 10 diyl group, 8-n-butyl-19-hexacosyne-1,26-diyl group, 11-s-butyl-20-hexacosyne-1,26-diyl group, 14-i-butyl-21-hexacosyne-1,26-diyl group, 5-ethyl-21-methyl-22-hexacosyne-1,26-diyl group, 7-methyl-23-hexacosyne-1,26-diyl group, 8,14-dimethyl-24-hexacosyne-1,26-diyl group;
- 15 7,16,24-trimethyl-2-heptacosyne-1,27-diyl group, 9-ethyl-3-heptacosyne-1,27-diyl group, 7,16-dimethyl-4-heptacosyne-1,27-diyl group, 9,13,21-trimethyl-5-heptacosyne-1,27-diyl group, 13-n-propyl-6-heptacosyne-1,27-diyl group, 10-i-propyl-7-heptacosyne-1,27-diyl group, 16-n-propyl-8-
- 20 heptacosyne-1,27-diyl group, 18-methyl-9-heptacosyne-1,27-diyl group, 9-i-propyl-10-heptacosyne-1,27-diyl group, 15-ethyl-7-methyl-11-heptacosyne-1,27-diyl group, 25-methyl-12-heptacosyne-1,27-heptacosyne-1,27-diyl group, 8,21-dimethyl-13-heptacosyne-1,27-diyl group, 5,11,23-trimethyl-
- 25 14-heptacosyne-1,27-diyl group, 9-ethyl-15-heptacosyne-1,27-diyl group, 8,20-dimethyl-16-heptacosyne-1,27-diyl group, 4,8,19-trimethyl-17-heptacosyne-1,27-diyl group, 7-n-propyl-18-heptacosyne-1,27-diyl group, 21-i-propyl-19-

- heptacosyne-1,27-diyl group, 14-n-propyl-20-heptacosyne-1,27-diyl group, 8-ethyl-21-heptacosyne-1,27-diyl group, 11-i-propyl-22-heptacosyne-1,27-diyl group, 5-ethyl-13-methyl-23-heptacosyne-1,27-diyl group, 16-methyl-24-
- 5 heptacosyne-1,27-diyl group, 7-ethyl-25-heptacosyne-1,27-diyl group;
- 14-ethyl-2-octacosyne-1,28-diyl group, 20-methyl-3-octacosyne-1,28-diyl group, 7,22-dimethyl-4-octacosyne-1,28-diyl group, 19-ethyl-5-octacosyne-1,28-diyl group, 11-
- 10 methyl-6-octacosyne-1,28-diyl group, 13,16-dimethyl-7-octacosyne-1,28-diyl group, 13-ethyl-8-octacosyne-1,28-diyl group, 6-methyl-9-octacosyne-1,28-diyl group, 9,16-dimethyl-10-octacosyne-1,28-diyl group, 7-ethyl-11-octacosyne-1,28-diyl group, 16-methyl-12-octacosyne-1,28-
- 15 diyl group, 6,15-dimethyl-13-octacosyne-1,28-diyl group, 22-ethyl-14-octacosyne-1,28-diyl group, 6-methyl-15-octacosyne-1,28-diyl group, 8,11-dimethyl-16-octacosyne-1,28-diyl group, 23-ethyl-17-octacosyne-1,28-diyl group, 4-methyl-18-octacosyne-1,28-diyl group, 7,14-dimethyl-19-
- 20 octacosyne-1,28-diyl group, 13-ethyl-20-octacosyne-1,28-diyl group, 8-methyl-21-octacosyne-1,28-diyl group, 11,17-dimethyl-22-octacosyne-1,28-diyl group, 10-ethyl-23-octacosyne-1,28-diyl group, 9-methyl-24-octacosyne-1,28-diyl group, 7,19-dimethyl-25-octacosyne-1,28-diyl group,
- 25 12-ethyl-26-octacosyne-1,28-diyl group;
- 15-methyl-2-nonacosyne-1,29-diyl group, 14-methyl-3-nonacosyne-1,29-diyl group, 12-methyl-4-nonacosyne-1,29-diyl group, 13-methyl-5-nonacosyne-1,29-diyl group, 11-

methyl-6-nonacosyne-1,29-diyl group, 10-methyl-7-
 nonacosyne-1,29-diyl group, 25-methyl-8-nonacosyne-1,29-
 diyl group, 24-methyl-9-nonacosyne-1,29-diyl group, 23-
 methyl-10-nonacosyne-1,29-diyl group, 22-methyl-11-
 5 nonacosyne-1,29-diyl group, 21-methyl-12-nonacosyne-1,29-
 diyl group, 20-methyl-13-nonacosyne-1,29-diyl group, 19-
 methyl-14-nonacosyne-1,29-diyl group, 18-methyl-15-
 nonacosyne-1,29-diyl group, 27-methyl-16-nonacosyne-1,29-
 diyl group, 26-methyl-17-nonacosyne-1,29-diyl group, 25-
 10 methyl-18-nonacosyne-1,29-diyl group, 24-methyl-19-
 nonacosyne-1,29-diyl group, 23-methyl-20-nonacosyne-1,29-
 diyl group, 20-methyl-21-nonacosyne-1,29-diyl group, 19-
 methyl-22-nonacosyne-1,29-diyl group, 18-methyl-23-
 nonacosyne-1,29-diyl group, 17-methyl-24-nonacosyne-1,29-
 15 diyl group, 16-methyl-25-nonacosyne-1,29-diyl group, 6-
 methyl-26-nonacosyne-1,29-diyl group, and 5-methyl-27-
 nonacosyne-1,29-diyl group.

Typically, optionally substituted straight-chained
 alkylene groups having 1 - 30 carbon atoms are preferred as
 20 G; optionally substituted straight-chained groups having 2
 - 15 carbon atoms are more preferred and straight-chained
 alkylene groups having 2 - 13 carbon atoms that may
 optionally be substituted by a hydroxyl group are further
 preferred; among these, ethane-1,2-diyl group, propane-1,3-
 25 diyl group, butane-1,4-diyl group, pentane-1,5-diyl group,
 hexane-1,6-diyl group, heptane-1,7-diyl group, octane-1,8-
 diyl group, nonane-1,9-diyl group, decane-1,10-diyl group,
 undecane-1,11-diyl group, dodecane-1,12-diyl group,

tridecane-1,13-diyl group, 2-hydroxypropane-1,3-diyl group, 3-hydroxy-octane-1,8-diyl group, 3-hydroxynonane-1,9-diyl group, 3-hydroxydecane-1,10-diyl group and the like are particularly preferred.

- 5 The optionally substituted straight-chained or branched alkylene group having 1 - 30 carbon atoms, the optionally substituted straight-chained or branched alkenylene group having 2 - 30 carbon atoms and the optionally substituted straight-chained or branched
- 10 alkynylene group having 2 - 30 carbon atoms, all being listed as candidates for G, are such that they bind to A in their 1-position and bind to E in their ω position or bind to E in their 1-position and bind to A in their ω position; preferably, they bind to A in their 1-position and bind to
- 15 E in their ω position.

E represents a single bond or -O- and preferably represents a single bond.

- J represents a single bond, an optionally substituted aromatic hydrocarbon group or an optionally substituted
- 20 heterocyclic group, with a single bond and an aromatic hydrocarbon group being preferred, and a single bond being more preferred.

- If J is an optionally substituted aromatic hydrocarbon group or an optionally substituted heterocyclic group,
- 25 exemplary substituents include $-(CH_2)_k-COOR^{7b}$, $-(CH_2)_1-$, $CONR^{8c}R^{9c}$, $-NR^{8d}R^{9d}$, hydroxyl group, etc. Here, k and 1 represent independently 0 or 1; R^{7b} represents a hydrogen atom or a straight-chained or branched alkyl group having 1

- 6 carbon atoms; R^{8c} , R^{9c} , R^{8c} and R^{9d} represent each independently a hydrogen atom or a straight-chained or branched alkyl group having 1 - 3 carbon atoms. Except in the case where Q is a single bond, these substituents are preferably absent and in the case where Q is a single bond,
 5 a preferred substituent is $-(CH_2)_k-COOR^{7b}$ (where k and R^{7b} have the same meanings as defined above). In the case where J is substituted, the number of substituents is from one to four, preferably one.

10 If J is an optionally substituted aromatic hydrocarbon group, the definition of the aromatic hydrocarbon group is the same as given for the aromatic hydrocarbon group in the case where it is used as Ar; preferred examples include p-phenylene group and m-phenylene group.

15 If J is an optionally substituted aromatic hydrocarbon group, preferred examples are unsubstituted p-phenylene group, unsubstituted m-phenylene group and -COOH substituted phenylene group.

If J is an optionally substituted heterocyclic group,
 20 the heterocyclic group means a 4-membered to 10-membered monocyclic or fused aliphatic or aromatic ring containing 1 - 4 hetero atoms which may be the same or different and are exemplified by an oxygen atom, a nitrogen atom and a sulfur atom; specific examples include oxetane, furan,
 25 dihydrofuran, tetrahydrofuran, pyran, dihydropyran, tetrahydropyran, dioxole, thiophene, dihydrothiophene, tetrahydrothiophene, thiopyran, dihydrothiopyran, tetrahydrothiopyran, pyrrole, dihydropyrrole, pyrrolidine,

pyridine, dihydropyridine, tetrahydropyridine, piperidine, pyrazole, 2-pyrazoline, pyrazolidine, imidazole, imidazolidine, pyrimidine, pyrazine, pyridazine, oxazoline, piperazine, 1,2,3-triazole, 1,2,4-triazole, tetrazole, 5 isoxazole, 1,3-oxazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2-thiazole, 1,3-thiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,3-dioxolan, 1,4-dioxane, oxazolidine, morpholine, indole, quinoline, isoquinoline, 10 benzopyran, benzofuran, benzothiophene, benzodiazole, benzoxazole and benzothiazole. Preferred examples include furan and oxazole. The heterocyclic group as J means a group having one bond each in two different positions in these hetero rings other than the positions having 15 substituents; preferred examples include furan-2,5-diyl group, 1,3-oxazole-2,4-diyl group and 1,3-oxazole-2,5-diyl group.

Preferred examples of the optionally substituted heterocyclic group as J include unsubstituted furan-2,5- 20 diyl group, unsubstituted 1,3-oxazole-2,4-diyl group and unsubstituted 1,3-oxazole-2,5-diyl group.

If J is an optionally substituted aromatic hydrocarbon group or an optionally substituted heterocyclic group, it may be bound to E via any of the two bonds as long as it is 25 bound to E via one bond and bound to Y via the other; preferably, J is bound to E in 4-position if it is 1,3-oxazole-2,4-diyl group and bound to E in 5-position if it is 1,3-oxazole-2,5-diyl group.

Y represents a single bond or -O- and preferably represents a single bond.

























L represents a single bond, a straight-chained or branched alkylene group having 1 - 10 carbon atoms, a
5 straight-chained or branched alkenylene group having 2 - 10 carbon atoms or a straight-chained or branched alkynylene group having 2 - 10 carbon atoms and a single bond is preferred; if J is an optionally substituted aromatic hydrocarbon group and Y is a single bond, L is preferably a
10 single bond or a straight-chained alkylene group having 1 - 5 carbon atoms, among which a single bond and a straight-chained alkylene group having 1 - 3 carbon atoms are preferred, with a single bond and propane-1,3-dily group being particularly preferred; if J is an optionally
15 substituted aromatic hydrocarbon group and Y is -O-, L is preferably one of straight-chained alkylene groups having 1 - 5 carbon atoms, among which straight-chained alkylene groups having 2 - 4 carbon atoms are preferred.





















L represents a straight-chained or branched alkylene
20 group having 1 - 10 carbon atoms, a straight-chained or branched alkenylene group having 2 - 10 carbon atoms, a straight-chained or branched alkynylene group having 2 - 10 carbon atoms, a straight-chained alkylene group having 1 - 5 carbon atoms, a straight-chained alkylene group having 1
25 - 3 carbon atoms or a straight-chained alkylene group having 2 - 4 carbon atoms; specific examples of these groups can appropriately be selected from the list of specific examples of G which represents an optionally

substituted straight-chained or branched alkylene group having 1 - 30 carbon atoms, an optionally substituted straight-chained or branched alkenylene group having 2 - 30 carbon atoms or an optionally substituted straight-chained
5 or branched alkynylene group having 2 - 30 carbon atoms, except that methylene group is added to the list.


Any of the two bonds of L may be bound to Y as long as it satisfies the condition that it is bound to Y via one bond and bound to Q via the other.

10 Q represents a single bond or one group selected from among the following formulae:





and



(where R^7 represents a hydrogen atom or a straight-chained or branched lower alkyl group having 1 - 6 carbon atoms,
 5 and R^8 R^9 , R^{10} and R^{11} represent each independently a
 hydrogen atom or a straight-chained or branched lower alkyl

group having 1 - 3 carbon atoms); Q is preferably Q^2 {where examples of Q^2 include a single bond, Q^{62} , Q^{63} , Q^{64} , Q^3 (where R^8 has the same meaning as defined above), Q^4 (where R^8 has the same meaning as defined above), Q^{17} (where R^7 has the same meaning as defined above), Q^{32} (where R^7 has the same meaning as defined above) and Q^{27} (where R^7 has the same meaning as defined above)}; considering the strength of antiandrogenic activity, cases where Q is Q^{62} , Q^{63} , Q^{64} and Q^3 , as well as Q^4 where R^8 is a hydrogen atom are more preferred, with Q^{62} , Q^{63} , Q^{64} and Q^3 being particularly preferred. If Q is Q^3 , the nitrogen atom and R^8 in Q^3 may preferably combine with Z to form a heterocyclic group. Considering peroral absorption, Q is more preferably Q^{17} where R^7 is a hydrogen atom, or Q^{32} where R^7 is a hydrogen atom, or Q^{27} where R^7 is a hydrogen atom. If Z is -COOH, Q may preferably be a single bond considering peroral absorption.

Further referring to Q, it is bound to L in the position marked with * and bound to Z in the position marked with **.

Z represents a hydrogen atom, an optionally substituted straight-chained or branched alkyl group having 1 - 10 carbon atoms, a straight-chained or branched alkenyl group having 2 - 10 carbon atoms that may optionally be substituted by a cycloalkyl group having 3 - 6 carbon atoms or a halogen atom, a straight-chained or branched alkynyl group having 2 - 10 carbon atoms that may optionally be

substituted by a halogen atom, $-O-R^d$ (where R^d represents a hydrogen atom or a protective group of a hydroxyl group), or $-COOH$.

If Z is an optionally substituted straight-chained or
5 branched alkyl group having 1 - 10 carbon atoms, exemplary
substituents include a halogen atom, a cycloalkyl group, a
phenyl group optionally substituted by a straight-chained
or branched alkyl group, a heterocyclic group, and a
hydroxyl group. Said heterocyclic group may be exemplified
10 by a furyl group. Examples of said halogen atom include a
fluorine atom, a chlorine atom, a bromine atom and an
iodine atom, with a fluorine atom being preferred. If said
optionally substituted straight-chained or branched alkyl
group having 1 - 10 carbon atoms is substituted by a
15 halogen atom, the number of substituent halogen atoms
ranges from one to ten, preferably from three to nine, and
substitution by five halogen atoms is particularly
preferred. In a preferred mode of substitution, all
hydrogen atoms on a certain carbon atom are substituted by
20 halogen atoms (as in the cases of trihalomethyl group,
1,1,3,3,3-pentahalopropyl group, etc.)

If Z is a straight-chained or branched alkenyl group
having 2 - 10 carbon atoms that may optionally be
substituted by a halogen atom or a straight-chained or
25 branched alkynyl group having 2 - 10 carbon atoms that may
optionally be substituted by a halogen atom, exemplary
halogen atoms include a fluorine atom, a chlorine atom, a
bromine atom and an iodine atom, with a fluorine atom being

preferred. The number of substituent halogen atoms ranges from one to ten, preferably from three to nine, and substitution by five halogen atoms is particularly preferred. In a preferred mode of substitution, all
5 hydrogen atoms on a certain carbon atom are substituted by halogen atoms.

If Z is an optionally substituted straight-chained or branched alkyl group having 1 - 10 carbon atoms, exemplary
10 straight-chained or branched alkyl groups having 1 - 10 carbon atoms include straight-chained alkyl groups, i.e., methyl group, ethyl group, n-propyl group, n-butyl group, n-pentyl group, n-hexyl group, n-heptyl group, n-octyl group, n-nonyl group and n-decyl group, as well as branched alkyl groups such as 1-methylethyl group, 1-methylpropyl
15 group, 2-methylpropyl group, 1-methylbutyl group, 2-methylbutyl group, 3-methylbutyl group, 1,1-dimethylpropyl group, 1,2-dimethylpropyl group, 2,2-dimethylpropyl group, 1-ethylpropyl group, 1-methylpentyl group, 2-methylpentyl group, 3-methylpentyl group, 4-methylpentyl group, 1,1-
20 dimethylbutyl group, 1,2-dimethylbutyl group, 1,3-dimethylbutyl group, 2,2-dimethylbutyl group, 2,3-dimethylbutyl group, 3,3-dimethylbutyl group, 1-ethylbutyl group, 2-ethylbutyl group, 1-methylhexyl group, 2-methylhexyl group, 3-methylhexyl group, 4-methylhexyl group,
25 5-methylhexyl group, 1-ethylpentyl group, 2-ethylpentyl group, 3-ethylpentyl group, 1,1-dimethylpentyl group, 1,2-dimethylpentyl group, 1,3-dimethylpentyl group, 1,4-dimethylpentyl group, 2,2-dimethylpentyl group, 2,3-

- dimethylpentyl group, 2,4-dimethylpentyl group, 3,3-dimethylpentyl group, 3,4-dimethylpentyl group, 3,3-dimethylpentyl group, 3,4-dimethylpentyl group, 4,4-dimethylpentyl group, 1-propylbutyl group, 1-ethyl-1-methylbutyl group, 1-ethyl-2-methylbutyl group, 1-ethyl-3-methylbutyl group, 2-ethyl-1-methylbutyl group, 2-ethyl-2-methylbutyl group, 2-ethyl-3-methylbutyl group, 1,1,2-trimethylbutyl group, 1,1,3-trimethylbutyl group, 1,2,2-trimethylbutyl group, 1,2,3-trimethylbutyl group, 1,3,3-trimethylbutyl group, 2,2,3-trimethylbutyl group, 2,3,3-trimethylbutyl group, 1-methylheptyl group, 2-methylheptyl group, 3-methylheptyl group, 4-methylheptyl group, 5-methylheptyl group, 6-methylheptyl group, 1-ethylhexyl group, 2-ethylhexyl group, 3-ethylhexyl group, 4-ethylhexyl group, 1,1-dimethylhexyl group, 1,2-dimethylhexyl group, 1,3-dimethylhexyl group, 1,4-dimethylhexyl group, 1,5-dimethylhexyl group, 2,2-dimethylhexyl group, 2,3-dimethylhexyl group, 2,4-dimethylhexyl group, 2,5-dimethylhexyl group, 3,3-dimethylhexyl group, 3,4-dimethylhexyl group, 3,5-dimethylhexyl group, 4,4-dimethylhexyl group, 4,5-dimethylhexyl group, 5,5-dimethylhexyl group;
- 1-propylpentyl group, 2-propylpentyl group, 1-ethyl-1-methylpentyl group, 1-ethyl-2-methylpentyl group, 1-ethyl-3-methylpentyl group, 1-ethyl-4-methylpentyl group, 2-ethyl-1-methylpentyl group, 2-ethyl-2-methylpentyl group, 2-ethyl-3-methylpentyl group, 2-ethyl-4-methylpentyl group, 3-ethyl-1-methylpentyl group, 3-ethyl-2-methylpentyl group,

3-ethyl-3-methylpentyl group, 3-ethyl-4-methylpentyl group,
 1,1,2-trimethylpentyl group, 1,1,3-trimethylpentyl group,
 1,1,4-trimethylpentyl group, 1,2,2-trimethylpentyl group,
 1,2,3-trimethylpentyl group, 1,2,4-trimethylpentyl group,
 5 1,3,3-trimethylpentyl group, 1,3,4-trimethylpentyl group,
 1,4,4-trimethylpentyl group, 2,2,3-trimethylpentyl group,
 2,2,4-trimethylpentyl group, 2,3,3-trimethylpentyl group,
 2,3,4-trimethylpentyl group, 2,4,4-trimethylpentyl group,
 3,3,4-trimethylpentyl group, 3,4,4-trimethylpentyl group,
 10 1-methyl-1-propylbutyl group, 2-methyl-1-propylbutyl group,
 3-methyl-1-propylbutyl group, 1,1-diethylbutyl group, 1,2-
 diethylbutyl group, 2,2-diethylbutyl group, 1,2-dimethyl-1-
 ethylbutyl group, 1,3-dimethyl-1-ethylbutyl group, 2,2-
 dimethyl-1-ethylbutyl group, 2,3-dimethyl-1-ethylbutyl
 15 group, 3,3-dimethyl-1-ethylbutyl group, 1,1-dimethyl-2-
 ethylbutyl group, 1,2-dimethyl-2-ethylbutyl group, 1,3-
 dimethyl-2-ethylbutyl group, 2,3-dimethyl-2-ethylbutyl
 group, 3,3-dimethyl-2-ethylbutyl group, 1,1-diethyl-2-
 methylpropyl group, 1-methyloctyl group, 2-methyloctyl
 20 group, 3-methyloctyl group,
 4-methyloctyl group, 5-methyloctyl group, 6-methyloctyl
 group, 7-methyloctyl group, 1-ethylheptyl group, 2-
 ethylheptyl group,
 3-ethylheptyl group, 4-ethylheptyl group, 5-ethylheptyl
 25 group, 1,1-dimethylheptyl group, 1,2-dimethylheptyl group,
 1,3-dimethylheptyl group, 1,4-dimethylheptyl group, 1,5-
 dimethylheptyl group, 1,6-dimethylheptyl group, 2,2-
 dimethylheptyl group, 2,3-dimethylheptyl group, 2,4-

dimethylheptyl group, 2,5-dimethylheptyl group, 2,6-
 dimethylheptyl group, 3,3-dimethylheptyl group, 3,4-
 dimethylheptyl group, 3,5-dimethylheptyl group, 3,6-
 dimethylheptyl group, 4,4-dimethylheptyl group, 4,5-
 5 dimethylheptyl group, 4,6-dimethylheptyl group, 5,5-
 dimethylheptyl group, 5,6-dimethylheptyl group, 6,6-
 dimethylheptyl group;
 1-propylhexyl group, 2-propylhexyl group, 3-propylhexyl
 group, 1-ethyl-1-methylhexyl group, 1-ethyl-2-methylhexyl
 10 group, 1-ethyl-3-methylhexyl group, 1-ethyl-4-methylhexyl
 group, 1-ethyl-5-methylhexyl group, 2-ethyl-1-methylhexyl
 group, 2-ethyl-2-methylhexyl group, 2-ethyl-3-methylhexyl
 group, 2-ethyl-4-methylhexyl group, 2-ethyl-5-methylhexyl
 group, 3-ethyl-1-methylhexyl group, 3-ethyl-2-methylhexyl
 15 group, 3-ethyl-3-methylhexyl group, 3-ethyl-4-methylhexyl
 group, 3-ethyl-5-methylhexyl group, 4-ethyl-1-methylhexyl
 group, 4-ethyl-2-methylhexyl group, 4-ethyl-3-methylhexyl
 group, 4-ethyl-4-methylhexyl group, 4-ethyl-5-methylhexyl
 group, 1,1,2-trimethylhexyl group, 1,1,3-trimethylhexyl
 20 group, 1,1,4-trimethylhexyl group, 1,1,5-trimethylhexyl
 group, 1,2,2-trimethylhexyl group, 1,2,3-trimethylhexyl
 group, 1,2,4-trimethylhexyl group, 1,2,5-trimethylhexyl
 group, 1,3,3-trimethylhexyl group, 1,3,4-trimethylhexyl
 group, 1,3,5-trimethylhexyl group, 1,4,4-trimethylhexyl
 25 group, 1,4,5-trimethylhexyl group, 1,5,5-trimethylhexyl
 group, 2,2,3-trimethylhexyl group, 2,2,4-trimethylhexyl
 group, 2,2,5-trimethylhexyl group, 2,3,3-trimethylhexyl
 group, 2,3,4-trimethylhexyl group, 2,3,5-trimethylhexyl

group, 2,4,4-trimethylhexyl group, 2,4,5-trimethylhexyl group, 2,5,5-trimethylhexyl group, 3,3,4-trimethylhexyl group, 3,3,5-trimethylhexyl group, 3,4,4-trimethylhexyl group, 3,4,5-trimethylhexyl group, 3,5,5-trimethylhexyl group, 4,4,5-trimethylhexyl group, 4,5,5-trimethylhexyl group, 1-methyl-nonyl group, 2-methyl-nonyl group, 3-methyl-nonyl group, 4-methyl-nonyl group, 5-methyl-nonyl group, 6-methyl-nonyl group, 7-methyl-nonyl group, 8-methyl-nonyl group, and 9-methyl-nonyl group; straight-
 10 chained alkyl groups having 1 - 10 carbon atoms are preferred, among which methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group and n-pentyl group are particularly preferred.

If Z is a straight-chained or branched alkenyl group
 15 having 2 - 10 carbon atoms that may optionally be substituted by a halogen atom, exemplary straight-chained or branched alkenyl groups having 2 - 10 carbon atoms include straight-chained alkenyl groups such as vinyl group, 1-propenyl group, 2-propenyl group, 1-butenyl group, 2-
 20 butenyl group, 3-butenyl group, 1,3-butadienyl group, 2-pentenyl group, 3-pentenyl group, 2,4-pentadienyl group, 2-hexenyl group, 3-hexenyl group, 4-hexenyl group, 2,4-hexadienyl group, 2-heptenyl group, 3-heptenyl group, 4-heptenyl group, 5-heptenyl group, 2,4-heptadienyl group,
 25 2,5-heptadienyl group, 3,5-heptadienyl group, 2-octenyl group, 3-octenyl group, 4-octenyl group, 5-octenyl group, 6-octenyl group, 2,4-octadienyl group, 2,5-octadienyl group, 2,6-octadienyl group, 2,4,6-octatrienyl group, 2-nonenyl

- group, 3-nonenyl group, 4-nonenyl group, 5-nonenyl group, 6-nonenyl group, 7-nonenyl group, 2-decenyl group, 3-decenyl group, 4-decenyl group, 5-decenyl group, 6-decenyl group, 7-decenyl group, 8-decenyl group;
- 5 as well as branched alkenyl groups such as 1-methylethenyl group, 2-methyl-1-propenyl group, 2-methyl-2-propenyl group, 2-methyl-1-butenyl group, 3-methyl-2-butenyl group, 2-methyl-3-butenyl group, 2,3-dimethyl-1,3-butadienyl group, 3-ethyl-2-propenyl group, 4-methyl-3-propenyl group, 3-
- 10 methyl-2,4-propadienyl group, 3,4-diethyl-2-hexenyl group, 4-methyl-3-hexenyl group, 2-methyl-4-hexenyl group, 3,5-dimethyl-2,4-hexadienyl group, 5-ethyl-3-methyl-2-heptenyl group, 5-methyl-3-heptenyl group, 4-n-propyl-4-heptenyl group, 3,6-dimethyl-5-heptenyl group, 5-ethyl-2,4-
- 15 heptadienyl group, 2,6-dimethyl-2,5-heptadienyl group, 4-ethyl-3,5-heptadienyl group, 4,6-dimethyl-2-octenyl group, 5-ethyl-3-octenyl group, 3-ethyl-4-octenyl group, 3-ethyl-5-octenyl group, 3,4-dimethyl-6-octenyl group, 5-ethyl-2,4-octadienyl group, 3-methyl-2,5-octadienyl group, 5-ethyl-
- 20 2,6-octadienyl group, 4-methyl-2,4,6-octatrienyl group, 5-methyl-2-nonenyl group, 6-methyl-3-nonenyl group, 7-methyl-4-nonenyl group, 3-methyl-5-nonenyl group, 4-methyl-6-nonenyl group, 3-methyl-7-nonenyl group, etc.

If Z is a straight-chained or branched alkynyl group

25 having 2 - 10 carbon atoms that may optionally be substituted by a halogen atom, exemplary straight-chained or branched alkynyl groups having 2 - 10 carbon atoms include straight-chained alkynyl groups such as ethynyl

- group, 1-propynyl group, 2-propynyl group, 1-butyryl group, 2-butyryl group, 3-butyryl group, 1,3-butadiynyl group, 2-pentyryl group, 3-pentyryl group, 2,4-pentadiynyl group, 2-hexynyl group, 3-hexynyl group, 4-hexynyl group, 2,4-hexadiynyl group, 2-heptyryl group, 3-heptyryl group, 4-heptyryl group, 5-heptyryl group, 2,4-heptadiynyl group, 2,5-heptadiynyl group, 3,5-heptadiynyl group, 2-octynyl group, 3-octynyl group, 4-octynyl group, 5-octynyl group, 6-octynyl group, 2,4-octadiynyl group, 2,5-octadiynyl group, 2,6-octadiynyl group, 2,4,6-octatriynyl group, 2-nonyryl group, 3-nonyryl group, 4-nonyryl group, 5-nonyryl group, 6-nonyryl group, 7-nonyryl group, 2-decyryl group, 3-decyryl group, 4-decyryl group, 5-decyryl group, 6-decyryl group, 7-decyryl group, 8-decyryl group;
- as well as branched alkynyl groups such as 1-methyl-2-propynyl group, 3-methyl-1-butyryl group, 2-methyl-3-butyryl group, 4-methyl-2-pentyryl group, 2-methyl-3-pentyryl group, 4-ethyl-2-hexynyl group, 5-methyl-3-hexynyl group, 2-methyl-4-hexynyl group, 5-ethyl-6-methyl-2-heptyryl group, 5-methyl-3-heptyryl group, 3-n-propyl-4-heptyryl group, 4,4-dimethyl-5-heptyryl group, 6-methyl-2,4-heptadiynyl group, 4-methyl-2,5-heptadiynyl group, 2-methyl-3,5-heptadiynyl group, 6,6-dimethyl-2-octynyl group, 6-methyl-3-octynyl group, 3-ethyl-4-octynyl group, 4-methyl-5-octynyl group, 4,8-dimethyl-6-octynyl group, 7-methyl-2,4-octadiynyl group, 4-methyl-2,5-octadiynyl group, 5-ethyl-2,6-octadiynyl group, 5-methyl-2-nonyryl group, 6-methyl-3-nonyryl group, 7-methyl-4-nonyryl group, 8-methyl-

5-nonynyl group, 4-methyl-6-nonynyl group, 3-methyl-7-nonynyl group, etc.

- If Z is $-O-R^d$, R^d is a hydrogen atom or a protective group for a hydroxyl group, and a hydrogen atom is preferred. The protective group for a hydroxyl group may be exemplified by the same protective groups for a hydroxyl group that were already mentioned for R^a , and preferred examples as well as particularly preferred examples are also the same as those mentioned for R^a .
- Typically, preferred examples of Z are a straight-chained or branched alkyl group having 1 - 10 carbon atoms that may optionally be substituted by a halogen atom, a hydroxyl group, and a straight-chained or branched alkyl group having 1 - 10 carbon atoms that is substituted by any one group selected from the group consisting of a cycloalkyl group having 3 - 6 carbon atoms, a hydroxyl group, a heterocyclic group and an optionally substituted phenyl group; straight-chained or branched alkyl groups having 3 - 10 carbon atoms that are substituted by a halogen atom are preferred and among these, straight-chained or branched alkyl groups having 3 - 8 carbon atoms that are substituted by a fluorine atom are particularly preferred, with 4,4,5,5,5-pentafluoropentyl group being most preferred. Considering peroral absorption, Z is preferably $-COOH$. Further considering peroral absorption, Z may preferably be a hydrogen atom if Q is Q^{17} where R^7 is a hydrogen atom.

If Q is Q^{65} , Q^{66} , Q^{67} , Q^{68} , Q^{69} or Q^{70} , Z is preferably a

hydrogen atom or an unsubstituted straight-chained or branched alkyl group having 1 - 3 carbon atoms and, among these, a hydrogen atom is particularly preferred. If Q is Q^{71} , Q^{72} , Q^{73} , Q^{74} , Q^{75} or Q^{76} , Z is preferably a hydrogen atom.

5 Compounds represented by the general formula (I) in which Z is $-O-R^d$ (where R^d has the same meaning as defined above) or $-COOH$ are also useful as intermediates for compounds represented by the general formula (I) in which Z is neither $-O-R^d$ (where R^d has the same meaning as defined
10 above) nor $-COOH$.

If Q is Q^3 , the nitrogen atom and R^8 in Q^3 may combine with Z to form a heterocyclic group, as exemplified by morpholino group, pyrrolidinyl group, piperidino group, etc.

Referring further to the general formula (I), it is
15 preferred that the dashed line in 4(5)-position signifies a single bond in combination with the solid line and X^2 signifies any one group selected from the group consisting of $-(CH_2)_pCO-NR^8Z^1$ (p represents an integer of at least 1, R^8 represents a straight-chained or branched lower alkyl group
20 having 1 - 6 carbon atoms, and Z^1 represents a hydrogen atom or a straight-chained or branched alkyl group having 1 - 10 carbon atoms that may optionally be substituted by a halogen atom), $-(CH_2)_p-SO_2-Z^1$ (p and Z^1 have the same meanings as defined above), $-(CH_2)_p-SO-Z^1$ (p and Z^1 have the
25 same meanings as defined above), $-Ph-O-(CH_2)_p-CO-NR^8Z^1$ (Ph represents a phenylene group and p, R^8 and Z^1 have the same meanings as defined above), and $-Ph-O-(CH_2)_p-H$ (p has the same meaning as defined above), with p being more referably

an integer of 1 - 13;

speaking of the group represented by $-NR^8Z^1$ in $-(CH_2)_p-CO-NR^8Z^1$,

amino group and n-pentylamino group are preferred if p is

5 5;

if p is 6, dimethylamino group and diethylamino group are preferred;

if p is 7, dimethylamino group, ethylamino group, n-butylamino group, i-propylamino group, cyclohexylamino group, N-n-butyl-N-methylamino group, diethylamino group, methylamino group, N-ethyl-N-methylamino group, N-methyl-N-n-propylamino group, N-methyl-N-i-propylamino group, N-t-butyl-N-methylamino group, n-propylamino group, n-hexylamino group, i-pentylamino group, i-butylamino group, 2,2-dimethylpropylamino group, 1-ethylpropylamino group, di-n-hexylamino group, amino group and n-pentylamino group are preferred, with dimethylamino group, ethylamino group, i-propylamino group, N-n-butyl-N-methylamino group, diethylamino group, methylamino group, N-ethyl-N-methylamino group, N-methyl-N-n-propylamino group, N-methyl-N-i-propylamino group and n-propylamino group being more preferred, and with dimethylamino group, ethylamino group, N-n-butyl-N-methylamino group, diethylamino group, methylamino group, N-ethyl-N-methylamino group and N-methyl-N-i-propylamino group being particularly preferred;

if p is 8, dimethylamino group, diethylamino group and N-n-butyl-N-methylamino group are preferred;

if p is 9, amino group, n-pentylamino group, dimethylamino

- group, diethylamino group, N-ethyl-N-methylamino group, N-n-butyl-N-methylamino group and N-methyl-N-n-propylamino group are preferred, with dimethylamino group, diethylamino group, N-n-butyl-N-methylamino group and N-methyl-N-n-propylamino group being more preferred, and dimethylamino group and N-methyl-N-n-propylamino being particularly preferred;
- if p is 10, dimethylamino group, diethylamino group, N-ethyl-N-methylamino group, N-n-butyl-N-methylamino group and N-methyl-N-n-propylamino group are preferred, with dimethylamino group being more preferred;
- if p is 11, dimethylamino group, diethylamino group, N-n-butyl-N-methylamino group, amino group and n-pentylamino group are preferred;
- if p is 13, amino group and n-pentylamino group are preferred;
- p in $-(CH_2)_p-SO_2-Z^1$ is preferably an integer of 5 - 13; the group represented by $-Z^1$ in $-(CH_2)_p-SO_2-Z^1$ is preferably 4,4,5,5,5-pentafluoropentyl group;
- p in $-(CH_2)_p-SO-Z^1$ is preferably an integer of 7 - 13; the group represented by $-Z^1$ in $-(CH_2)_p-SO-Z^1$ is preferably 4,4,5,5,5-pentafluoropentyl group;
- p in $-Ph-O-(CH_2)_p-CO-NR^8Z^1$ is preferably an integer of 1 - 7; the group represented by $-NR^8Z^1$ in $-Ph-O-(CH_2)_p-CO-NR^8Z^1$ is preferably amino group or n-pentylamino group;
- speaking further of that group, it is preferably amino group if p is 1; if p is 3, amino group and n-pentylamino group are

preferred;

if p is 7, amino group and n-pentylamino group are preferred;

p in $-\text{Ph}-\text{O}-(\text{CH}_2)_p-\text{H}$ is preferably 1.

- 5 Referring further to the general formula (I), it is preferred that the dashed line in 4(5) position signifies a single bond or a double bond in combination with the solid line and X^2 signifies any one group selected from the group consisting of $-(\text{CH}_2)_p-\text{COOH}$ (p is an integer of at least 1),
- 10 $-(\text{CH}_2)_p-\text{OH}$ (p has the same meaning as defined above), $-\text{Ph}-\text{O}-(\text{CH}_2)_p-\text{COOH}$ (Ph represents a phenylene group and p has the same meaning as defined above), $-(\text{CH}_2)_p-\text{CO}-\text{NR}^8\text{Z}^2$ (p has the same meaning as defined above, R^8 represents a hydrogen atom or a straight-chained or branched lower alkyl group
- 15 having 1 - 6 carbon atoms, Z^2 represents a straight-chained or branched alkyl group having 1 - 10 carbon atoms that is substituted by any one group selected from the group consisting of a cycloalkyl group, a hydroxyl group, a carboxyl group, a heterocyclic group and a phenyl group, or
- 20 $-\text{NR}^8\text{Z}^2$ may be such that N, R^8 and Z^2 combine together to form a hetero ring), $-(\text{CH}_2)_p-\text{Ph}-\text{O}-(\text{CH}_2)_q-\text{CO}-\text{NR}^8\text{Z}^3$ (Ph, p and R^8 have the same meanings as defined above, q represents an integer of at least 1, and Z^3 represents a hydrogen atom or a straight-chained or branched alkyl group having 1 - 10
- 25 carbon atoms that may optionally be substituted by any one group selected from the group consisting of a cycloalkyl group, a hydroxyl group, a carboxyl group, a heterocyclic group and a phenyl group, or $-\text{NR}^8\text{Z}^3$ may be such that N, R^8

- and Z^3 combine together to form a hetero ring) and $-(CH_2)_p-CH(COOH)-(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), with p being more preferably an integer of 1 - 13;
 p in $-(CH_2)_p-COOH$ is preferably an integer of 5 - 13;
 5 p in $-(CH_2)_p-OH$ is preferably an integer of 7 - 9;
 p in $-Ph-O-(CH_2)_p-COOH$ is preferably an integer of 1 - 7;
 p in $-(CH_2)_p-CO-NR^8Z^3$ is preferably an integer of 6 - 11;
 the group represented by $-NR^8Z^3$ in $-(CH_2)_p-CO-NR^8Z^3$ is preferably exemplified by, cyclohexylmethylamino group,
 10 cyclopropylmethylamino group, 3-hydroxypropylamino group, t-butylbenzylamino group, 2,2-diphenylethylamino group, N-methyl-N-benzylamino group, phenylamino group, benzylamino group, 2-phenylethylamino group, piperidino group, pyrrolidinyl group and morpholino group, with N-methyl-N-
 15 benzylamino group, benzylamino group, 2-phenylethylamino group, piperidino group, pyrrolidinyl group and morpholino group being more preferred, and piperidino group, pyrrolidinyl group and morpholino group being particularly preferred;
 20 p in $-(CH_2)_p-Ph-O-(CH_2)_q-CO-NR^8Z^3$ is preferably 3;
 q in $-(CH_2)_p-Ph-O-(CH_2)_q-CO-NR^8Z^3$ is preferably 3 or 4;
 the group represented by $-NR^8Z^3$ in $-(CH_2)_p-Ph-O-(CH_2)_q-CO-NR^8Z^3$ is preferably exemplified by methylamino group, dimethylamino group and pyrrolidinyl group;
 25 p in $-(CH_2)_p-CH(COOH)-(CH_2)_3-CF_2-CF_3$ is preferably 8;
 p in $-(CH_2)_p-Ph-O-(CH_2)_q-COOH$ is preferably 3;
 q in $-(CH_2)_p-Ph-O-(CH_2)_q-COOH$ is preferably 3 or 4.

Referring further to the general formula (I), it is

preferred that the dashed line in 4(5) position signifies a single bond or a double bond in combination with the solid line and X^1 signifies any one group selected from the group consisting of $-(CH_2)_p-COOH$ (p is an integer of at least 1),

5 $-(CH_2)_p-CH(COOH)-(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), $-(CH_2)_p-CH(COOMe)-(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), $-O-(CH_2)_p-COOH$ (p has the same meaning as defined above), $-O-(CH_2)_p-CH(COOH)-(CH_2)_3-$

10 CF_2-CF_3 (p has the same meaning as defined above), $-(CH_2)_p-S-$
 $(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), $-(CH_2)_p-SO-(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), $-O-(CH_2)_p-SO-(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), $-O-(CH_2)_p-SO_2-(CH_2)_3-CF_2-CF_3$ (p has the same meaning as defined above), $-Ph-O-CH_3$ (Ph represents a

15 phenylene group), $-Ph-O-(CH_2)_p-COOH$ (Ph and p have the same meanings as defined above), $-(CH_2)_p-CO-NR^8Z^3$ (p has the same meaning as defined above, R^8 represents a hydrogen atom or a straight-chained or branched lower alkyl group having 1 -

20 6 carbon atoms, Z^3 represents a hydrogen atom or a
 straight-chained or branched alkyl group having 1 - 10 carbon atoms that may optionally be substituted by any one group selected from the group consisting of a cycloalkyl group, a hydroxyl group, a carboxyl group, a heterocyclic group and a phenyl group, or $-NR^8Z^3$ may be such that N , R^8

25 and Z^3 combine together to form a hetero ring), $-Ph-O-(CH_2)_p-CO-NR^8Z^3$ (Ph , p , R^8 , Z^3 and $-NR^8Z^3$ have the same meanings as defined above) and $-O-(CH_2)_p-CO-NR^8Z^3$ (p , R^8 , Z^3 and $-NR^8Z^3$ have the same meanings as defined above), with p

- being more preferably an integer of 3 - 13;
- p in $-(\text{CH}_2)_p\text{-COOH}$ is preferably an integer of 7 - 11;
- p in $-(\text{CH}_2)_p\text{-CH(COOH)-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably 8;
- p in $-(\text{CH}_2)_p\text{-CH(COOMe)-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably 8;
- 5 p in $-\text{O-}(\text{CH}_2)_p\text{-COOH}$ is preferably an integer of 5 - 13;
- p in $-\text{O-}(\text{CH}_2)_p\text{-CH(COOH)-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably 8;
- p in $-(\text{CH}_2)_p\text{-S-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably 10;
- p in $-(\text{CH}_2)_p\text{-SO-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably 10;
- p in $-\text{O-}(\text{CH}_2)_p\text{-SO-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably an integer of
- 10 5 - 13;
- p in $-\text{O-}(\text{CH}_2)_p\text{-SO}_2\text{-}(\text{CH}_2)_3\text{-CF}_2\text{-CF}_3$ is preferably an integer of
- 7 - 13;
- p in $-\text{Ph-O-}(\text{CH}_2)_p\text{-COOH}$ is preferably an integer of 3 - 7;
- p in $-(\text{CH}_2)_p\text{-CO-NR}^8\text{Z}^3$ is preferably an integer of 7 - 11;
- 15 the group represented by $-\text{NR}^8\text{Z}^3$ in $-(\text{CH}_2)_p\text{-CO-NR}^8\text{Z}^3$ is preferably exemplified by, amino group, n-pentylamino group, dimethylamino group, methylamino group, N-ethyl-N-methylamino group, N-methyl-N-n-propylamino group, diethylamino group, benzylamino group, N-n-butyl-N-
- 20 methylamino group, 2-hydroxyethylamino group, morpholino group and piperidino group;
- p in $-\text{Ph-O-}(\text{CH}_2)_p\text{-CO-NR}^8\text{Z}^3$ is preferably 7;
- the group represented by $-\text{NR}^8\text{Z}^3$ in $-\text{Ph-O-}(\text{CH}_2)_p\text{-CO-NR}^8\text{Z}^3$ is preferably exemplified by amino group and n-pentylamino
- 25 group;
- p in $-\text{O-}(\text{CH}_2)_p\text{-CO-NR}^8\text{Z}^3$ is preferably an integer of 5 - 13;
- the group represented by $-\text{NR}^8\text{Z}^3$ in $-\text{O-}(\text{CH}_2)_p\text{-CO-NR}^8\text{Z}^3$ is preferably exemplified by amino group and n-pentylamino

group.

Specifically, preferred examples of X^1 and X^2 are a hydrogen atom, 10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl group, 11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyl group, 12-(4,4,5,5,5-pentafluoropentylsulfinyl)dodecyl group, 10-(4,4,5,5,5-pentafluoropentylsulfonyl)decyl group, 11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyl group, 12-(4,4,5,5,5-pentafluoropentylsulfonyl)dodecyl group, 10-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}decyl group, 11-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}undecyl group, 9-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}nonyl group, 10-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}decyl group, 9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyloxy group, 10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyloxy group, 11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyloxy group, 9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyloxy group, 10-(4,4,5,5,5-pentafluoropentylsulfonyl)decyloxy group, 11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyloxy group; 9-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}nonyloxy group, 10-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}decyloxy group, 8-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}octyloxy group, 9-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}nonyloxy group, 4-{8-(4,4,5,5,5-pentafluoropentylsulfinyl)octyloxy}phenyl group, 4-{9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyloxy}phenyl group, 4-{8-(4,4,5,5,5-pentafluoropentylsulfonyl)octyloxy}phenyl group, 4-{9-

- (4,4,5,5,5-pentafluoropentylsulfonyl)nonyloxy}phenyl group,
 4-[8-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}octyloxy]phenyl group, 4-[9-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}nonyloxy]phenyl group, 4-[7-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}heptyloxy]phenyl group, 4-[8-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}octyloxy]phenyl group, 6-[4-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}phenyl]hexyl group, 5-[4-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}phenyl]pentyloxy group, tridecyloxy group, 11-carboxy-15,15,16,16,16-pentafluorohexadecyl) group, 4-{{2-hydroxy-3-(4,4,5,5,5-pentafluoropentylsulfinylethyloxy)propyl}oxy}phenyl group, 4-hydroxy-9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl group, 10-carboxy-14,14,15,15,15-pentafluoropentadecyloxy group, 9-carboxy-13,13,14,14,14-pentafluorotetradecyloxy group, 6-carboxy-10,10,11,11,11-pentafluoroundecyl group, 10-carboxy-14,14,15,15,15-pentafluoropentadecyl group, 14-carboxy-18,18,19,19,19-pentafluorononadecyl group, 9-carboxynonyloxy group, 6-carboxyhexyl group, 10-carboxydecyl group, 14-carboxytetradecyl group, 3-{4-(4-carboxybutyl)phenyl}propyl group, 3-{4-(4-carboxy-8,8,9,9,9-pentafluorononyl)phenyl}propyl group, 5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl group, 9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl group, 13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyl group, 4-hydroxy-10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl group,

4-hydroxy-15,15,16,16,16-pentafluorohexadecyl group, 9-
 {N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}nonyl group,
 and 8-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}octyl group;
 5-carboxypentyl group, 7-carboxyheptyl group, 9-
 5 carboxynonyl group, 11-carboxyundecyl group, 13-
 carboxytridecyl group, 9-carboxy-13,13,14,14,14-
 pentafluorotetradecyl group, 9-methoxycarbonyl-
 13,13,14,14,14-pentafluorotetradecyl group, 5-
 carboxypentyloxy group, 7-carboxyheptyloxy group, 10-
 10 carboxydecyloxy group, 11-carboxyundecyloxy group, 13-
 carboxytridecyloxy group, 23-carboxytricosanyloxy group, 7-
 (N,N-dimethylaminocarbonyl)heptyl group, 7-(N-
 ethylaminocarbonyl)heptyl group, 7-{N-
 (cyclopropylmethyl)aminocarbonyl}heptyl group, 7-{N-
 15 (cyclohexylmethyl)aminocarbonyl}heptyl group, 7-(N-
 butylaminocarbonyl)heptyl group, 7-(N-
 isopropylaminocarbonyl)heptyl group, 7-(N-t-
 butylaminocarbonyl)heptyl group, 7-(N-
 cyclohexylaminocarbonyl)heptyl group, 7-{N-(3-
 20 hydroxypropyl)aminocarbonyl}heptyl group, 7-(N-methyl-N-
 butylaminocarbonyl)heptyl group, 7-(N,N-
 diethylaminocarbonyl)heptyl group, 7-
 (piperidinocarbonyl)heptyl group, 7-{N-(4-t-
 butylbenzyl)aminocarbonyl}heptyl group, 7-{N-(2,2-
 25 diphenylethyl)aminocarbonyl}heptyl group, 7-{N-(2-
 furylmethyl)aminocarbonyl}heptyl group, 7-(N-
 methylaminocarbonyl)heptyl group, 7-(N-methyl-N-
 ethylaminocarbonyl)heptyl group, 7-(N-methyl-N-

- propylaminocarbonyl)heptyl group, 7-(N-methyl-N-isopropylaminocarbonyl)heptyl group, 7-(N-methyl-N-benzylaminocarbonyl)heptyl group, 7-(1-pyrrolidinylcarbonyl)heptyl group, 7-
- 5 (morpholinocarbonyl)heptyl group, 7-(N-methyl-N-t-butylaminocarbonyl)heptyl group, 7-(N-cyclopropylaminocarbonyl)heptyl group, 6-(N,N-dimethylaminocarbonyl)hexyl group, 6-(N,N-diethylaminocarbonyl)hexyl group, 6-
- 10 (piperidinocarbonyl)hexyl group, 8-(N,N-dimethylaminocarbonyl)octyl group, 8-(N,N-diethylaminocarbonyl)octyl group, 8-(N-methyl-N-butylaminocarbonyl)octyl group, 8-(N-benzylaminocarbonyl)octyl group, 8-{N-(2-
- 15 hydroxyethyl)aminocarbonyl}octyl group, 8-(piperidinocarbonyl)octyl group, 9-(N,N-dimethylaminocarbonyl)nonyl group, 9-(N,N-diethylaminocarbonyl)nonyl group, 9-(1-pyrrolidinylcarbonyl)nonyl group, 9-(N-methyl-N-
- 20 ethylaminocarbonyl)nonyl group, 9-(N-methyl-N-butylaminocarbonyl)nonyl group, 9-(N-benzylaminocarbonyl)nonyl group, 9-(piperidinocarbonyl)nonyl group, 9-{N-(2-
- hydroxyethyl)aminocarbonyl}nonyl group, 9-(N-methyl-N-
- 25 propylaminocarbonyl)nonyl group, 9-(morpholinocarbonyl)nonyl group, 10-(N,N-dimethylaminocarbonyl)decyl group, 10-(N,N-diethylaminocarbonyl)decyl group, 10-(N-methyl-N-

- ethylaminocarbonyl)decyl group, 10-(N-methyl-N-propylaminocarbonyl)decyl group, 10-(N-methyl-N-butylaminocarbonyl)decyl group, 10-(morpholinocarbonyl)decyl group, 11-(N,N-
- 5 dimethylaminocarbonyl)undecyl group, 11-(N,N-diethylaminocarbonyl)undecyl group, 11-(piperidinocarbonyl)undecyl group, 11-(N-benzylaminocarbonyl)undecyl group, 11-(N-methyl-N-butylaminocarbonyl)undecyl group, 11-{N-(2-
- 10 hydroxyethyl)aminocarbonyl}undecyl group, 7-{N-(2-hydroxyethyl)aminocarbonyl}heptyl group, 7-(N-propylaminocarbonyl)heptyl group, 7-(N-hexylaminocarbonyl)heptyl group, 7-(N-isopentylaminocarbonyl)heptyl group, 7-(N-
- 15 isobutylaminocarbonyl)heptyl group, 7-(N-neopentylaminocarbonyl)heptyl group, 7-{N-(3-pentyl)aminocarbonyl}heptyl group, 7-(N,N-dihexylaminocarbonyl)heptyl group, 7-(N-phenylaminocarbonyl)heptyl group, 7-(N-
- 20 benzylaminocarbonyl)heptyl group, 7-{N-(2-phenylethyl)aminocarbonyl}heptyl group, 5-(aminocarbonyl)pentyl group, 5-(N-pentylaminocarbonyl)pentyl group, 7-(aminocarbonyl)heptyl group, 7-(N-pentylaminocarbonyl)heptyl group, 9-
- 25 (aminocarbonyl)nonyl group, 9-(N-pentylaminocarbonyl)nonyl group, 11-(aminocarbonyl)undecyl group, 11-(N-pentylaminocarbonyl)undecyl group, 13-(aminocarbonyl)tridecyl group, 13-(N-

- pentylaminocarbonyl)tridecyl group, 8-(N-methyl-N-ethylaminocarbonyl)octyl group, 8-(N-methyl-N-propylaminocarbonyl)octyl group, 8-(morpholinocarbonyl)octyl group, 8-(N-methylaminocarbonyl)octyl group, 10-(4,4,5,5,5-pentafluoropentylsulfanyl)decyl group, 7-hydroxyheptyl group, 8-hydroxyoctyl group, 9-hydroxynonyl group, 7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyl group, 7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyl group, 9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyl group, 13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyl group; 4-(carboxymethoxy)phenyl group, 4-(3-carboxypropoxy)phenyl group, 4-(7-carboxyheptyloxy)phenyl group, 4-(carbamoylmethoxy)phenyl group, 4-(3-carbamoylpropoxy)phenyl group, 4-(7-carbamoylheptyloxy)phenyl group, 4-(3-N-pentylcarbamoylpropoxy)phenyl group, 4-(7-N-pentylcarbamoylheptyloxy)phenyl group, 4-methoxyphenyl group;
- 20 5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl group, 7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy group, 13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyloxy group; 7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy group, 13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyloxy group;
- 25 4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl group, 4-{7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy}phenyl group, 4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)pentyl group, 4-{7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy}phenyl group,

- 4-{7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy}phenyl group;
- 3-{3-(3-carboxypropoxy)phenyl}propyl group, 3-{3-(4-carboxybutoxy)phenyl}propyl group, 3-[3-{3-N-
- 5 methylaminocarbonyl}propoxy}phenyl]propyl group, 3-[3-{3-N,N-dimethylaminocarbonyl}propoxy}phenyl]propyl group, 3-[3-{3-(1-pyrrolidinylcarbonyl)propoxy}phenyl]propyl group, 3-[3-{4-N-methylaminocarbonyl}butoxy}phenyl]propyl group, 3-[3-{4-(N,N-dimethylaminocarbonyl)butoxy}phenyl]propyl
- 10 group, 3-[3-{4-(1-pyrrolidinylcarbonyl)butoxy}phenyl]propyl group;
- 5-(aminocarbonyl)pentylloxy group, 5-(N-pentylaminocarbonyl)pentylloxy group, 7-(aminocarbonyl)heptyloxy group, 7-(N-
- 15 pentylaminocarbonyl)heptyloxy group, 9-(aminocarbonyl)nonyloxy group, 9-(N-pentylaminocarbonyl)nonyloxy group, 11-(aminocarbonyl)undecyloxy group, 11-(N-pentylaminocarbonyl)undecyloxy group, 13-
- 20 (aminocarbonyl)tridecyloxy group, and 13-(N-pentylaminocarbonyl)tridecyloxy group. More preferred are 10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl group, 11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyl group, 11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyl group, 9-
- 25 (4,4,5,5,5-pentafluoropentylsulfinyl)nonyloxy group, 11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyloxy group, 9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyloxy group, 11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyloxy group, 9-

- carboxy-13,13,14,14,14-pentafluorotetradecyloxy group, 9-carboxynonyloxy group, 5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl group, 9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl group;
- 5 5-carboxypentyl group, 7-carboxyheptyl group, 9-carboxynonyl group, 11-carboxyundecyl group, 13-carboxytridecyl group, 9-carboxy-13,13,14,14,14-pentafluorotetradecyl group, 9-methoxycarbonyl-13,13,14,14,14-pentafluorotetradecyl group, 5-
- 10 carboxypentyloxy group, 7-carboxyheptyloxy group, 10-carboxydecyloxy group, 11-carboxyundecyloxy group, 13-carboxytridecyloxy group, 23-carboxytricosanyloxy group, 7-(N,N-dimethylaminocarbonyl)heptyl group, 7-(N-ethylaminocarbonyl)heptyl group, 7-(N-
- 15 (cyclopropylmethyl)aminocarbonyl}heptyl group, 7-(N-(cyclohexylmethyl)aminocarbonyl}heptyl group, 7-(N-butylaminocarbonyl)heptyl group, 7-(N-isopropylaminocarbonyl)heptyl group, 7-(N-t-butylaminocarbonyl)heptyl group, 7-(N-
- 20 cyclohexylaminocarbonyl)heptyl group, 7-(N-(3-hydroxypropyl)aminocarbonyl}heptyl group, 7-(N-methyl-N-butylaminocarbonyl)heptyl group, 7-(N,N-diethylaminocarbonyl)heptyl group, 7-(piperidinocarbonyl)heptyl group, 7-(N-(4-t-
- 25 butylbenzyl)aminocarbonyl}heptyl group, 7-(N-(2,2-diphenylethyl)aminocarbonyl}heptyl group, 7-(N-(2-furylmethyl)aminocarbonyl}heptyl group, 7-(N-methylaminocarbonyl)heptyl group, 7-(N-methyl-N-

- ethylaminocarbonyl)heptyl group, 7-(N-methyl-N-propylaminocarbonyl)heptyl group, 7-(N-methyl-N-isopropylaminocarbonyl)heptyl group, 7-(N-methyl-N-benzylaminocarbonyl)heptyl group, 7-(1-pyrrolidinylcarbonyl)heptyl group, 7-(morpholinocarbonyl)heptyl group, 7-(N-methyl-N-t-butylaminocarbonyl)heptyl group, 7-(N-cyclopropylaminocarbonyl)heptyl group, 6-(N,N-dimethylaminocarbonyl)hexyl group, 6-(N,N-diethylaminocarbonyl)hexyl group, 6-(piperidinocarbonyl)hexyl group, 8-(N,N-dimethylaminocarbonyl)octyl group, 8-(N,N-diethylaminocarbonyl)octyl group, 8-(N-methyl-N-butylaminocarbonyl)octyl group, 8-(N-benzylaminocarbonyl)octyl group, 8-(N-(2-hydroxyethyl)aminocarbonyl)octyl group, 8-(piperidinocarbonyl)octyl group, 9-(N,N-dimethylaminocarbonyl)nonyl group, 9-(N,N-diethylaminocarbonyl)nonyl group, 9-(1-pyrrolidinylcarbonyl)nonyl group, 9-(N-methyl-N-ethylaminocarbonyl)nonyl group, 9-(N-methyl-N-butylaminocarbonyl)nonyl group, 9-(N-benzylaminocarbonyl)nonyl group, 9-(piperidinocarbonyl)nonyl group, 9-(N-(2-hydroxyethyl)aminocarbonyl)nonyl group, 9-(N-methyl-N-propylaminocarbonyl)nonyl group, 9-(morpholinocarbonyl)nonyl group, 10-(N,N-dimethylaminocarbonyl)decyl group, 10-(N,N-

- diethylaminocarbonyl)decyl group, 10-(N-methyl-N-ethylaminocarbonyl)decyl group, 10-(N-methyl-N-propylaminocarbonyl)decyl group, 10-(N-methyl-N-butylaminocarbonyl)decyl group, 10-
- 5 (morpholinocarbonyl)decyl group, 11-(N,N-dimethylaminocarbonyl)undecyl group, 11-(N,N-diethylaminocarbonyl)undecyl group, 11-(piperidinocarbonyl)undecyl group, 11-(N-benzylaminocarbonyl)undecyl group, 11-(N-methyl-N-
- 10 butylaminocarbonyl)undecyl group, 11-(N-(2-hydroxyethyl)aminocarbonyl)undecyl group, 7-(N-(2-hydroxyethyl)aminocarbonyl)heptyl group, 7-(N-propylaminocarbonyl)heptyl group, 7-(N-hexylaminocarbonyl)heptyl group, 7-(N-
- 15 isopentylaminocarbonyl)heptyl group, 7-(N-isobutylaminocarbonyl)heptyl group, 7-(N-neopentylaminocarbonyl)heptyl group, 7-(N-(3-pentyl)aminocarbonyl)heptyl group, 7-(N,N-dihexylaminocarbonyl)heptyl group, 7-(N-
- 20 phenylaminocarbonyl)heptyl group, 7-(N-benzylaminocarbonyl)heptyl group, 7-(N-(2-phenylethyl)aminocarbonyl)heptyl group, 5-(aminocarbonyl)pentyl group, 5-(N-pentylaminocarbonyl)pentyl group, 7-(aminocarbonyl)heptyl
- 25 group, 7-(N-pentylaminocarbonyl)heptyl group, 9-(aminocarbonyl)nonyl group, 9-(N-pentylaminocarbonyl)nonyl group, 11-(aminocarbonyl)undecyl group, 11-(N-pentylaminocarbonyl)undecyl group, 13-

- (aminocarbonyl)tridecyl group, 13-(N-pentylaminocarbonyl)tridecyl group, 8-(N-methyl-N-ethylaminocarbonyl)octyl group, 8-(N-methyl-N-propylaminocarbonyl)octyl group, 8-
- 5 (morpholinocarbonyl)octyl group, 8-(N-methylaminocarbonyl)octyl group, 10-(4,4,5,5,5-pentafluoropentylsulfanyl)decyl group, 7-hydroxyheptyl group, 8-hydroxyoctyl group, 9-hydroxynonyl group, 7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyl group, 7-
- 10 (4,4,5,5,5-pentafluoropentylsulfonyl)heptyl group, 9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyl group, 13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyl group; 4-(carboxymethoxy)phenyl group, 4-(3-carboxypropoxy)phenyl group, 4-(7-carboxyheptyloxy)phenyl group, 4-
- 15 (carbamoylmethoxy)phenyl group, 4-(3-carbamoylpropoxy)phenyl group, 4-(7-carbamoylheptyloxy)phenyl group, 4-(3-N-pentylcarbamoylpropoxy)phenyl group, 4-(7-N-pentylcarbamoylheptyloxy)phenyl group, 4-methoxyphenyl
- 20 group;
- 5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl group, 7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy group, 13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyloxy group; 7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy group, 13-
- 25 (4,4,5,5,5-pentafluoropentylsulfonyl)tridecyloxy group; 4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl}phenyl group, 4-{7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy}phenyl group, 4-{5-

(4,4,5,5,5-pentafluoropentylsulfonyl)pentylloxy}phenyl group,
 4-{7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy group;
 3-{3-(3-carboxypropoxy)phenyl}propyl group, 3-{3-(4-
 carboxybutoxy)phenyl}propyl group, 3-[3-{3-N-
 5 methylaminocarbonyl}propoxy}phenyl]propyl group, 3-[3-{3-
 N,N-dimethylaminocarbonyl}propoxy}phenyl]propyl group, 3-
 [3-{3-(1-pyrrolidinylcarbonyl)propoxy}phenyl]propyl group,
 3-[3-{4-(N-methylaminocarbonyl)butoxy}phenyl]propyl group,
 3-[3-{4-(N,N-dimethylaminocarbonyl)butoxy}phenyl]propyl
 10 group, and 3-[3-{4-(1-
 pyrrolidinylcarbonyl)butoxy}phenyl]propyl group;
 as well as 5-(aminocarbonyl)pentylloxy group, 5-(N-
 pentylaminocarbonyl)pentylloxy group, 7-
 (aminocarbonyl)heptyloxy group, 7-(N-
 15 pentylaminocarbonyl)heptyloxy group, 9-
 (aminocarbonyl)nonyloxy group, 9-(N-
 pentylaminocarbonyl)nonyloxy group, 11-
 (aminocarbonyl)undecyloxy group, 11-(N-
 pentylaminocarbonyl)undecyloxy group, 13-
 20 (aminocarbonyl)tridecyloxy group, and 13-(N-
 pentylaminocarbonyl)tridecyloxy group.

Particularly preferred are 7-(N,N-
 dimethylaminocarbonyl)heptyl group, 7-(N-
 ethylaminocarbonyl)heptyl group, 7-(N-
 25 isopropylaminocarbonyl)heptyl group, 7-(N-methyl-N-
 butylaminocarbonyl)heptyl group, 7-(N,N-
 diethylaminocarbonyl)heptyl group, 7-
 (piperidinocarbonyl)heptyl group, 7-{N-(2-

- furylmethyl)aminocarbonyl}heptyl group, 7-(N-methylaminocarbonyl)heptyl group, 7-(N-methyl-N-ethylaminocarbonyl)heptyl group, 7-(N-methyl-N-propylaminocarbonyl)heptyl group, 7-(N-methyl-N-isopropylaminocarbonyl)heptyl group, 7-(N-methyl-N-benzylaminocarbonyl)heptyl group, 7-(1-pyrrolidinylcarbonyl)heptyl group, 7-(morpholinocarbonyl)heptyl group, 9-(N,N-dimethylaminocarbonyl)nonyl group, 9-(N,N-diethylaminocarbonyl)nonyl group, 9-(N-methyl-N-butylaminocarbonyl)nonyl group, 9-(N-methyl-N-propylaminocarbonyl)nonyl group, 9-(morpholinocarbonyl)nonyl group, 10-(N,N-dimethylaminocarbonyl)decyl group, 7-{N-(2-hydroxyethyl)aminocarbonyl}heptyl group, 7-(N-propylaminocarbonyl)heptyl group, 7-(N-benzylaminocarbonyl)heptyl group, 7-{N-(2-phenylethyl)aminocarbonyl}heptyl group, 3-[3-{3-N-methylaminocarbonyl}propoxy}phenyl]propyl group, 3-[3-{3-(N,N-dimethylaminocarbonyl)propoxy}phenyl]propyl group, and 3-[3-{4-(1-pyrrolidinylcarbonyl)butoxy}phenyl]propyl group.
- It should however be noted that X^1 and X^2 are not a hydrogen atom at the same time. Particularly preferred cases are such that X^1 is a hydrogen atom and X^2 is any one of the groups listed above except a hydrogen atom, as well as where X^1 is any one of the groups listed above except a hydrogen atom and X^2 is a hydrogen atom.

Preferred examples of the compound represented by the

general formula (I) are listed below:

- 17 β -hydroxy-11 β -{10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyloxy}-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -{11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyloxy}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyloxy}-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -{11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyloxy}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-carboxy-13,13,14,14,14-pentafluorotetradecyloxy)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-carboxynonyloxy)-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -{12-(4,4,5,5,5-pentafluoropentylsulfinyl)dodecyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{10-(4,4,5,5,5-pentafluoropentylsulfonyl)decyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyl}-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -{12-(4,4,5,5,5-pentafluoropentylsulfonyl)dodecyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[10-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}decyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[11-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-{N-(5,5,6,6,6-

- pentafluorohexanoyl)amino}nonyl]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[10-{N-(5,5,6,6,6-
 pentafluorohexanoyl)amino}decyl]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(9-(4,4,5,5,5-
 5 pentafluoropentylsulfinyl)nonyloxy}-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(10-(4,4,5,5,5-
 pentafluoropentylsulfinyl)decyloxy}-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(11-(4,4,5,5,5-
 pentafluoropentylsulfinyl)undecyloxy}-5 α -androstan-3-one;
 10 17 β -hydroxy-11 β -(9-(4,4,5,5,5-
 pentafluoropentylsulfonyl)nonyloxy}-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(10-(4,4,5,5,5-
 pentafluoropentylsulfonyl)decyloxy}-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(11-(4,4,5,5,5-
 15 pentafluoropentylsulfonyl)undecyloxy}-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[9-{N-(4,4,5,5,5-
 pentafluoropentyl)aminocarbonyl}nonyloxy]-5 α -androstan-3-
 one;
 17 β -hydroxy-11 β -[10-{N-(4,4,5,5,5-
 20 pentafluoropentyl)aminocarbonyl}decyloxy]-5 α -androstan-3-
 one;
 17 β -hydroxy-11 β -[8-{N-(5,5,6,6,6-
 pentafluorohexanoyl)amino}octyloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[9-{N-(5,5,6,6,6-
 25 pentafluorohexanoyl)amino}nonyloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[4-{8-(4,4,5,5,5-
 pentafluoropentylsulfinyl)octyloxy}phenyl]-5 α -androstan-3-
 one;

- 17 β -hydroxy-11 β -[4-{9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyloxy}phenyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[4-{8-(4,4,5,5,5-pentafluoropentylsulfonyl)octyloxy}phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-{9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyloxy}phenyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -(4-[8-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}octyloxy]phenyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(4-[9-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}nonyloxy]phenyl)-5 α -
- 15 androstan-3-one;
- 17 β -hydroxy-11 β -(4-[7-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}heptyloxy]phenyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(4-[8-{N-(5,5,6,6,6-pentafluorohexanoyl)amino}octyloxy]phenyl)-5 α -androstan-3-
- 20 one;
- 17 β -hydroxy-11 β -(6-[4-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}phenyl]hexyl)-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -(5-[4-{N-(4,4,5,5,5-pentafluoropentyl)aminocarbonyl}phenyl]pentyloxy)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -tridecyloxy-5 α -androstan-3-one;

- 17 β -hydroxy-11 β -(11-carboxy-15,15,16,16,16-pentafluorohexadecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-{{2-hydroxy-3-(4,4,5,5,5-pentafluoropentylsulfinylethyloxy)propyl}oxy}phenyl]-5 α -
 5 androstan-3-one;
- 17 β -hydroxy-11 β -(4-hydroxy-9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(10-carboxy-14,14,15,15,15-pentafluoropentadecyloxy)-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -(9-carboxy-13,13,14,14,14-pentafluorotetradecyloxy)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(6-carboxy-10,10,11,11,11-pentafluoroundecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(10-carboxy-14,14,15,15,15-pentafluoropentadecyl)-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -(14-carboxy-18,18,19,19,19-pentafluorononadecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-carboxynonyloxy)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(6-carboxyhexyl)-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -(10-carboxydecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(14-carboxytetradecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[3-{4-(4-carboxybutyl)phenyl}propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[3-{4-(4-carboxy-8,8,9,9,9-pentafluorononyl)phenyl}propyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -(5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-(4,4,5,5,5-

- pentafluoropentylsulfinyl)nonyl}-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(13-(4,4,5,5,5-
 pentafluoropentylsulfinyl)tridecyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(4-hydroxy-10-(4,4,5,5,5-
 5 pentafluoropentylsulfinyl)decyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(4-hydroxy-15,15,16,16,16-
 pentafluorohexadecyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[9-{N-(4,4,5,5,5-
 pentafluoropentyl)aminocarbonyl)nonyl}-5 α -androstan-3-one;
 10 and
 17 β -hydroxy-11 β -[8-{N-(5,5,6,6,6-
 pentafluorohexanoyl)amino}octyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(11-(4,4,5,5,5-
 pentafluoropentylsulfinyl)undecyl)-5 α -androstan-3-one;
 15 17 β -hydroxy-7 β -(11-(4,4,5,5,5-
 pentafluoropentylsulfonyl)undecyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(5-(4,4,5,5,5-
 pentafluoropentylsulfinyl)pentyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(9-(4,4,5,5,5-
 20 pentafluoropentylsulfinyl)nonyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(5-carboxypentyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(7-carboxyheptyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(9-carboxynonyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(11-carboxyundecyl)-5 α -androstan-3-one;
 25 17 β -hydroxy-7 β -(13-carboxytridecyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 β -(9-carboxy-13,13,14,14,14-
 pentafluorotetradecyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(9-carboxy-13,13,14,14,14-

- pentafluorotetradecyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(9-methoxycarbonyl-13,13,14,14,14-pentafluorotetradecyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(5-carboxypentyloxy)-5 α -androstan-3-one;
 5 17 β -hydroxy-11 β -(7-carboxyheptyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(10-carboxydecyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(11-carboxyundecyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(13-carboxytridecyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(23-carboxytricosanyloxy)-5 α -androstan-3-
 10 one;
 17 β -hydroxy-7 α -{7-(N,N-dimethylaminocarbonyl)heptyl}-5 α -
 androstan-3-one;
 17 β -hydroxy-7 α -{7-(N-ethylaminocarbonyl)heptyl}-5 α -
 androstan-3-one;
 15 17 β -hydroxy-7 α -[7-{N-(cyclopropylmethyl)aminocarbonyl}heptyl]-5 α -androstan-3-
 one;
 17 β -hydroxy-7 α -[7-{N-(cyclohexylmethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 20 17 β -hydroxy-7 α -[7-(N-butylaminocarbonyl)heptyl]-5 α -
 androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-(isopropylaminocarbonyl)heptyl)-5 α -
 androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-t-butylaminocarbonyl)heptyl]-5 α -
 25 androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-cyclohexylaminocarbonyl)heptyl]-5 α -
 androstan-3-one;
 17 β -hydroxy-7 α -[7-{N-(3-

- hydroxypropyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-butylaminocarbonyl)heptyl]-
 5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N,N-diethylaminocarbonyl)heptyl]-5 α -
 5 androstan-3-one;
 17 β -hydroxy-7 α -[7-(piperidinocarbonyl)heptyl]-5 α -androstan-
 3-one;
 17 β -hydroxy-7 α -[7-{N-(4-t-
 butylbenzyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 10 17 β -hydroxy-7 α -[7-{N-(2,2-
 diphenylethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-{N-(2-furylmethyl)aminocarbonyl}heptyl]-
 5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-{7-(N-methylaminocarbonyl)heptyl]-5 α -
 15 androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-ethylaminocarbonyl)heptyl]-
 5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-propylaminocarbonyl)heptyl]-
 5 α -androstan-3-one;
 20 17 β -hydroxy-7 α -[7-(N-methyl-N-
 isopropylaminocarbonyl)heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-benzylaminocarbonyl)heptyl]-
 5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(1-pyrrolidinylcarbonyl)heptyl]-5 α -
 25 androstan-3-one;
 17 β -hydroxy-7 α -[7-(morpholinocarbonyl)heptyl]-5 α -androstan-
 3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-t-butylaminocarbonyl)heptyl]-

- 5 α -androstan-3-one;
- 17 β -hydroxy-7 β -[7-(N-cyclopropylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[6-(N,N-dimethylaminocarbonyl)hexyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[6-(N,N-diethylaminocarbonyl)hexyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[6-(piperidinocarbonyl)hexyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-(N,N-dimethylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-(N,N-diethylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-(N-methyl-N-butylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-(N-benzylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-{N-(2-hydroxyethyl)aminocarbonyl}octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-(piperidinocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N,N-dimethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N,N-diethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(1-pyrrolidinylcarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-methyl-N-ethylaminocarbonyl)nonyl]-5 α -

- androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-methyl-N-butylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-benzylaminocarbonyl)nonyl]-5 α -
- 5 androstan-3-one;
- 17 β -hydroxy-7 α -[9-(piperidinocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-(2-hydroxyethyl)aminocarbonyl)nonyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[9-(N-methyl-N-propylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(morpholinocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-(N,N-dimethylaminocarbonyl)decyl]-5 α -
- 15 androstan-3-one;
- 17 β -hydroxy-7 α -[10-(N,N-diethylaminocarbonyl)decyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-(N-methyl-N-ethylaminocarbonyl)decyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-7 α -[10-N-methyl-N-propylaminocarbonyl)decyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-(N-methyl-N-butylaminocarbonyl)decyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-(morpholinocarbonyl)decyl]-5 α -androstan-
- 25 3-one;
- 17 β -hydroxy-7 α -[11-(N,N-dimethylaminocarbonyl)undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[11-(N,N-diethylaminocarbonyl)undecyl]-5 α -

- androstan-3-one;
- 17 β -hydroxy-7 α -[11-(piperidinocarbonyl)undecyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[11-(N-benzylaminocarbonyl)undecyl]-5 α -
5 androstan-3-one;
- 17 β -hydroxy-7 α -[11-(N-methyl-N-butylaminocarbonyl)undecyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[11-{N-(2-
hydroxyethyl)aminocarbonyl}undecyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[7-{N-(2-hydroxyethyl)aminocarbonyl}heptyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-propylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-hexylaminocarbonyl)heptyl]-5 α -
15 androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-isopentylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-isobutylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 20 17 β -hydroxy-7 α -[7-(N-neopentylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-{N-(3-pentyl)aminocarbonyl}heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N,N-dihexylaminocarbonyl)heptyl]-5 α -
25 androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-phenylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-benzylaminocarbonyl)heptyl]-5 α -

- androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-(2-phenylethyl)aminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(7-carboxyheptyl)-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -(8-carboxyoctyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-carboxynonyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(11-carboxyundecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[5-(aminocarbonyl)pentyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[5-(N-pentylaminocarbonyl)pentyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(aminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-pentylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[9-(aminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(aminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-pentylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -[9-(N-pentylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[11-(aminocarbonyl)undecyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[11-(aminocarbonyl)undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[11-(N-pentylaminocarbonyl)undecyl]-5 α -androstan-3-one;

- 17 β -hydroxy-11 β -[11-(N-pentylaminocarbonyl)undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[13-(aminocarbonyl)tridecyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[13-(N-pentylaminocarbonyl)tridecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{7-(N,N-dimethylaminocarbonyl)heptyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-{7-(N-methylaminocarbonyl)heptyl}]-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -[7-(N-methyl-N-ethylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(N-methyl-N-propylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -[7-(morpholinocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[8-(N,N-dimethylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[8-(N-methylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -[8-(N-methyl-N-ethylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[8-(N-methyl-N-propylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[8-(morpholinocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N,N-dimethylaminocarbonyl)nonyl]-5 α -androstan-3-one;

- 17 β -hydroxy-11 β -[9-(N,N-diethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N-methyl-N-butylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[9-(N-benzylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(piperidinocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-{N-(2-hydroxyethyl)aminocarbonyl}nonyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -[10-(4,4,5,5,5-pentafluoropentylsulfanyl)decyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -(7-hydroxyheptyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -(8-hydroxyoctyl)-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -(9-hydroxynonyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-7 α -[7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -[4-(carboxymethoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(3-carboxypropoxy)phenyl]-5 α -androstan-3-

- one;
- 17 β -hydroxy-11 β -[4-(3-carboxypropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(7-carboxyheptyloxy)phenyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[4-(7-carboxyheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(carbamoylmethoxy)phenyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[4-(carbamoylmethoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(3-carbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-(3-carbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[4-(7-carbamoylheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-(7-carbamoylheptyloxy)phenyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-7 α -[4-(3-N-pentylcarbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-(3-N-pentylcarbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(7-N-pentylcarbamoylheptyloxy)phenyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[4-(7-N-pentylcarbamoylheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-methoxyphenyl]-5 α -androstan-3-one;

- 17 β -hydroxy-11 β -[4-methoxyphenyl]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentylloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyloxy]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentylloxy}phenyl]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[4-{7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy}phenyl]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)pentylloxy}phenyl]-5 α -androstan-3-one;
 17 β -hydroxy-11 β -[4-{7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy}phenyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[3-{3-(3-carboxypropoxy)phenyl}propyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[3-{3-(4-carboxybutoxy)phenyl}propyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[3-[3-{3-(N-

- methylaminocarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{3-(N,N-dimethylaminocarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[3-[3-{3-(1-pyrrolidinylcarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{4-(N-methylaminocarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[3-[3-{4-(N,N-dimethylaminocarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[3-[3-{4-(1-pyrrolidinylcarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[5-(aminocarbonyl)pentyloxy]-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -[5-(N-pentylaminocarbonyl)pentyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(aminocarbonyl)heptyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(N-pentylaminocarbonyl)heptyloxy]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[9-(aminocarbonyl)nonyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N-pentylaminocarbonyl)nonyloxy]-5 α -

- androstan-3-one;
- 17 β -hydroxy-11 β -[11-(aminocarbonyl)undecyloxy]-5 α -androstan-one;
- 17 β -hydroxy-11 β -[11-(N-pentylaminocarbonyl)undecyloxy]-5 α -
- 5 androstan-3-one;
- 17 β -hydroxy-11 β -[13-(aminocarbonyl)tridecyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[13-(N-pentylaminocarbonyl)tridecyloxy]-5 α -androstan-3-one;
- 10 more preferred are the following:
- 17 β -hydroxy-11 β -{10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyloxy}-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -{11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyloxy}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyloxy}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -{11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyloxy}-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -(9-carboxy-13,13,14,14,14-pentafluorotetradecyloxy)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-carboxynonyloxy)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -{11-(4,4,5,5,5-pentafluoropentylsulfinyl)undecyl}-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -{11-(4,4,5,5,5-pentafluoropentylsulfonyl)undecyl}-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -{5-(4,4,5,5,5-

- pentafluoropentylsulfinyl)pentyl}-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(5-carboxypentyl)-5 α -androstan-3-one;
 5 17 β -hydroxy-7 α -(7-carboxyheptyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(9-carboxynonyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(11-carboxyundecyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(13-carboxytridecyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(9-carboxy-13,13,14,14,14-pentafluorotetradecyl)-5 α -androstan-3-one;
 10 17 β -hydroxy-11 β -(9-carboxy-13,13,14,14,14-pentafluorotetradecyl)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(9-methoxycarbonyl-13,13,14,14,14-pentafluorotetradecyl)-5 α -androstan-3-one;
 15 17 β -hydroxy-11 β -(5-carboxypentyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(7-carboxyheptyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(10-carboxydecyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(11-carboxyundecyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-11 β -(13-carboxytridecyloxy)-5 α -androstan-3-one;
 20 17 β -hydroxy-11 β -(23-carboxytricosanyloxy)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(7-(N,N-dimethylaminocarbonyl)heptyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -(7-(N-ethylaminocarbonyl)heptyl)-5 α -androstan-3-one;
 25 17 β -hydroxy-7 α -(7-(N-(cyclopropylmethyl)aminocarbonyl)heptyl)-5 α -androstan-3-one;

- 17 β -hydroxy-7 α -[7-{N-(cyclohexylmethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-butylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[7-(N-(isopropylaminocarbonyl)heptyl)-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-t-butylaminocarbonyl)heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-cyclohexylaminocarbonyl)heptyl]-5 α -
- 10 androstan-3-one;
 17 β -hydroxy-7 α -[7-{N-(3-hydroxypropyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-butylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[7-(N,N-diethylaminocarbonyl)heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(piperidinocarbonyl)heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-{N-(4-t-butylbenzyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-{N-(2,2-diphenylethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-{N-(2-furylmethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -[7-(N-methylaminocarbonyl)heptyl]-5 α -androstan-3-one;
 17 β -hydroxy-7 α -[7-(N-methyl-N-ethylaminocarbonyl)heptyl]-5 α -androstan-3-one;

- 17 β -hydroxy-7 α -[7-(N-methyl-N-propylaminocarbonyl)heptyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-methyl-N-
isopropylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[7-(N-methyl-N-benzylaminocarbonyl)heptyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(1-pyrrolidinylcarbonyl)heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-(morpholinocarbonyl)heptyl]-5 α -androstan-
10 3-one;
- 17 β -hydroxy-7 α -[7-(N-methyl-N-t-butylaminocarbonyl)heptyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-cyclopropylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 15 17 β -hydroxy-7 α -[6-(N,N-dimethylaminocarbonyl)hexyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[6-(N,N-diethylaminocarbonyl)hexyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[6-(piperidinocarbonyl)hexyl]-5 α -androstan-
20 3-one;
- 17 β -hydroxy-7 α -[8-(N,N-dimethylaminocarbonyl)octyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[8-(N,N-diethylaminocarbonyl)octyl]-5 α -
androstan-3-one;
- 25 17 β -hydroxy-7 α -[8-(N-methyl-N-butylaminocarbonyl)octyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[8-(N-benzylaminocarbonyl)octyl]-5 α -
androstan-3-one;

- 17 β -hydroxy-7 α -[8-{N-(2-hydroxyethyl)aminocarbonyl}octyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[8-(piperidinocarbonyl)octyl]-5 α -androstan-
3-one;
- 5 17 β -hydroxy-7 α -[9-(N,N-dimethylaminocarbonyl)nonyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N,N-diethylaminocarbonyl)nonyl]-5 α -
androstan-3-one;
- 10 17 β -hydroxy-7 α -[9-(1-pyrrolidinylcarbonyl)nonyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-methyl-N-ethylaminocarbonyl)nonyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-methyl-N-butylaminocarbonyl)nonyl]-5 α -
androstan-3-one;
- 15 17 β -hydroxy-7 α -[9-(N-benzylaminocarbonyl)nonyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[9-(piperidinocarbonyl)nonyl]-5 α -androstan-
3-one;
- 17 β -hydroxy-7 α -[9-{N-(2-hydroxyethyl)aminocarbonyl}nonyl]-
20 5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-methyl-N-propylaminocarbonyl)nonyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(morpholinocarbonyl)nonyl]-5 α -androstan-
3-one;
- 25 17 β -hydroxy-7 α -[10-(N,N-dimethylaminocarbonyl)decyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[10-(N,N-diethylaminocarbonyl)decyl]-5 α -
androstan-3-one;

- 17 β -hydroxy-7 α -[10-(N-methyl-N-ethylaminocarbonyl)decyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-N-methyl-N-propylaminocarbonyl)decyl]-
5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[10-(N-methyl-N-butylaminocarbonyl)decyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-(morpholinocarbonyl)decyl]-5 α -androstan-
3-one;
- 17 β -hydroxy-7 α -[11-(N,N-dimethylaminocarbonyl)undecyl]-5 α -
10 androstan-3-one;
- 17 β -hydroxy-7 α -[11-(N,N-diethylaminocarbonyl)undecyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[11-(piperidinocarbonyl)undecyl]-5 α -
androstan-3-one;
- 15 17 β -hydroxy-7 α -[11-(N-benzylaminocarbonyl)undecyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[11-(N-methyl-N-butylaminocarbonyl)undecyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[11-{N-(2-
20 hydroxyethyl)aminocarbonyl}undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-{N-(2-hydroxyethyl)aminocarbonyl}heptyl]-
5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-propylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 25 17 β -hydroxy-7 α -[7-(N-hexylaminocarbonyl)heptyl]-5 α -
androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-isopentylaminocarbonyl)heptyl]-5 α -
androstan-3-one;

- 17 β -hydroxy-7 α -[7-(N-isobutylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-neopentylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[7-{N-(3-pentyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N,N-dihexylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-phenylaminocarbonyl)heptyl]-5 α -
- 10 androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-benzylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-{N-(2-phenylethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -(7-carboxyheptyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(8-carboxyoctyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(9-carboxynonyl)-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -(11-carboxyundecyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[5-(aminocarbonyl)pentyl]-5 α -androstan-3-
- 20 one;
- 17 β -hydroxy-7 α -[5-(N-pentylaminocarbonyl)pentyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(aminocarbonyl)heptyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -[7-(N-pentylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(aminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(aminocarbonyl)nonyl]-5 α -androstan-3-

- one;
- 17 β -hydroxy-7 α -[9-(N-pentylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N-pentylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[11-(aminocarbonyl)undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[11-(aminocarbonyl)undecyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[11-(N-pentylaminocarbonyl)undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[11-(N-pentylaminocarbonyl)undecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[13-(aminocarbonyl)tridecyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[13-(N-pentylaminocarbonyl)tridecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(N,N-dimethylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -[7-{7-(N-methylaminocarbonyl)heptyl}-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(N-methyl-N-ethylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(N-methyl-N-propylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[7-(morpholinocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[8-(N,N-dimethylaminocarbonyl)octyl]-5 α -

- androstan-3-one;
- 17 β -hydroxy-11 β -[8-(N-methylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[8-(N-methyl-N-ethylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[8-(N-methyl-N-propylaminocarbonyl)octyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[8-(morpholinocarbonyl)octyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -[9-(N,N-dimethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N,N-diethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N-methyl-N-butylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -[9-(N-benzylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(piperidinocarbonyl)nonyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -[9-{N-(2-hydroxyethyl)aminocarbonyl}nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[10-(4,4,5,5,5-pentafluoropentylsulfanyl)decyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -(7-hydroxyheptyl)-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -(8-hydroxyoctyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -(9-hydroxynonyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyl]-5 α -androstan-3-one;

- 17 β -hydroxy-7 α -[13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-7 α -[9-(4,4,5,5,5-pentafluoropentylsulfonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(carboxymethoxy)phenyl]-5 α -androstan-3-
- 10 one;
- 17 β -hydroxy-7 α -[4-(3-carboxypropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-(3-carboxypropoxy)phenyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[4-(7-carboxyheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-(7-carboxyheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(carbamoylmethoxy)phenyl]-5 α -androstan-3-
- 20 one;
- 17 β -hydroxy-7 α -[4-(carbamoylmethoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(3-carbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-11 β -[4-(3-carbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(7-carbamoylheptyloxy)phenyl]-5 α -androstan-3-one;

- 17 β -hydroxy-11 β -[4-(7-carbamoylheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(3-N-pentylcarbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[4-(3-N-pentylcarbamoylpropoxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-(7-N-pentylcarbamoylheptyloxy)phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-(7-N-pentylcarbamoylheptyloxy)phenyl]-
- 10 5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[4-methoxyphenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-methoxyphenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentylloxy]-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -[7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy]-5 α -androstan-3-one;
- 20 17 β -hydroxy-11 β -[13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentylloxy}phenyl]-5 α -androstan-3-
- 25 one;
- 17 β -hydroxy-11 β -[4-{7-(4,4,5,5,5-pentafluoropentylsulfinyl)heptyloxy}phenyl]-5 α -androstan-3-one;


- 17 β -hydroxy-11 β -[4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)pentyloxy}phenyl]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[4-{7-(4,4,5,5,5-pentafluoropentylsulfonyl)heptyloxy}phenyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-{3-(3-carboxypropoxy)phenyl}propyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[3-{3-(4-carboxybutoxy)phenyl}propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{3-(N-methylaminocarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 15 17 β -hydroxy-7 α -[3-[3-{3-(N,N-dimethylaminocarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{3-(1-pyrrolidinylcarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-7 α -[3-[3-{4-(N-methylaminocarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{4-(N,N-dimethylaminocarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-
- 25 one;
- 17 β -hydroxy-7 α -[3-[3-{4-(1-pyrrolidinylcarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-one;

- 17 β -hydroxy-11 β -[5-(aminocarbonyl)pentyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[5-(N-pentylaminocarbonyl)pentyloxy]-5 α -androstan-3-one;
- 5 17 β -hydroxy-11 β -[7-(aminocarbonyl)heptyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[7-(N-pentylaminocarbonyl)heptyloxy]-5 α -androstan-3-one;
- 10 17 β -hydroxy-11 β -[9-(aminocarbonyl)nonyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N-pentylaminocarbonyl)nonyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[11-(aminocarbonyl)undecyloxy]-5 α -androstan-3-one;
- 15 17 β -hydroxy-11 β -[11-(N-pentylaminocarbonyl)undecyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[13-(aminocarbonyl)tridecyloxy]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[13-(N-pentylaminocarbonyl)tridecyloxy]-5 α -androstan-3-one;
- 20 particularly preferred are the following:
- 17 β -hydroxy-7 α -{7-(N,N-dimethylaminocarbonyl)heptyl}-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -{7-(N-ethylaminocarbonyl)heptyl}-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -[7-(N-(isopropylaminocarbonyl)heptyl)-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-methyl-N-butylaminocarbonyl)heptyl]-


- 5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N,N-diethylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(piperidinocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-{N-(2-furylmethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-{7-(N-methylaminocarbonyl)heptyl}-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[7-(N-methyl-N-ethylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-methyl-N-propylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-methyl-N-
- 15 isopropylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-methyl-N-benzylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(1-pyrrolidinylcarbonyl)heptyl]-5 α -androstan-3-one;
- 20 17 β -hydroxy-7 α -[7-(morpholinocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N,N-dimethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N,N-diethylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 25 17 β -hydroxy-7 α -[9-(N-methyl-N-butylaminocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(N-methyl-N-propylaminocarbonyl)nonyl]-


- 5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[9-(morpholinocarbonyl)nonyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[10-(N,N-dimethylaminocarbonyl)decyl]-5 α -
5. androstan-3-one;
- 17 β -hydroxy-7 α -[7-{N-(2-hydroxyethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-(N-propylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 10 17 β -hydroxy-7 α -[7-(N-benzylaminocarbonyl)heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[7-{N-(2-phenylethyl)aminocarbonyl}heptyl]-5 α -androstan-3-one;
- 17 β -hydroxy-11 β -[9-(N,N-diethylaminocarbonyl)nonyl]-5 α -
- 15 androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{3-(N-methylaminocarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{3-(N,N-
- 20 dimethylaminocarbonyl)propoxy}phenyl]propyl]-5 α -androstan-3-one;
- 17 β -hydroxy-7 α -[3-[3-{4-(1-pyrrolidinylcarbonyl)butoxy}phenyl]propyl]-5 α -androstan-3-one.

25 The structures of these compounds are shown below:




















$\{-Ar-A-R^1\}$;








$\{-ArA-R^1\}$;


































{-Ar-A-R¹};



































$\{-ArA-R^1\}$;





$\xi - \text{ArA-R}^1 ;$


If the compounds represented by the general formula (I) contain one or more asymmetric carbon atom in their molecule, those compounds which have R and S configurations

as absolute configuration for each of the contained asymmetric carbon atoms, as well as all mixtures of those compounds at any proportions are included within the scope of the invention.

5 Speaking of the substance of the invention which acts as antagonist against but not as agonist for the androgen receptor, the expression "not acting as agonist" means that in the following androgen receptor gene assay, the transcriptional activity value of the substance at any
10 concentration of 0.1 nmol/L - 10 μ mol/L is from one to five times the transcriptional activity value for no addition of the substance which is taken as unity:

Twenty-four hours before transfection, 1.0×10^5 HeLa cells (purchased from Dainippon Pharmaceutical Co., Ltd.)
15 are cultured in phenol red free Dulbecco's modified Eagle medium (DMEM) containing 5% of charcoal-treated FBS (DCC-FBS) in 12-well microplates. Five hundred nanograms/well of MMTV-Luc vector (the reporter plasmid of luciferase having mouse tumor long terminal repeats containing the
20 androgen response element: GM-CAT vector (A.T.C.C. No. 67282) purchased from A.T.C.C. provided that the chloramphenicol acetyl transferase gene was replaced by the firefly luciferase gene), 100 ng/well of pSG5-hAR (the expression vector of the human androgen receptor which
25 harbors the androgen receptor gene under the control of SV40 promoter) and 5 ng/well of Renilla Luc vector (a vector for internal standard incorporating the sea pansy luciferase gene) are transfected into the HeLa cells. The

transfection is performed in a liquid culture of the phenol red free DMEM using 3 mL/well of lipofectoamine (GibcoBRL). Nine hours after the transfection, the liquid culture is replaced by phenol red free DMEM/3% DCC-FBS containing 10 μ mol/L of a compound of the invention which is represented by the general formula (I) or the substance of the invention which acts as antagonist against but not as agonist for the androgen receptor. The transcriptional activity value is measured 48 hours after the replacement of the liquid culture. Transcriptional activity is measured with a dual-luciferase reporter assay system (Promega). The transcriptional activity value is defined as the value for firefly luciferase divided by the value for sea pansy luciferase. To implement this assay, reference may be had to J. Biol. Chem., vol. 270, pp. 19998-20003, 1995.

WO97/49709 mentions hydroxyflutamide (the essence of the in vivo activity of flutamide) and bicaltamide as substances that act as antagonist against but not as agonist for the androgen receptor; however, according to the definition given in that publication, the expression "not acting as agonist" means that in an androgen reporter gene assay using CV-1 cells, the agonist efficiency value represented by the following formula is 0 - 20% at a concentration of 10 μ mol/L or above and this definition is clearly and strictly distinguished from the definition of the expression "not acting as agonist" which is given in the present invention:

Agonist efficiency (%) = (transcriptional activity value of screened non-steroid compound)/(maximum transcriptional activity value by DHT) x 100

In the androgen receptor reporter gene assay used in
5 defining the expression "not acting as agonist" in the invention, each of hydroxyflutamide and bicaltamide was found to act as agonist at a concentration of 10 μ mol/L (see Example 1 in this specification).

The expression "acting as antagonist" means that in
10 the following androgen receptor gene assay, the transcriptional activity value of 0.1 nmol/L of dihydrotestosterone (DHT) is inhibited to 0 - 50% at any concentration of 0.1 nmol/L - 10 μ mol/L:

Twenty-four hours before transfection, 1.0×10^5 HeLa
15 cells are cultured in phenol red free DMEM/5% DCC-FBS on 12-well microplates. Five hundred nanograms/well of MMTV-Luc vector, 100 ng/well of pSG5/hAR and 5 ng/well of Renilla Luc vector are transfected into the HeLa cells. The transfection is performed in a liquid culture of the
20 phenol red free DMEM using 3 mL/well of lipofectoamine. Nine hours after the transfection, the liquid culture is replaced by phenol red free DMEM/3% DCC-FBS containing 0.1 nmol/L of DHT and 1.0 mol/L of a compound of the invention which is represented by the general formula (I) or the
25 substance of the invention which acts as antagonist against but not as agonist for the androgen receptor. The transcriptional activity value is measured 48 hours after the replacement of the liquid culture. Transcriptional

activity is measured with a dual-luciferase reporter assay system. The transcriptional activity value is defined as the value for firefly luciferase divided by the value for sea pansy luciferase. To implement this assay, reference
5 may be had to J. Biol. Chem., vol. 270, pp. 19998-20003, 1995.

Specific examples of the substance of the invention which acts as antagonist against but not as agonist for the androgen receptor may include compounds of the invention
10 which are represented by the general formula (I).

The compounds of the invention which are represented by the general formula (I) and the substance of the invention which acts as antagonist against but not as agonist for the androgen receptor can also be obtained as
15 their pharmaceutically acceptable salts. Pharmaceutically acceptable salts include inorganic acid salts such as hydrochlorides, hyrobromides, hydroiodides, sulfates and phosphates; organic acid salts such as formates, acetates, oxalates, maleates, fumarates, methanesulfonates,
20 benzenesulfonates, p-toluenesulfonates, succinates, malonates, citrates, gluconates, mandelates, benzoates, salicylates, trifluoroacetates, tartrates, propionates and glutarates; inorganic base salts such as sodium salts, potassium salts, magnesium salts and zinc salts; and
25 organic base salts such as ammonium salts.


The compounds of the invention which are represented by the general formula (I), their pharmaceutically acceptable salts, as well as the substance of the invention

which acts as antagonist against but not as agonist for the androgen receptor, and its pharmaceutically acceptable salts can also be obtained as their prodrugs. The prodrugs mean those compounds which undergo rapid transformation in living body to generate, typically by hydrolysis in the blood, the compounds of the invention which are represented by the general formula (I), their pharmaceutically acceptable salts, as well as the substance of the invention which acts as antagonist against but not as agonist for the androgen receptor, and its pharmaceutically acceptable salts. T. Higuchi and V. Stella give detailed accounts of the concept of prodrugs in "Prodrugs as Novel Delivery Systems", vol. 14 of the A.C.S. Symposium Series, American Chemical Society (1975). These prodrugs may or may not have activity on their own but they usually have little activity. Reference may also be had to D.E.V. Wilman, "Prodrugs in Cancer Chemotherapy" in Biochemical Society Transactions, vol. 14, pp. 375-382, the 615th Meeting, Belfast, 1986, and V.J. Stella et al., "Prodrugs: Chemical Methods for Targeted Drug Delivery" in Directed Drug Delivery, ed. by R. Borchardt et al., pp. 247-267, Humana Press, 1985. If compounds of the invention which are represented by the general formula (I) have the -COOH partial structure, specific examples of prodrugs include esters, carbonates, carbamates, etc. of such compounds.

The compounds of the invention which are represented by the general formula (I) can typically be produced by process A to process W, process B' to process L', process

S' to process W', process U", process W" and process W'" that are set forth below, or depending on the end compound, partial modifications of process A to process W, process B' to process L', process S' to process W', process U", process W" and process W'" may be employed.

In the chemical formulae listed in process A to process W, process B' to process L', process S' to process W', process U", process W" and process W'", R² represents the general formula (IV)



(wherein G² represents a single bond, a straight-chained or branched alkylene group having 1 - 26 carbon atoms, a straight-chained or branched alkenylene group having 2 - 26 carbon atoms or a straight-chained or branched alkynylene group having 2 - 26 carbon atoms; E, J, Y, L, Q² and Z have the same meanings as defined above, provided that R⁷ and R⁸ in Q² are preferably a hydrogen atom); R³ represents a substituted silyl group, preferably t-butyldimethylsilyl group; X³ represents a halogen atom or a substituted sulfonate group, preferably p-toluenesulfonate group or methanesulfonate group; R¹² represents a straight-chained or branched alkyl group having 1 - 6 carbon atoms, preferably methyl group and ethyl group; R^e and R^f, when taken together with the carbon atoms in 3- and 17-positions to which they are bound, represent protected -(C=O)-, preferably 1,3-dioxane, 1,3-dioxolan, 1,3-dithian, etc., particularly preferably 1,3-dioxolan, etc.; L² represents a straight-chained or branched lower alkylene group having 1 - 10

carbon atoms, preferably ethane-1,2-diyl group, propane-1,3-diyl group and butane-1,4-diyl group.

R^4 represents the general formula (V)



- 5 (wherein G^3 represents a straight-chained or branched alkylene group having 1 - 27 carbon atoms, a straight-chained or branched alkenylene group having 2 - 27 carbon atoms or a straight-chained or branched alkynylene group having 2 - 27 carbon atoms; E, J, Y, L, Q^2 and Z have the same meanings as defined above); R^5 represents a halogen atom, preferably a bromine atom or an iodine atom; R^6 represents a substituted silyl group, preferably trimethylsilyl group; R^{13} represents a straight-chained or branched alkyl group having 1 - 6 carbon atoms that may optionally be substituted by a halogen atom, preferably trifluoromethyl group or 1,1,2,2,3,3,4,4,4-nonafluorobutyl group; R^{14} represents a group represented by $-MgR^5$, $-ZnR^5$ or $-Sn(R^7)_3$, preferably a group represented by $-Sn(R^7)_3$; G^4 represents a straight-chained or branched alkylene group having 1 - 30 carbon atoms, a straight-chained or branched alkenylene group having 2 - 30 carbon atoms or a straight-chained or branched alkynylene group having 2 - 30 carbon atoms; the wavy line represents a single bond of trans configuration or cis configuration, preferably trans configuration, with respect to the double bond.

Process A is for producing compound (6) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula

(II) in which Ar is a single bond, A is -O- and R¹ is -CH₂-CH=CH-CH₂-R², X² is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the

5 dashed line together with the solid line is a single bond or a double bond; compound (7) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R¹ is -(CH₂)₄-R², X² is a hydrogen

10 atom, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; compound (9)


15 represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R¹ is -(CH₂)₄-G²-S(O)-Z, X² is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the

20 dashed line together with the solid line is a single bond or a double bond; compound (10) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R¹ is -(CH₂)₄-G²-S(O)₂-Z, X² is a

25 hydrogen atom, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; and

compound (147) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{R}^2$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a double bond.

Process A



Step A1 is for producing compound (2) and implemented by reacting compound (1) with compound (133) in an inert solvent in the presence of a base.

5 The inert solvent to be used is not limited in any particular way as long as it does not participate in the

reaction of interest; examples are halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, preferably halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, with dichloromethane being more preferred. The base to be used may be exemplified by organic bases such as diisopropylethylamine, 4-dimethylaminopyridine, pyridine, triethylamine and N-methylmorpholine, preferably diisopropylethylamine. The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 50°C, preferably 10°C - 30°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes to 24 hours, preferably 30 minutes - 15 hours.

Step A2 is for synthesizing compound (3) and implemented by reacting compound (2) with a reducing agent in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and alcoholic solvents such as methanol and ethanol, preferably ether and tetrahydrofuran, with ether being more preferred. The reducing agent to be used may be exemplified by: metal hydrogen complex compounds such as aluminum lithium hydride,

trimethoxyaluminum lithium hydride, tri-t-butoxyaluminum
 lithium hydride, aluminum lithium hydride-trichloroaluminum
 (alane), aluminum lithium hydride-boron trifluoride,
 aluminum hydride magnesium chloride, magnesium aluminum
 5 hydride, sodium aluminum hydride, sodium triethoxyaluminum
 hydride, sodium bis(methoxyethoxy)aluminum hydride, sodium
 boron hydride, sodium boron hydride-palladium/carbon,
 sodium boron hydrogensulfide, sodium boron hydrogencyanide,
 sodium trimethoxyboron hydride, lithium boron hydride,
 10 lithium boron hydrogencyanide, lithium triethylboron
 hydride, lithium tri-s-butylboron hydride, lithium tri-t-
 butylboron hydride, calcium boron hydride, potassium boron
 hydride, potassium triisopropoxyboron hydride, potassium
 tri-s-butylboron hydride, zinc boron hydride,
 15 tetramethylammonium boron hydride, and tetra-n-
 butylammonium cyanoboron hydride; metal hydrides such as
 diisobutylaluminum hydride, triphenyltin hydride, tri-n-
 butyltin hydride, diphenyltin hydride, di-n-butyltin
 hydride, triethyltin hydride, trimethyltin hydride,
 20 trichlorosilane/tri-n-butylamine, trichlorosilane/tri-n-
 propylamine, triethylsilane, trimethylsilane,
 diphenylsilane, phenylsilane, polymethylhydrosiloxane,
 dimethylphenylsilane, di-n-butylsilane, and
 methylphenylsilane; borane derivatives such as diborane,
 25 dimethylamine-borane, trimethylamine-borane,
 ethylenediamine-borane, pyridine-borane, dimethylsulfide-
 borane, 2,3-dimethyl-2-butylborane (thexylborane), bis-3-
 methyl-2-butylborane (disiamylborane),

diisopinocanepherylborane, dicyclohexylborane, and 9-borabicyclo[3,3,1]nonane (9-BBN); preferred examples are metal hydrogen complex compounds such as aluminum lithium hydride, trimethoxyaluminum lithium hydride, tri-t-
5 butoxyaluminum lithium hydride, aluminum lithium hydride-trichloroaluminum (alane), aluminum lithium hydride-boron trifluoride, aluminum hydride magnesium chloride, magnesium aluminum hydride, sodium aluminum hydride, sodium triethoxyaluminum hydride, sodium
10 bis(methoxyethoxy)aluminum hydride, sodium boron hydride, sodium boron hydride-palladium/carbon, sodium boron hydrogensulfide, sodium boron hydrogencyanide, sodium trimethoxyboron hydride, lithium boron hydride, lithium boron hydrogencyanide, lithium triethylboron hydride,
15 lithium tri-s-butylboron hydride, lithium tri-t-butylboron hydride, calcium boron hydride, potassium boron hydride, potassium triisopropoxyboron hydride, potassium tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium boron hydride, and tetra-n-butylammonium cyanoboron hydride,
20 with aluminum lithium hydride being more preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of -30°C ~ 100°C, preferably 0°C ~ 70°C. The reaction time which varies with the reaction temperature and the like is typically in the
25 range of 10 minutes - 48 hours, preferably 30 minutes - 24 hours.

As a by-product of this step, there is formed a compound having the hydroxyl group in 11-position of

compound (3) oriented in α configuration and this compound may be used to prepare compounds having X^1 in compound (6), compound (7), compound (9) and compound (10) oriented in α configuration.

5 Step A3 is for producing compound (4) and implemented by reacting compound (3) with a base in an inert solvent to make a salt of compound (3) and then reacting it with compound (134) in an inert solvent.

10 The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples are halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, 15 xylene, quinoline and chlorobenzene, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, and N-methylpyrrolidone; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, 20 as well as dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, and N-methylpyrrolidone. The base to be used may be exemplified by metal hydrides such as sodium hydride, potassium hydride and calcium hydride, alkyl lithium compounds such as 25 methyl lithium, ethyl lithium, n-butyl lithium and t-butyl lithium, metal hydroxides such as lithium hydroxide, sodium hydroxide, potassium hydroxide, potassium hydroxide, calcium hydroxide, barium hydroxide and cesium hydroxide,

metal amides such as sodium amide, potassium bistrimethylsilylamide, sodium bistrimethylsilylamide and lithium diisopropylamide, amines such as triethylamine, diisopropylethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene, 5 pyridine, dimethylamionopyridine and pyrazine, as well as sodium tetraborate, sodium iodide, lithium hexamethyldisilazane, sodium hexamethyldisilazane and potassium hexamethyldisilazane; preferred examples are metal hydrides such as sodium hydride, potassium hydride 10 and calcium hydride, and alkyl lithium compounds such as methyllithium, ethyllithium, n-butyllithium and t-butyllithium. The reaction temperature which varies with the type of solvent and the like is typically in the range of -30°C ~ 100°C, preferably 0°C ~ 70°C. The reaction 15 temperature which varies with the reaction temperature and the like is typically in the range of 10 minutes - 48 hours, preferably 30 minutes - 24 hours.

Step A4 is for producing compound (5) and implemented by reacting compound (4) with compound (135) in an inert 20 solvent in the presence of an organometallic catalyst.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are halogen-containing solvents such as dichloromethane and chloroform, ether 25 solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, with dichloromethane and dimethoxyethane being more preferred.

The organometallic catalyst to be used is preferably
benzylidene-bis(tricyclohexylphosphine)-dichlororuthenium.
The reaction temperature which varies with the type of
solvent and the like is typically in the range of -30°C ~
5 100°C , preferably 0°C ~ 80°C . The reaction time which
varies with the reaction temperature and the like is
typically in the range of 10 minutes - 48 hours, preferably
30 minutes - 24 hours.

Step A5 is for producing compound (6) and implemented
10 by reacting compound (5) with an acid in an aqueous solvent.

The inert solvent to be used is not limited in any
particular way as long as it does not interfere with the
reaction; examples are mixed solvents consisting of water
and ether solvents such as ether, tetrahydrofuran and
15 dioxane, alcoholic solvents such as methanol and ethanol,
or ketonic solvents such as acetone, with hydrous acetone
being preferred. The acid to be used may be exemplified by
inorganic acids such as hydrochloric acid, hydrobromic acid,
hydroiodic acid, sulfuric acid and phosphoric acid, and
20 organic acids such as acetic acid, p-toluenesulfonic acid,
pyridinium-p-toluenesulfonate, with hydrochloric acid being
preferred. The reaction temperature which varies with the
type of solvent and the like is typically in the range of
 0°C - 100°C (preferably 30°C - 80°C). The reaction time
25 which varies with the reaction temperature and the like is
typically in the range of 15 minutes - 24 hours (preferably
30 minutes - 10 hours).

Step A6 is for producing compound (7) and implemented

by performing catalytic reduction in an alcoholic solvent or an inert solvent.

The solvent to be used may be exemplified by alcoholic solvents such as methanol, ethanol, n-propanol, i-propanol, 5 n-butanol, s-butanol, t-butanol, pentanol, hexanol, cyclopropanol, cyclobutanol, cyclopentanol, cyclohexanol, ethylene glycol, 1,3-propanediol, 1,4-butanediol and 1,5-pentenediol, ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as 10 benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N- 15 methylpyrrolidone, ethyl acetate, acetonitrile and nitromethane; preferred examples are ethanol, dioxane, benzene and ethyl acetate.

The condition to be used in catalytic reduction is a homogeneous system such as hydrogen- 20 chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-chlorotris(triparamethoxyphenylphosphine)rhodium(I), hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I), hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II) 25 acetate, hydrogen-chlorohydridetris(triphenylphosphine) ruthenium(II), hydrogen-carboxylatohydridetris(triphenylphosphine)ruthenium(II), hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I),

hydrogen-platinum(II)-tin chloride complex, hydrogen-pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine cobalt(II) complex, hydrogen-bis(dimethylglyoximate)cobalt(II) complex, hydrogen-methyl benzoate-tricarbonylchromium complex, hydrogen-bis(tricarbonylcyclopentadienylchromium), hydrogen-pentacarbonyliron, hydrogen-bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-hydridecarbonylcobalt complex, hydrogen-octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium, hydrogen-chromium(III) acetylacetonato-triisobutylaluminum, hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an inhomogeneous system condition such as hydrogen-platinum dioxide, hydrogen-platinum/carbon, hydrogen-palladium/carbon, hydrogen-palladium/barium sulfate, hydrogen-palladium/calcium carbonate, hydrogen-Raney nickel, hydrogen-copper chromite, hydrogen-rhodium/carbon, hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide, or hydrogen-ruthenium/carbon; preferred examples are hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-palladium/carbon, hydrogen-palladium/calcium carbonate, etc.

The reaction temperature is typically in the range of 0°C - 100°C, preferably 0°C - 60°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 10 minutes - 6 hours.

Step A8 is for producing compound (9) in the case

where Q^2 in R^2 in compound (7) is -S- and implemented by reacting compound (7) with an oxidizing agent in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction and examples include halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, alcoholic solvents such as methanol and ethanol, ether solvents such as tetrahydrofuran, as well as water, and mixtures thereof; preferred examples are dichloromethane, methanol and a mixture of tetrahydrofuran and water.

The oxidizing agent to be used may be exemplified by organic peroxides such as t-butyl perbenzoate, t-butyl peracetate, t-butyl hydroperoxide, t-amyl hydroperoxide, dibenzoyl peroxide, di-p-nitrobenzoyl peroxide and di-p-chlorobenzoyl peroxide, organic peracids such as perbenzoic acid, metachloroperbenzoic acid, p-nitroperbenzoic acid, monoperoxyphthalic acid, performic acid, peracetic acid, trifluoroperacetic acid and peroxyauric acid, halogens such as hypochlorous acid, sodium hypochlorite, potassium hypobromite, potassium hypoiodite, sodium chlorate, potassium chlorate, sodium bromate, potassium bromate, sodium iodate, potassium iodate, perchloryl fluoride, orthoperiodic acid, sodium metaperiodate, potassium metaperiodate, N-bromoacetamide, N-bromosuccinimide, N-bromophthalimide, isocyanuric chloride, isocyanuric bromide,

N-bromocaprolactam, 1-chlorobenzotriazole, 1,3-dibromo-5,5-dimethylhydantoin, sodium N-chloro-p-toluenesulfonamide (chloramine T), sodium N-chlorobenzenesulfonamide (chloramine B), t-butyl hypochlorite, t-butyl hypobromite, 5 t-butyl hypoiodite, iodosylbenzene acetate and iodosylbenzene, as well as peroxomonosulfuric acid, OXONE (registered trademark) and hydrogen peroxide; preferred examples are sodium periodate and OXONE (registered trademark).

10 The reaction temperature which varies with the type of solvent and the like is typically in the range of -20°C ~ 30°C (preferably -10°C ~ 10°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 15 30 minutes - 15 hours).

Step A9 is for producing compound (10) in the case where Q^2 in R^2 in compound (7) is -S- and implemented by reacting compound (7) with an oxidizing agent in an inert solvent.

20 The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction and examples include halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, aromatic solvents such as benzene, toluene, 25 xylene, quinoline and chlorobenzene, alcoholic solvents such as methanol and ethanol, ether solvents such as tetrahydrofuran, as well as water, and mixtures thereof; preferred examples are dichloromethane, methanol and a

mixture of tetrahydrofuran and water.

The oxidizing agent to be used may be exemplified by organic peroxides such as t-butyl perbenzoate, t-butyl peracetate, t-butyl hydroperoxide, t-amyl hydroperoxide, 5 dibenzoyl peroxide, di-p-nitrobenzoyl peroxide and di-p-chlorobenzoyl peroxide, organic peracids such as perbenzoic acid, metachloroperbenzoic acid, p-nitroperbenzoic acid, monoperoxyphthalic acid, performic acid, peracetic acid, trifluoroperacetic acid and peroxyauric acid, halogens 10 such as hypochlorous acid, sodium hypochlorite, potassium hypobromite, potassium hypiodite, sodium chlorate, potassium chlorate, sodium bromate, potassium bromate, sodium iodate, potassium iodate, perchloryl fluoride, orthoperiodic acid, sodium metaperiodate, potassium 15 metaperiodate, N-bromoacetamide, N-bromosuccinimide, N-bromophthalimide, isocyanuric chloride, isocyanuric bromide, N-bromocaprolactam, 1-chlorobenzotriazole, 1,3-dibromo-5,5-dimethylhydantoin, sodium N-chloro-p-toluenesulfonamide (chloramine T), sodium N-chlorobenzenesulfonamide 20 (chloramine B), t-butyl hypochlorite, t-butyl hypobromite, t-butyl hypiodite, iodosylbenzene acetate and iodosylbenzene, as well as peroxomonosulfuric acid, OXONE (registered trademark) and hydrogen peroxide; a preferred example is OXONE (registered trademark).

25 The reaction temperature which varies with the type of solvent and the like is typically in the range of 0 °C ~ 100 °C (preferably 10°C ~ 50°C). The reaction time which varies with the reaction temperature and the like is

typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

Step A10 is for producing compound (145) and implemented by reacting compound (144) with a base in an inert solvent to make a salt of compound (144) and then reacting it with compound (134) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

A compound having the hydroxyl group in 11-position of compound (144) oriented in β configuration is commercially available and using this compound in place of compound (144), one can obtain a compound having X^1 in compound (7) oriented in β configuration.

Step A11 is for producing compound (146) and implemented by reacting compound (145) with compound (135) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the aforementioned step A4 in process A.

Step A12 is for producing compound (147) and implemented by reacting compound (146) with a reducing agent in an optionally miscible inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not interfere with the reaction; examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, alcoholic solvents such as methanol and ethanol, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, and amines such as pyridine and

triethylamine; preferred examples are alcoholic solvents such as methanol and ethanol, with methanol being more preferred. The reducing agent to be used may be exemplified by: metal hydrogen complex compounds such as

5 aluminum lithium hydride, trimethoxyaluminum lithium hydride, tri-t-butoxyaluminum lithium hydride, aluminum lithium hydride-trichloroaluminum (alane), aluminum lithium hydride-boron trifluoride, aluminum hydride magnesium chloride, magnesium aluminum hydride, sodium aluminum

10 hydride, sodium triethoxyaluminum hydride, sodium bis(methoxyethoxy)aluminum hydride, sodium boron hydride, sodium boron hydride-palladium/carbon, sodium boron hydrogensulfide, sodium boron hydrogencyanide, sodium trimethoxyboron hydride, lithium boron hydride, lithium

15 boron hydrogencyanide, lithium triethylboron hydride, lithium tri-s-butylboron hydride, lithium tri-t-butylboron hydride, calcium boron hydride, potassium boron hydride, potassium triisopropoxyboron hydride, potassium tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium

20 boron hydride, and tetra-n-butylammonium cyanoboron hydride; metal hydrides such as diisobutylaluminum hydride, triphenyltin hydride, tri-n-butyltin hydride, diphenyltin hydride, di-n-butyltin hydride, triethyltin hydride, trimethyltin hydride, trichlorosilane/tri-n-butylamine,

25 trichlorosilane/tri-n-propylamine, triethylsilane, trimethylsilane, diphenylsilane, phenylsilane, polymethylhydrosiloxane, dimethylphenylsilane, di-n-butylsilane, and methylphenylsilane; borane derivatives

such as diborane, dimethylamine-borane, trimethylamine-borane, ethylenediamine-borane, pyridine-borane, dimethylsulfide-borane, 2,3-dimethyl-2-butylborane (thexylborane), bis-3-methyl-2-butylborane (disiamylborane),
5 diisopinocamphephenylborane, dicyclohexylborane, and 9-borabicyclo[3,3,1]nonane (9-BBN); preferred examples are metal hydrogen complex compounds such as aluminum lithium hydride, trimethoxyaluminum lithium hydride, tri-t-butoxyaluminum lithium hydride, aluminum lithium hydride-
10 trichloroaluminum (alane), aluminum lithium hydride-boron trifluoride, aluminum hydride magnesium chloride, magnesium aluminum hydride, sodium aluminum hydride, sodium triethoxyaluminum hydride, sodium bis(methoxyethoxy)aluminum hydride, sodium boron hydride,
15 sodium boron hydride-palladium/carbon, sodium boron hydrosulfide, sodium boron hydrogencyanide, sodium trimethoxyboron hydride, lithium boron hydride, lithium boron hydrogencyanide, lithium triethylboron hydride, lithium tri-s-butylboron hydride, lithium tri-t-butylboron
20 hydride, calcium boron hydride, potassium boron hydride, potassium triisopropoxyboron hydride, potassium tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium boron hydride, and tetra-n-butylammonium cyanoboron hydride, with sodium boron hydride being more preferred. The
25 reaction temperature which varies with the type of solvent and the like is typically in the range of -30°C ~ 100°C , preferably 0°C ~ 70°C . The reaction time which varies with the reaction temperature and the like is typically in the


range of 10 minutes - 48 hours, preferably 30 minutes - 24 hours.

Step A13 is for producing compound (7) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the
5 aforementioned step A6 in process A.

Process B is for producing compound (17) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula
10 (II) in which Ar is a single bond, A is -O- and R^1 is -G-S-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double
15 bond; compound (18) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is -G-S(O)-Z-, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together
20 with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; and compound (19) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general
25 formula (II) in which Ar is a single bond, A is -O- and R^1 is -G-S(O)₂-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the

dashed line together with the solid line is a single bond or a double bond.

Process B



5 Step B1 is for producing compound (13) and implemented by reacting compound (3) with a base in an inert solvent to make a salt of compound (3) and then reacting it with compound (136) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

10 Step B2 is for producing compound (14) and implemented by reacting compound (13) with a deprotecting agent, namely by removing the substituted silyl group, in an inert solvent.

15 The inert solvent to be used is not limited in an particular way as long as it does not interfere with the reaction; examples include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, as well as

dimethylformamide and water, with tetrahydrofuran being preferred. The deprotecting agent to be used is not limited in any particular way and may be exemplified by fluorides such as hydrogen fluoride, hydrogen fluoride-pyridine, sodium fluoride, potassium fluoride and tetra-n-butylammonium fluoride, and organic acids such as formic acid, acetic acid and p-toluenesulfonic acid, with tetra-n-butylammonium fluoride being preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 80°C (preferably 0°C - 50°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

Step B3 is for producing compound (15) and implemented by reacting compound (14) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (14) with a halogenating agent in an inert solvent.

The amine-containing solvent to be used is not limited in any particularly way and may be exemplified by triethylamine, diisopropylethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene and pyridine, with pyridine and triethylamine being preferred.

The sulfonyl chloride compound to be used is not limited in any particular way and may be exemplified by p-toluenesulfonyl chloride, benzenesulfonyl chloride, methanesulfonyl chloride and trifluoromethanesulfonyl chloride, with methanesulfonyl chloride and

trifluoromethanesulfonyl chloride being preferred.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction of interest; examples include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, nitrile-containing solvents such as acetonitrile, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone and ethyl acetate, with benzene and dichloromethane being preferred.

The halogenating agent to be used may be exemplified by chlorinating agents such as carbon tetrachloride-triphenylphosphine, thionyl chloride, sulfuryl chloride, N-chlorosuccinimide-triphenylphosphine, N-chlorosuccinimide-dimethyl sulfide, phosphorus trichloride and phosphorus pentachloride, and brominating agents such as carbon tetrabromide-triphenylphosphine, N-bromosuccinimide-triphenylphosphine, N-bromosuccinimide-dimethyl sulfide, phosphorus tribromide and phosphorus pentabromide, and preferred examples are carbon tetrabromide-triphenylphosphine and thionyl chloride. The reaction temperature is typically in the range of 0°C - 80°C, preferably 10°C - 40°C. The reaction time which varies with the reaction temperature is typically in the range of 10 minutes - 10 hours, preferably 30 minutes - 3 hours.

Step B4 is for producing compound (16) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and then reacting it with compound (15) in an
5 alcoholic solvent.

The alcoholic solvent to be used is not limited in any particular way and may be exemplified by methanol, ethanol, n-propanol, i-propanol and mixed solvents containing them, and preferred examples are methanol and a methanol-
10 tetrahydrofuran mixed solvent.

The metal alkoxide to be used is not limited in any particular way and may be exemplified by sodium methoxide and sodium ethoxide, with sodium methoxide being preferred.

The reaction temperature which varies with the solvent
15 and other conditions is typically in the range of 0°C - 80°C, preferably 10°C - 40°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 10 hours, preferably 30 minutes - 8 hours.

20 Step B5 is for producing compound (17) and implemented by reacting compound (16) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.


Step B6 is for producing compound (18) and implemented
25 by reacting compound (17) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.

Step B7 is for producing compound (19) and implemented

by reacting compound (18) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

Process B' is an alternative method for producing compound (243) having the dashed line in compound (17) forming a double bond together with the solid line and compound (244) having the dashed line in compound (17) forming a single bond together with the solid line.

Process B'



10

Step B'1 is for producing compound (149) and implemented by reacting compound (148) with a base in an inert solvent to make a salt of compound (148) and then reacting it with compound (136) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

Step B'2 is for producing compound (150) and

implemented by reacting compound (149) with a deprotecting agent, namely by removing the substituted silyl group, in an inert solvent. The reaction is performed as in the aforementioned step B2 in process B.

5 Step B'3 is for producing compound (151) and implemented by reacting compound (150) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (150) with a halogenating agent in an inert solvent. The reaction is performed as in the
10 aforementioned step B3 in process B.

 Step B'4 is for producing compound (152) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and then reacting it with
15 compound (151) in an alcoholic solvent. The reaction is performed as in the aforementioned step B4 in process B.

 Step B'5 is for producing compound (153) and implemented by reacting compound (152) with an acid in an aqueous solvent. The reaction is performed as in the
20 aforementioned step A5 in process A.

 Step B'6 is for producing compound (243) and implemented by reacting compound (153) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step A12 in
25 process A.

 Step B'7 is for producing compound (244) and implemented by performing catalytic reduction of compound (243) in an alcoholic solvent or an inert solvent or

reducing compound (243) with a reducing agent in an optionally miscible inert solvent.

The solvent to be used in performing catalytic reduction

5 may be exemplified by alcoholic solvents such as methanol, ethanol, n-propanol, i-propanol, n-butanol, s-butanol, t-butanol, pentanol, hexanol, cyclopropanol, cyclobutanol, cyclopentanol, cyclohexanol, ethylene glycol, 1,3-propanediol, 1,4-butanediol and 1,5-pentanediol, ether
10 solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, amine-containing solvents such as pyridine
15 and triethylamine, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, ethyl acetate, acetonitrile and nitromethane; preferred examples are ethanol, ether, dioxane, and pyridine.

20 The condition to be used in catalytic reduction is a homogeneous system such as hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-chlorotris(triparamethoxyphenylphosphine)rhodium(I),
25 hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I), hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II) acetate, hydrogen-chlorohydridetris(triphenylphosphine)ruthenium(II),

- hydrogen-
- carboxylatohydridetris(triphenylphosphine)ruthenium(II),
- hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I),
- hydrogen-platinum(II)-tin chloride complex, hydrogen-
- 5 pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine
- cobalt(II) complex, hydrogen-
- bis(dimethylglyoximato)cobalt(II) complex, hydrogen-methyl
- benzoate-tricarbonylchromium complex, hydrogen-
- bis(tricarbonylcyclopentadienylchromium), hydrogen-
- 10 pentacarbonyliron, hydrogen-
- bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-
- hydridecarbonylcobalt complex, hydrogen-
- octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium,
- hydrogen-chromium(III) acetylacetonato-triisobutylaluminum,
- 15 hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or
- hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an
- inhomogeneous system condition such as hydrogen-platinum
- dioxide, hydrogen-platinum/carbon, hydrogen-
- palladium/carbon, hydrogen-palladium/barium sulfate,
- 20 hydrogen-palladium/calcium carbonate, hydrogen-Raney nickel,
- hydrogen-copper chromite, hydrogen-rhodium/carbon,
- hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide, or
- hydrogen-ruthenium/carbon; preferred examples are hydrogen-
- chlorotris(triphenylphosphine)rhodium(I), hydrogen-
- 25 palladium/carbon, hydrogen-palladium/calcium carbonate, etc.

The reaction temperature is typically in the range of 0°C - 100°C, preferably 0°C - 60°C. The reaction time which varies with the reaction temperature and the like is

typically in the range of 10 minutes - 24 hours, preferably 10 minutes - 6 hours.

The inert solvent to be used in the reaction with the reducing agent is not limited in any particular way as long as it does not participate in the reaction; examples include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, amine-containing solvents such as pyridine and triethylamine, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, acetonitrile and nitromethane; preferred examples are tetrahydrofuran, benzene, toluene and pyridine.

The reducing agent to be used may be exemplified by metals such as sodium/liquid ammonia, lithium/liquid ammonia, lithium/methylamine, lithium/ethylamine, lithium/ethylenediamine, sodium/hexamethylphosphamide-t-butanol, sodium/ethanol, sodium/t-butanol-tetrahydrofuran and sodium/toluene-t-amyl alcohol, metal hydrides such as triphenyltin hydride, tri-n-butyltin hydride, diphenyltin hydride, di-n-butyltin hydride, triethyltin hydride, trimethyltin hydride, trichlorosilane/tri-n-butylamine, trichlorosilane/tri-n-propylamine, triethylsilane, trimethylsilane, diphenylsilane, phenylsilane, polymethylhydrosiloxane, dimethylphenylsilane, di-n-butylsilane and methylphenylsilane, metal hydrogen complex


compounds such as lithium aluminum hydride/copper(I) iodide, trimethoxyaluminum lithium hydride/copper(I) bromide, tri-
 t-butoxyaluminum lithium hydride/copper(I) bromide, sodium
 boron hydride, sodium boron hydride-palladium/carbon,
 5 sodium boron hydrogensulfide, sodium boron hydrogencyanide,
 sodium trimethoxyboron hydride, lithium boron hydride,
 lithium boron hydrogencyanide, lithium triethylboron
 hydride, lithium tri-s-butylboron hydride, lithium tri-t-
 butylboron hydride, calcium boron hydride, potassium boron
 10 hydride, potassium triisopropoxyboron hydride, potassium
 tri-s-butylboron hydride, zinc boron hydride,
 tetramethylammonium boron hydride and tetra-n-butylammonium
 cyanoboron hydride; preferred examples are sodium/liquid
 ammonia, lithium/liquid ammonia, triphenyltin hydride, tri-
 15 n-butyltin hydride, lithium aluminum hydride/copper(I)
 iodide, trimethoxyaluminum lithium hydride/copper(I)
 bromide, sodium boron hydride and potassium tri-s-
 butylboron hydride.

The reaction temperature which varies with the type of
 20 reducing agent is typically in the range of -80°C ~ 100°C ,
 preferably -78°C ~ 80°C . The reaction time which varies
 with the reaction temperature and the like is typically in
 the range of 10 minutes - 24 hours, preferably 10 minutes -
 6 hours.

25 Process C is for producing compound (25) represented
 by the general formula (I) in which X^1 is a group of β
 configuration that is represented by the general formula
 (II) in which Ar is a single bond, A is -O- and R^1 is -G-

CONH-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

Process C



Step C1 is for producing compound (22) and implemented by reacting compound (15) with a cyanylating agent in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, methyl acetate, acetonitrile and

nitromethane; a preferred example is dimethyl sulfoxide.

The cyanlyating agent to be used may be exemplified by lithium cyanide, sodium cyanide, potassium cyanide, etc. and sodium cyanide is preferred. The reaction temperature
5 is typically in the range of 0°C - 80°C, preferably 10°C - 40°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 1 hour - 15 hours.

Step C2 is for producing compound (23) and implemented
10 by hydrolyzing compound (22) in the presence of a base.

The solvent to be used is not limited in any particular way as long as it is used in ordinary hydrolytic reaction; examples can be alcoholic solvents such as methanol and ethanol, ether solvents such as
15 tetrahydrofuran and dioxane, water, and mixtures thereof; preferred examples are water and hydrous alcoholic solvents such as water-ethanol.

The base to be used is not limited in any particular way as long as it does not affect other portions of the
20 compound; preferred examples are metal hydroxides such as lithium hydroxide, sodium hydroxide, potassium hydroxide, calcium hydroxide, barium hydroxide and cesium hydroxide, with sodium hydroxide and potassium hydroxide being particularly preferred.

25 The reaction temperature is typically in the range of 0°C - 100°C, preferably 50°C - 100°C.

The reaction time which varies with the reaction temperature and the like is typically in the range of 10

minutes - 48 hours, preferably 5 hours - 48 hours.

Step C3 is for producing compound (24) and implemented by reacting compound (23) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with
5 compound (138) or acid addition salts thereof in an inert solvent.

The reaction is performed by, for example, the acid halide method, the mixed acid anhydride method, the active ester method or the condensation method. The acid halide
10 method is implemented by reacting compound (23) with a halogenating agent (e.g. thionyl chloride, chloride oxalate, phosphorus pentachloride, etc.) in an inert solvent to prepare an acid halide which is then reacted with compound (138) or an acid addition salt in an inert solvent in the
15 presence or absence of a base (preferably in it's presence). The base to be used may be exemplified by organic amines such as triethylamine, N-methylmorpholine, pyridine and 4-dimethylaminopyridine, alkali metal bicarbonates such as sodium bicarbonate and potassium bicarbonate, and alkali
20 metal carbonates such as sodium carbonate and potassium carbonate; organic amines are preferred (with triethylamine being particularly preferred).

The solvent to be used is not limited in any particular way as long as it does not participate in the
25 reaction; examples include hydrocarbon solvents such as hexane, cyclohexane, benzene, toluene and xylene, halogen-containing solvents such as dichloromethane, 1,2-dichloroethane and carbon tetrachloride, ether solvents

such as ether, tetrahydrofuran and dioxane, ketonic solvents such as acetone, amide-containing solvents such as N,N-dimethylacetamide, N,N-dimethylformamide and N-methyl-2-pyrrolidone, and sulfoxide-containing solvents such as dimethyl sulfoxide; preferred examples are hydrocarbon solvents, halogen-containing solvents and ether solvents, and more preferred examples are ether solvents (with tetrahydrofuran being particularly preferred). The reaction temperature varies with the type of solvent and the like; however, for both the reaction of a halogenating agent with compound (23) and the reaction of an acid halide with compound (138) or its acid addition salt, the range is typically between -20 °C and 150 °C; preferably, the temperature for the reaction between a halogenating agent and compound (23) is in the range of -10°C ~ 50°C, and the temperature for the reaction between an acid halide and compound (138) or its acid addition salt is in the range of 0°C - 100°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably, 30 minutes - 15 hours).

The mixed acid anhydride method is implemented by reacting a C₁-C₆ alkyl halogenocarbonate (where C₁-C₆ refers to a straight-chained or branched alkyl group having 1 - 6 carbon atoms), a di-C₁-C₆ alkylcyanophosphoric acid or a diarylphosphorylazide with compound (23) to prepare a mixed acid anhydride which is then reacted with compound (138) or an acid addition salt thereof. The reaction for preparing

a mixed acid anhydride is performed by reacting a C_1-C_6 alkyl halogenocarbonate such as methyl chlorocarbonate, ethyl chlorocarbonate, isobutyl chlorocarbonate or hexyl chlorocarbonate (preferably ethyl chlorocarbonate or isobutyl chlorocarbonate), a di- C_1-C_6 alkylcyanophosphoric acid such as dimethylcyanophosphoric acid, diethylcyanophosphoric acid or dihexylcyanophosphoric acid or a diarylphosphoric acid azide such as diphenylphosphoric acid azide, di-(p-nitrophenyl)phosphoric acid azide or dinaphthylphosphoric acid azide (preferably diphenylphosphoric acid azide) with compound (23), preferably in an inert solvent in the presence of a base.

The base and the inert solvent to be used are the same as those used when the acid halide method is employed in the step under consideration. The reaction temperature which varies with the type of solvent and the like is typically in the range of $-20^{\circ}\text{C} \sim 50^{\circ}\text{C}$ (preferably $0^{\circ}\text{C} - 30^{\circ}\text{C}$). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

The reaction between a mixed acid anhydride and compound (138) or its acid addition salt is performed in an inert solvent in the presence or absence (preferably the presence) of a base, and the base and the inert solvent to be used are the same as those used in the above acid halide method. The reaction temperature which varies with the type of solvent and the like is typically in the range of $-20^{\circ}\text{C} \sim 50^{\circ}\text{C}$ (preferably $0^{\circ}\text{C} - 30^{\circ}\text{C}$). The reaction time

which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours). If a di-C₁-C₆ alkylcyanophosphoric acid or a diarylphosphoric acid azide is used in the process under consideration, compound (23) may be directly reacted with compound (138) or its acid addition salt in the presence of a base.

The active esterification method is implemented by reacting compound (23) with an active esterifying agent (e.g. N-hydroxy compounds such as N-hydroxysuccinimide and N-hydroxybenzotriazole) in the presence of a condensing agent (e.g. dicyclohexylcarbodiimide or carbonyldiimidazole) to prepare an active ester which is then reacted with compound (138) or an acid addition salt thereof. The reaction for preparing an active ester is preferably performed in an inert solvent and the inert solvent to be used may be exemplified by ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as dimethylformamide, ethyl acetate, acetonitrile, etc.; preferred examples are dichloromethane, acetonitrile, ethyl acetate, etc. The reaction temperature varies with the type of solvent and the like; the temperature for the active esterification reaction is typically in the range of -20°C ~ 50°C (preferably -10°C ~ 30°C), and the temperature for the reaction between the active ester compound and compound (138) or its acid addition salt is typically in


the range of -20°C ~ 50°C (preferably -10°C ~ 30°C). The reaction time varies with the reaction temperature and the like; however, for both reactions, it is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15
5 hours).

The condensation method is performed by directly reacting compound (23) with compound (138) or an acid addition salt thereof in the presence of a condensing agent [e.g. dicyclohexylcarbodiimide, carbonyldiimidazole, or 1-
10 (N,N-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride]. This reaction is performed as in the aforementioned reaction for preparing the active ester.

Step C4 is for producing compound (25) and implemented by reacting compound (24) with an acid in an aqueous
15 solvent and this reaction is performed as in the aforementioned step A5 in process A.

Process C' is a method of producing compound (245) having the dashed line in compound (25) forming a double bond together with the solid line, and compound (158)
20 having the dashed line in compound (25) forming a single bond together with the solid line.

Process C'



Step C'1 is for producing compound (154) and implemented by reacting compound (151) with a cyanylating agent in an inert solvent. The reaction is performed as in the aforementioned step C1 in process C.

Step C'2 is for producing compound (155) and implemented by hydrolyzing compound (154) in the presence of a base. The reaction is performed as in the aforementioned step C2 in process C.

Step c'3 is for producing compound (156) and implemented by reacting compound (155) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (138) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

Step C'4 is for producing compound (157) and implemented by reacting compound (156) with an acid in an

aqueous solvent. The reaction is performed as in the
aforementioned step A5 in process A.

Step C'5 is for producing compound (245) and
implemented by reacting compound (157) with a reducing
5 agent in an optionally miscible inert solvent. The
reaction is performed as in the aforementioned step A12 in
process A.

Step C'6 is for producing compound (158) and
implemented by performing catalytic reduction of compound
10 (245) in an alcoholic solvent or an inert solvent or
reacting compound (245) with a reducing agent in an
optionally miscible inert solvent.

The solvent to be used in performing catalytic
reduction may be exemplified by alcoholic solvents such as
15 methanol, ethanol, n-propanol, i-propanol, n-butanol, s-
butanol, t-butanol, pentanol, hexanol, cyclopropanol,
cyclobutanol, cyclopentanol, cyclohexanol, ethylene glycol,
1,3-propanediol, 1,4-butanediol and 1,5-pentanediol, ether
solvents such as ether, tetrahydrofuran, dioxane and
20 dimethoxyethane, aromatic solvents such as benzene, toluene,
xylene, quinoline and chlorobenzene, halogen-containing
solvents such as dichloromethane, chloroform and carbon
tetrachloride, amine-containing solvents such as pyridine
and triethylamine, as well as cyclohexane, dimethyl
25 sulfoxide, dimethylacetamide, dimethylimidazolidinone,
dimethylformamide, N-methylpyrrolidone, ethyl acetate,
acetonitrile and nitromethane; preferred examples are
ethanol, ether, dioxane, and pyridine.

The condition to be used in catalytic reduction is a homogeneous system such as hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-chlorotris(triparamethoxyphenylphosphine)rhodium(I), hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I), hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II) acetate, hydrogen-chlorohydridetris(triphenylphosphine)ruthenium(II), hydrogen-carboxylatohydridetris(triphenylphosphine)ruthenium(II), hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I), hydrogen-platinum(II)-tin chloride complex, hydrogen-pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine cobalt(II) complex, hydrogen-bis(dimethylglyoximato)cobalt(II) complex, hydrogen-methyl benzoate-tricarbonylchromium complex, hydrogen-bis(tricarbonylcyclopentadienylchromium), hydrogen-pentacarbonyliron, hydrogen-bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-hydridecarbonylcobalt complex, hydrogen-octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium, hydrogen-chromium(III) acetylacetonato-triisobutylaluminum, hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an inhomogeneous system condition such as hydrogen-platinum dioxide, hydrogen-platinum/carbon, hydrogen-palladium/carbon, hydrogen-palladium/barium sulfate,

hydrogen-palladium/calcium carbonate, hydrogen-Raney nickel, hydrogen-copper chromite, hydrogen-rhodium/carbon, hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide, or hydrogen-ruthenium/carbon; preferred examples are hydrogen-
5 chlorotris(triphenylphosphine)rhodium(I), hydrogen-palladium/carbon, hydrogen-palladium/calcium carbonate, etc.

The reaction temperature is typically in the range of 0°C - 100°C, preferably 0°C - 60°C. The reaction time which varies with the reaction temperature and the like is
10 typically in the range of 10 minutes - 24 hours, preferably 10 minutes - 6 hours.

The inert solvent to be used in the reaction with the reducing agent is not limited in any particular way as long as it does not participate in the reaction; examples
15 include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, amine-containing
20 solvents such as pyridine and triethylamine, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, acetonitrile and nitromethane; preferred examples are tetrahydrofuran, benzene, toluene and pyridine.

25 The reducing agent to be used may be exemplified by metals such as sodium/liquid ammonia, lithium/liquid ammonia, lithium/methylamine, lithium/ethylamine, lithium/ethylenediamine, sodium/hexamethylphosphamide-t-


butanol, sodium/ethanol, sodium/t-butanol-terahydrofuran, sodium/toluene-t-amyl alcohol, metal hydrides such as triphenyltin hydride, tri-n-butyltin hydride, diphenyltin hydride, di-n-butyltin hydride, triethyltin hydride, 5 trimethyltin hydride, trichlorosilane/tri-n-butylamine, trichlorosilane/tri-n-propylamine, triethylsilane, trimethylsilane, diphenylsilane, phenylsilane, polymethylhydrosiloxane, dimethylphenylsilane, di-n-butylsilane and methylphenylsilane; metal hydrogen complex 10 compounds such as lithium aluminum hydride/copper(I) iodide, trimethoxyaluminum lithium hydride/copper(I) bromide, tri-t-butoxyaluminum lithium hydride/copper(I) bromide, sodium boron hydride, sodium boron hydride-palladium/carbon, sodium boron hydrogensulfide, sodium boron hydrogencyanide, 15 sodium trimethoxyboron hydride, lithium boron hydride, lithium boron hydrogencyanide, lithium triethylboron hydride, lithium tri-s-butylboron hydride, lithium tri-t-butylboron hydride, calcium boron hydride, potassium boron hydride, potassium triisopropoxyboron hydride, potassium 20 tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium boron hydride and tetra-n-butylammonium cyanoboron hydride; preferred examples are sodium/liquid ammonia, lithium/liquid ammonia, triphenyltin hydride, tri-n-butyltin hydride, lithium aluminum hydride/copper(I) 25 iodide, trimethoxyaluminum lithium hydride/copper(I) bromide, sodium boron hydride and potassium tri-s-butylboron hydride.

The reaction temperature which varies with the type of

reducing agent is typically in the range of $-80^{\circ}\text{C} \sim 100^{\circ}\text{C}$, preferably $-78^{\circ}\text{C} \sim 80^{\circ}\text{C}$. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 10 minutes - 6 hours.

Process D is for producing compound (30) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is $-\text{O}-$ and R^1 is $-\text{G}-\text{NHCO}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond.

15 Process D



Step D1 is for producing compound (27) and implemented

by reacting compound (14) with phthalimide in an inert solvent in the presence of an azodicarboxylic acid dialkyl ester (preferably diethyl azodicarboxylate) and a phosphine compound (preferably triphenylphosphine).

5 The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and aromatic solvents such as benzene, toluene, xylene, quinoline and
10 chlorobenzene, with tetrahydrofuran being preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 50°C (preferably 10°C - 30°C). The reaction time which varies with the reaction temperature and the like is typically in
15 the range of 15 minutes - 48 hours (preferably 30 minutes - 24 hours).

Step D2 is an alternative step for producing compound (27) and implemented by reacting compound (15) with a metal salt of phthalimide (preferably phthalimide potassium) in
20 an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic
25 solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, as well as dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone,

dimethylformamide, and N-methylpyrrolidone; a preferred example is tetrahydrofuran.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 50°C (preferably 10°C - 30°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 48 hours (preferably 30 minutes - 24 hours).

Step D3 is for producing compound (28) and implemented by reacting compound (27) with an amine-containing compound (preferably hydrazine) in an alcoholic solvent.

The alcoholic solvent to be used is not limited in any particular way as long as it does not interfere with the reaction and examples are methanol, ethanol, n-propyl alcohol and i-propyl alcohol, with ethanol being preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 50°C (preferably 10°C - 30°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 48 hours (preferably 30 minutes - 24 hours).


Step D4 is for producing compound (29) and implemented by reacting compound (139) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (28) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

Step D5 is for producing compound (30) and implemented

by reacting compound (29) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

Process D' is a method of producing compound (246) having the dashed line in compound (30) forming a double bond together with the solid line, and compound (163) having the dashed line in compound (30) forming a single bond together with the solid line.

Process D'



10

Step D'1 is for producing compound (159) and implemented by reacting compound (150) with phthalimide in an inert solvent in the presence of an azodicarboxylic acid dialkyl ester (preferably diethyl azodicarboxylate) and a

phosphine compound (preferably triphenylphosphine). The reaction is performed as in the aforementioned step D1 in process D.

Step D'2 is an alternative step for producing compound (159) and implemented by reacting compound (151) with a metal salt of phthalimide (preferably phthalimide potassium) in an inert solvent. The reaction is performed as in the aforementioned step D2 in process D.

Step D'3 is for producing compound (160) and implemented by reacting compound (159) with an amine-containing compound (preferably hydrazine) in an alcoholic solvent. The reaction is performed as in the aforementioned step D3 in process D.

Step D'4 is for producing compound (161) and implemented by reacting compound (139) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (160) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step D4 in process D.

Step D'5 is for producing compound (162) and implemented by reacting compound (161) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step C4 in process C.


Step D'6 is for producing compound (246) and implemented by reacting compound (162) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step A12 in process A.

Step D'7 is for producing compound (163) and implemented by performing catalytic reduction of compound (246) in an alcoholic solvent or an inert solvent or by reacting compound (246) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step C'6 in process C'.

Process E is for producing compound (35) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}=\text{CH}-\text{CH}_2-\text{R}^4$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond; compound (36) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(\text{CH}_2)_3-\text{R}^4$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond or a double bond; compound (38) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(\text{CH}_2)_3-\text{G}^3-\text{S}(\text{O})-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position

to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (39) represented by the general formula (I) in which X^1 is a group of β configuration that is
 5 represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(CH_2)_3-G^3-S(O)_2-Z$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line
 10 together with the solid line is a single bond or a double bond.

Process E



Step E1 is for producing compound (32) and implemented by reacting compound (134) with a metal (preferably magnesium) or an alkylolithium (preferably t-butyllithium) in an inert solvent to make a reactive derivative of compound (134) and reacting it with compound (2) in an inert solvent. The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and tetrahydrofuran is more preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 80°C (preferably 10°C - 50°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

The reaction of interest can also be implemented by reacting compound (2) with compound (140) in an inert solvent in the presence of an activator.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and halogen-containing solvents such as dichloromethane; more preferred examples are tetrahydrofuran and dichloromethane.

The activator to be used is not limited in any particular way as long as it does not interfere with the reaction; examples are fluorides such as tetra-n-

butylammonium fluoride, and Lewis acids such as aluminum trichloride, ethylaluminum dichloride, titanium tetrachloride, boron trifluoride, and trimethylsilyl trifluoromethanesulfonate; a preferred example is

5 trimethylsilyl trifluoromethanesulfonate.

The reaction temperature which varies with the type of solvent and the like is typically in the range of -78°C ~ 50°C (preferably -60°C ~ 30°C). The reaction time which varies with the reaction temperature and the like is

10 typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 5 hours).

Step E2 is for producing compound (33) and implemented by reacting compound (32) with a reducing agent in an inert solvent in the presence of an additive.

15 The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples include aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane,

20 chloroform and carbon tetrachloride, and ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane; preferred examples are benzene, toluene and dichloromethane.

The additive to be used is not limited in any

25 particular way as long as it does not interfere with the reaction and preferred examples are 1,1'-thiocarbonyl diimidazole, phosgene, benzoyl chloride, zinc iodide, boron trifluoride (BF_3), etc.

The reducing agent to be used may be exemplified by metal hydrides such as triphenyltin hydride, tri-n-butyltin hydride, diphenyltin hydride, di-n-butyltin hydride, triethyltin hydride, trimethyltin hydride,

5 trichlorosilane/tri-n-butylamine, trichlorosilane/tri-n-propylamine, triethylsilane, trimethylsilane, diphenylsilane, phenylsilane, polymethylhydrosiloxane, dimethylphenylsilane, di-n-butylsilane and methylphenylsilane; and metal hydrogen complex compounds

10 such as lithium aluminum hydride, trimethoxyaluminum lithium hydride, tri-t-butoxyaluminum lithium hydride, lithium aluminum hydride-trichloroaluminum (alane), lithium aluminum hydride-boron trifluoride, aluminum hydride magnesium chloride, magnesium aluminum hydride, sodium

15 aluminum hydride, sodium triethoxyaluminum hydride, sodium bis(methoxyethoxy)aluminum hydride, sodium boron hydride, sodium boron hydride-palladium/carbon, sodium boron hydrogensulfide, sodium boron hydrogencyanide, sodium trimethoxyboron hydride, lithium boron hydride, lithium

20 boron hydrogencyanide, lithium triethylboron hydride, lithium tri-s-butylboron hydride, lithium tri-t-butylboron hydride, calcium boron hydride, potassium boron hydride, potassium triisopropoxyboron hydride, potassium tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium

25 boron hydride, and tetra-n-butylammonium cyanoboron hydride; preferred examples are tri-n-butyltin hydride, triethylsilane and sodium boron hydrogencyanide.

The reaction temperature which varies with the type of

solvent and the like is typically in the range of 0°C ~ 150°C (preferably 10°C ~ 100°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

In this step, a compound having the allyl group in the 11-position of compound (33) oriented in α configuration forms as a by-product and this may be used to give compounds having X^1 in compound (35), compound (36), compound (38) and compound (39) oriented in α configuration.

Step E3 is for producing compound (34) and implemented by reacting compound (33) with compound (141) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the aforementioned step A4 in process A.

Step E4 is for producing compound (35) and implemented by reacting compound (34) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

Step E5 is for producing compound (36) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Step E7 is for producing compound (38) in the case where Q^2 in R^4 in compound (36) is -S- and implemented by reacting compound (36) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.


Step E8 is for producing compound (39) in the case where Q^2 in R^4 in compound (36) is -S- and implemented by reacting compound (36) with an oxidizing agent in an inert solvent. The reaction is performed as in the

5 aforementioned step A9 in process A.

Process E' is an alternative method of producing compound (247) having the dashed line in compound (35) forming a double bond together with the solid line, and compound (248) having the dashed line in compound (36)

10 forming a single bond together with the solid line.

Process E'



Step E'1 is for producing compound (165) and implemented by reacting compound (134) with a metal

15 (preferably magnesium) or an alkylolithium (preferably t-butylolithium) in an inert solvent to make a reactive derivative of compound (134) and reacting it with compound

(164) in an inert solvent. Alternatively, step E'1 may be implemented by reacting compound (164) with compound (140) in an inert solvent in the presence of an activator. The reaction is performed as in the aforementioned step E1 in
5 process E.

Step E'2 is for producing compound (166) and implemented by reacting compound (166) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step E2
10 in process E.

In this step, a compound having the allyl group in the 11-position of compound (166) oriented in α configuration forms as a by-product and this may be used to give compounds having X^1 in compound (35) and compound (36)
15 oriented in α configuration.

Step E'3 is for producing compound (167) and implemented by reacting compound (166) with compound (141) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the
20 aforementioned step E3 in process E.

Step E'4 is for producing compound (168) and implemented by reacting compound (167) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step D'5 in process D'.


25 Step E'5 is for producing compound (247) and implemented by reacting compound (168) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step A12 in

process A.

Step E'6 is for producing compound (36) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Process F is for producing compound (49) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{G}^4-\text{S}(\text{O})-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond, and compound (52) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{G}^4-\text{S}(\text{O})_2-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond.

Process F



Step F1 is for producing compound (42) and implemented by reacting compound (142) with an alkyllithium (preferably n-butyllithium) in an inert solvent to make a reactive derivative of compound (142) and reacting it with compound (2) in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and tetrahydrofuran is more preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0 °C - 80

°C (preferably 10 °C - 50 °C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

5 Step F2 is for producing compound (43) and implemented by reacting compound (42) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step E2 in process E.

10 In this step, a compound having the substituent in the 11-position of compound (43) oriented in α configuration forms as a by-product and this may be used to give compounds having X^1 in compound (49) and compound (52) oriented in α configuration.

15 Step F3 is for producing compound (44) and implemented by reacting compound (43) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the aforementioned step B2 in process B.

20 Step F4 is for producing compound (45) and implemented by reacting compound (44) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (44) with a halogenating agent in an inert solvent. The reaction is performed as in the aforementioned step B3 in process B.

25 Step F5 is for producing compound (46) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and reacting it with compound (45) in an alcoholic

solvent. The reaction is performed as in the
aforementioned step B4 in process B.

Step F6 is for producing compound (47) and implemented
by reacting compound (46) with an oxidizing agent in an
5 inert solvent. The reaction is performed as in the
aforementioned step A8 in process A.

Step F7 is for producing compound (48) and implemented
by reacting compound (47) with an acid in an aqueous
solvent. The reaction is performed as in the
10 aforementioned step A5 in process A.

Step F8 is for producing compound (49) and implemented
by performing catalytic reduction of compound (48) in an
alcoholic solvent or an inert solvent.

The solvent to be used may be exemplified by alcoholic
15 solvents such as methanol, ethanol, n-propanol, i-propanol,
n-butanol, s-butanol, t-butanol, pentanol, hexanol,
cyclopropanol, cyclobutanol, cyclopentanol, cyclohexanol,
ethylene glycol, 1,3-propanediol, 1,4-butanediol, and 1,5-
pentanediol, ether solvents such as ether, tetrahydrofuran,
20 dioxane and dimethoxyethane, aromatic solvents such as
benzene, toluene, xylene, quinoline and chlorobenzene,
halogen-containing solvents such as dichloromethane,
chloroform and carbon tetrachloride, as well as cyclohexane,
dimethyl sulfoxide, dimethylacetamide,
25 dimethylimidazolidinone, dimethylformamide, N-
methylpyrrolidone, ethyl acetate, acetonitrile and
nitromethane; a preferred example is ethyl acetate.

The condition to be used in catalytic reduction is a

- homogeneous system such as hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-chlorotris(triparamethoxyphenylphosphine)rhodium(I),
- 5 hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I), hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II) acetate, hydrogen-chlorohydridetris(triphenylphosphine)ruthenium(II), hydrogen-
- 10 carboxylatohydridetris(triphenylphosphine)ruthenium(II), hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I), hydrogen-platinum(II)-tin chloride complex, hydrogen-pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine cobalt(II) complex, hydrogen-
- 15 bis(dimethylglyoximato)cobalt(II) complex, hydrogen-methyl benzoate-tricarbonylchromium complex, hydrogen-bis(tricarbonylcyclopentadienylchromium), hydrogen-pentacarbonyliron, hydrogen-bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-
- 20 hydridecarbonylcobalt complex, hydrogen-octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium, hydrogen-chromium(III) acetylacetonato-triisobutylaluminum, hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an
- 25 inhomogeneous system condition such as hydrogen-platinum dioxide, hydrogen-platinum/carbon, hydrogen-palladium/carbon, hydrogen-palladium/barium sulfate, hydrogen-palladium/calcium carbonate, hydrogen-Raney nickel,

hydrogen-copper chromite, hydrogen-rhodium/carbon, hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide, or hydrogen-ruthenium/carbon; a preferred example is hydrogen palladium/carbon.


5 The reaction temperature is typically in the range of 0°C - 100°C, preferably 0°C - 60°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 10 minutes - 6 hours.

10 As an ancillary to this reaction, conversion to a single bond may occasionally be effected if the dashed line forms a double bond together with the solid line.

Step F11 is for producing compound (52) and implemented by reacting compound (49) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step B7 in process B.

Process F' is an alternative method of producing compound (249) having the dashed line in compound (48) forming a double bond together with the solid line.

20 Process F'



Step F'1 is for producing compound (169) and implemented by reacting compound (142) with an alkylolithium (preferably n-butyllithium) in an inert solvent to make a reactive derivative of compound (142) and reacting it with compound (164) in an inert solvent. The reaction is performed as in the aforementioned step F1 in process F.

Step F'2 is for producing compound (170) and implemented by reacting compound (169) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step F2 in process F.

In this step, a compound having the substituent in the 11-position of compound (170) oriented in α configuration forms as a by-product and this may be used to give a compound having X^1 in compound (249) oriented in α

configuration.

Step F'3 is for producing compound (171) and implemented by reacting compound (170) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the
5 aforementioned step F3 in process F.

Step F'4 is for producing compound (172) and implemented by reacting compound (171) with a sulfonyl chloride compound in an amine-containing solvent or
10 reacting compound (171) with a halogenating agent in an inert solvent. The reaction is performed as in the aforementioned step F4 in process F.

Step F'5 is for producing compound (173) and implemented by reacting compound (137) with a metal
15 alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and reacting it with compound (172) in an alcoholic solvent. The reaction is performed as in the aforementioned step F5 in process F.

Step F'6 is for producing compound (174) and
20 implemented by reacting compound (173) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step F6 in process F.


Step F'7 is for producing compound (175) and implemented by reacting compound (174) with an acid in an
25 aqueous solvent. The reaction is performed as in the aforementioned step E'4 in process E.

Step F'8 is for producing compound (249) and implemented by reacting compound (175) with a reducing

agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step E'5 in process E.

Process G is for producing compound (56) represented
5 by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{G}^4-\text{COOH}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the
10 carbon atom in 3-position to which they are bound, are - $(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond, and compound (57) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general
15 formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{G}^4-\text{CONH}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are - $(\text{C}=\text{O})-$, and the dashed line together with the solid line is
20 a single bond or a double bond.

Process G



Step G1 is for producing compound (53) and implemented by reacting compound (45) with a cyanylating agent in an inert solvent. The reaction is performed as in the
5
aforementioned step C1 in process C.

Step G2 is for producing compound (54) and implemented by hydrolyzing compound (53) in the presence of a base. The reaction is performed as in the aforementioned step C2
10
in process C.

Step G3 is for producing compound (55) and implemented by reacting compound (54) with an acid in an aqueous solvent. The reaction is performed as in the
aforementioned step A5 in process A.


Step G4 is for producing compound (56) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the
15
aforementioned step F8 in process F.

As an ancillary to this reaction, conversion to a single bond may occasionally be effected if the dashed line
20
forms a double bond together with the solid line.

Step G5 is for producing compound (57) and implemented by reacting compound (56) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (138) or acid addition salts thereof in an inert solvent. The reaction is performed as in the

Process G' is an alternative method of producing compound (250) having the dashed line in compound (55) forming a double bond together with the solid line.

10 Process G'



Step G'1 is for producing compound (176) and implemented by reacting compound (172) with a cyanylating agent in an inert solvent. The reaction is performed as in the

Step G'2 is for producing compound (177) and implemented by hydrolyzing compound (176) in the presence of a base. The reaction is performed as in the


20 Step G'3 is for producing compound (178) and

implemented by reacting compound (177) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step F'7 in process F'.

Step G'4 is for producing compound (250) and
5 implemented by reacting compound (178) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step F'8 in process F'.

Process H is for producing compound (63) represented by the general formula (I) in which X^1 is a group of β
10 configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{G}^4-\text{NHCO}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are
15 $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond.

Process H



Step H1 is for producing compound (59) and implemented by reacting compound (44) with phthalimide in an inert solvent in the presence of an azodicarboxylic acid dialkyl ester (preferably diethyl azodicarboxylate) and a phosphine compound (preferably triphenylphosphine). The reaction is performed as in the aforementioned step D1 in process D.

Step H2 is an alternative step for producing compound (59) and implemented by reacting compound (45) with a metal salt of phthalimide (preferably phthalimide potassium) in an inert solvent. The reaction is performed as in the aforementioned step D2 in process D.

Step H3 is for producing compound (60) and implemented by reacting compound (59) with an amine-containing compound

(preferably hydrazine) in an alcoholic solvent. The reaction is performed as in the aforementioned step D3 in process D.

Step H4 is for producing compound (61) and implemented
5 by reacting compound (139) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (60) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

10 Step H5 is for producing compound (62) and implemented by reacting compound (61) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.


Step H6 is for producing compound (63) and implemented
15 by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step F8 in process F.

As an ancillary to this reaction, conversion to a single bond may occasionally be effected if the dashed line
20 forms a double bond together with the solid line.

Process H' is an alternative method of producing compound (251) having the dashed line in compound (62) forming a double bond together with the solid line.

Process H'

25



Step H'1 is for producing compound (179) and implemented by reacting compound (171) with phthalimide in an inert solvent in the presence of an azodicarboxylic acid dialkyl ester (preferably diethyl azodicarboxylate) and a phosphine compound (preferably triphenylphosphine). The reaction is performed as in the aforementioned step H1 in process H.

Step H'2 is an alternative step for producing compound (179) and implemented by reacting compound (172) with a metal salt of phthalimide (preferably phthalimide potassium) in an inert solvent. The reaction is performed as in the aforementioned step H2 in process H.

Step H'3 is for producing compound (180) and

implemented by reacting compound (179) with an amine-containing compound (preferably hydrazine) in an alcoholic solvent. The reaction is performed as in the aforementioned step H3 in process H.

5 Step H'4 is for producing compound (181) and implemented by reacting compound (139) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (180) or acid addition salts thereof in an inert solvent. The reaction is performed as
10 in the aforementioned step H4 in process H.


Step H'5 is for producing compound (182) and implemented by reacting compound (181) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step G'3 in process G'.

15 Step H'6 is for producing compound (251) and implemented by reacting compound (182) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step G'4 in process G'.

20 Process I is for producing compound (70) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -CH₂-
25 CH=CH-CH₂-R², X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond

or a double bond; compound (71) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-(CH_2)_4-R^2$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond or a double bond; compound (73) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-(CH_2)_4-G^2-S(O)-Z$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (74) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-(CH_2)_4-G^2-S(O)_2-Z$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

Process I



Step I1 is for producing compound (65) and implemented by reacting compound (143) with a metal (preferably magnesium) or an alkyllithium (preferably n-butyllithium) in an inert solvent to make a reactive derivative of compound (143) and reacting it with compound (2) in an inert solvent. The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, and tetrahydrofuran is more preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 80°C (preferably

10°C - 50°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

Step I2 is for producing compound (66) and implemented
5 by reacting compound (65) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step E2 in process E.

In this step, a compound having the $-C_6H_4-OR^3$ in the 11-position of compound (66) oriented in α configuration
10 forms as a by-product and this may be used to give compounds having X^1 in compound (70), compound (71), compound (73) and compound (74) oriented in α configuration.

For the synthesis of compound (66) and a compound having the $-C_6H_4-OR^3$ in the 11-position of compound (66)
15 oriented in α configuration, reference may be had to the methods of introducing a variety of aromatic hydrocarbon groups as disclosed in Tetrahedron, vol. 52, 1529-1542, 1996.

Step I3 is for producing compound (67) and implemented
20 by reacting compound (66) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent.

The inert solvent to be used is not limited in an particular way as long as it does not interfere with the reaction; examples include ether solvents such as ether,
25 tetrahydrofuran, dioxane and dimethoxyethane, as well as dimethylformamide and water, with tetrahydrofuran being preferred. The deprotecting agent to be used is not limited in any particular way and may be exemplified by

fluorides such as hydrogen fluoride, hydrogen fluoride-pyridine, sodium fluoride, potassium fluoride and tetra-n-butylammonium fluoride, and organic acids such as formic acid, acetic acid and p-toluenesulfonic acid, with tetra-n-
5 butylammonium fluoride being preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 80°C (preferably 0°C - 50°C). The reaction time which varies with the reaction temperature and the like is typically in
10 the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

Step I4 is for producing compound (68) and implemented by reacting compound (67) with a base in an inert solvent to make a salt of compound (67) and then reacting it with
15 compound (134) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

Step I5 is for producing compound (69) and implemented by reacting compound (68) with compound (135) in an inert solvent in the presence of an organometallic catalyst. The
20 reaction is performed as in the aforementioned step A4 in process A.

Step I6 is for producing compound (70) and implemented by reacting compound (69) with an acid in an aqueous solvent. The reaction is performed as in the
25 aforementioned step A5 in process A.

Step I7 is for producing compound (71) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the


aforementioned step A6 in process A.

Step I9 is for producing compound (73) in the case where Q^2 in R^2 in compound (71) is -S- and implemented by reacting compound (71) with an oxidizing agent in an inert solvent. The reaction is performed as in step A8 in process A.

Step I10 is for producing compound (74) in the case where Q^2 in R^2 in compound (74) is -S- and implemented by reacting compound (71) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

Process I' is an alternative method of producing compound (252) having the dashed line in compound (70) forming a double bond together with the dashed line.

15 Process I'



Step I'1 is for producing compound (184) and

implemented by reacting compound (143) with a metal (preferably magnesium) or an alkylolithium (preferably n-butyllithium) in an inert solvent to make a reactive derivative of compound (143) and reacting it with compound
5 (164) in an inert solvent. The reaction is performed as in the aforementioned step I1 in process I.

Step I'2 is for producing compound (185) and implemented by reacting compound (184) with a reducing agent in an inert solvent in the presence of an additive.
10 The reaction is performed as in the aforementioned step I2 in process I.

In this step, a compound having the $-C_6H_4-OR^3$ in the 11-position of compound (185) oriented in α configuration forms as a by-product and this may be used to give a
15 compound having X^1 in compound (252) oriented in α configuration.

For the synthesis of compound (185) and a compound having the $-C_6H_4-OR^3$ in the 11-position of compound (185) oriented in α configuration, reference may be had to the
20 methods of introducing a variety of aromatic hydrocarbon groups as disclosed in Tetrahedron, vol. 52, 1529-1542, 1996.

Step I'3 is for producing compound (186) and implemented by reacting compound (185) with a deprotecting
25 agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the aforementioned step I3 in process I.

Step I'4 is for producing compound (187) and

implemented by reacting compound (186) with a base in an inert solvent to make a salt of compound (186) and then reacting it with compound (134) in an inert solvent. The reaction is performed as in the aforementioned step I4 in
5 process I.

Step I'5 is for producing compound (188) and implemented by reacting compound (187) with compound (135) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the
10 aforementioned step I5 in process I.

Step I'6 is for producing compound (189) and implemented by reacting compound (188) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step H'5 in process H'.


15 Step I'7 is for producing compound (252) and implemented by reacting compound (189) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step H'6 in process H'.

20 Process J is for producing compound (81) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -G-S-Z,
25 X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond;

compound (82) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -
5 O- and R^1 is -G-S(O)-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are - (C=O), and the dashed line together with the solid line is a single bond or a double bond; and compound (83)
10 represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -G²-S(O)₂-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and
15 R^c , when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond.

Process J

20



Step J1 is for producing compound (77) and implemented by reacting compound (67) with a base in an inert solvent to make a salt of compound (67) and reacting it with compound (136) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

Step J2 is for producing compound (78) and implemented by reacting compound (77) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the aforementioned step B2 in process B.

Step J3 is for producing compound (79) and implemented by reacting compound (78) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (78) with a halogenating agent in an inert solvent. The reaction is performed as in the aforementioned step B3 in

process B.

Step J4 is for producing compound (80) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and reacting it with compound (79) in an alcoholic solvent. The reaction is performed as in the aforementioned step B4 in process B.


Step J5 is for producing compound (81) and implemented by reacting compound (80) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

Step J6 is for producing compound (82) and implemented by reacting compound (81) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.

Step J7 is for producing compound (83) and implemented by reacting compound (82) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step B7 in process B.

Process J' is an alternative method for producing compound (253) having the dashed line in compound (81) forming a double bond together with the solid line, and compound (254) having the dashed line in compound (81) forming a single bond together with the solid line.

Process J'



Step J'1 is for producing compound (190) and implemented by reacting compound (186) with a base in an inert solvent to make a salt of compound (186) and reacting it with compound (136) in an inert solvent. The reaction is performed as in the aforementioned step J1 in process J.

Step J'2 is for producing compound (191) and implemented by reacting compound (190) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the aforementioned step J2 in process J.

Step J'3 is for producing compound (192) and implemented by reacting compound (191) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (191) with a halogenating agent in an inert solvent. The reaction is performed as in the

aforementioned step J3 in process J.

Step J'4 is for producing compound (193) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive
 5 derivative of compound (137) and reacting it with compound (192) in an alcoholic solvent. The reaction is performed as in the aforementioned step J4 in process J.

Step J'5 is for producing compound (194) and implemented by reacting compound (193) with an acid in an
 10 aqueous solvent. The reaction is performed as in the aforementioned step I'6 in process I'.


Step J'6 is for producing compound (253) and implemented by reacting compound (194) with a reducing agent in an optionally miscible inert solvent. The
 15 reaction is performed as in the aforementioned step I'7 in process I'.

Step J'7 is for producing compound (254) and implemented by performing catalytic reduction of compound (253) in an alcoholic solvent or an inert solvent or
 20 reacting compound (253) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step C'6 in process C'.

Process K is for producing compound (89) represented by the general formula (I) in which X^1 is a group of β
 25 configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -G-CONH-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and

R^o , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

5 Process K



Step K1 is for producing compound (86) and implemented by reacting compound (79) with a cyanylating agent in an inert solvent. The reaction is performed as in the
10 aforementioned step C1 in process C.


Step K2 is for producing compound (87) and implemented by hydrolyzing compound (86) in the presence of a base. The reaction is performed as in the aforementioned step C2 in process C.

15 Step K3 is for producing compound (88) and implemented by reacting compound (87) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (138) or acid addition salts thereof in an inert solvent. The reaction is performed as in the
20 aforementioned step C3 in process C.

Step K4 is for producing compound (89) and implemented by reacting compound (88) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

- 5 Process K' is an alternative method for producing compound (255) having the dashed line in compound (89) forming a double bond together with the solid line, and compound (199) having the dashed line in compound (89) forming a single bond together with the solid line.

10 Process K'



- Step K'1 is for producing compound (195) and implemented by reacting compound (192) with a cyanylating agent in an inert solvent. The reaction is performed as in the aforementioned step K1 in process K.
- 15

Step K'2 is for producing compound (196) and implemented by hydrolyzing compound (195) in the presence

of a base. The reaction is performed as in the
aforementioned step K2 in process K.

Step K'3 is for producing compound (197) and
implemented by reacting compound (196) or reactive
5 derivatives thereof (acid halides, mixed acid anhydrides or
active esters) with compound (138) or acid addition salts
thereof in an inert solvent. The reaction is performed as
in the aforementioned step K3 in process K.

Step K'4 is for producing compound (198) and
10 implemented by reacting compound (197) with an acid in an
aqueous solvent. The reaction is performed as in the
aforementioned step J'5 in process J'.


Step K'5 is for producing compound (255) and
implemented by reacting compound (198) with a reducing
15 agent in an optionally miscible inert solvent. The
reaction is performed as in the aforementioned step J'6 in
process J'.

Step K'6 is for producing compound (199) and
implemented by performing catalytic reduction of compound
20 (255) in an alcoholic solvent or an inert solvent or
reacting compound (255) with a reducing agent in an
optionally miscible inert solvent. The reaction is
performed as in the aforementioned step C'6 in process C'.

Process L is for producing compound (94) represented
25 by the general formula (I) in which X^1 is a group of β
configuration that is represented by the general formula
(II) in which Ar is an aromatic hydrocarbon group
(preferably a p-phenylene group), A is -O- and R^1 is -G-

NHCO-Z, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

Process L



Step L1 is for producing compound (91) and implemented by reacting compound (78) with phthalimide in an inert solvent in the presence of an azodicarboxylic acid dialkyl ester (preferably diethyl azodicarboxylate) and a phosphine compound (preferably triphenylphosphine). The reaction is performed as in the aforementioned step D1 in process D.

Step L2 is an alternative step for producing compound (91) and implemented by reacting compound (79) with a metal salt of phthalimide (preferably phthalimide potassium) in an inert solvent. The reaction is performed as in the

aforementioned step D2 in process D.


Step L3 is for producing compound (92) and implemented by reacting compound (91) with an amine-containing compound (preferably hydrazine) in an alcoholic solvent. The
5 reaction is performed as in the aforementioned step D3 in process D.

Step L4 is for producing compound (93) and implemented by reacting compound (139) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with
10 compound (92) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

Step L5 is for producing compound (94) and implemented by reacting compound (93) with an acid in an aqueous
15 solvent. The reaction is performed as in the aforementioned step A5 in process A.

Process L' is a method of producing compound (256) having the dashed line in compound (94) forming a double bond together with the solid line, and compound (199)
20 having the dashed line in compound (94) forming a single bond together with the solid line.

Process L'



Step L'1 is for producing compound (200) and implemented by reacting compound (191) with phthalimide in an inert solvent in the presence of an azodicarboxylic acid dialkyl ester (preferably diethyl azodicarboxylate) and a phosphine compound (preferably triphenylphosphine). The reaction is performed as in the aforementioned step L1 in process L.

Step L'2 is an alternative step for producing compound (200) and implemented by reacting compound (192) with a metal salt of phthalimide (preferably phthalimide potassium) in an inert solvent. The reaction is performed as in the aforementioned step L2 in process L.

Step L'3 is for producing compound (201) and

implemented by reacting compound (200) with an amine-containing compound (preferably hydrazine) in an alcoholic solvent. The reaction is performed as in the aforementioned step L3 in process L.

5 Step L'4 is for producing compound (202) and implemented by reacting compound (139) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (201) or acid addition salts thereof in an inert solvent. The reaction is performed as
10 in the aforementioned step L4 in process L.

 Step L'5 is for producing compound (203) and implemented by reacting compound (202) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step K'4 in process K'.


15 Step L'6 is for producing compound (256) and implemented by reacting compound (203) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step K'5 in
process K.

20 Step L'7 is for producing compound (199) and implemented by performing catalytic reduction of compound (256) in an alcoholic solvent or an inert solvent or by reacting compound (256) with a reducing agent in an optionally miscible inert solvent. The reaction is
25 performed as in the aforementioned step C'6 in process C'.

 Process M is for producing compound (102) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the

general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-(CH_2)_4-R^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together
5 with the solid line is a single bond or a double bond; compound (104) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is
10 $-(CH_2)_4-G^2-S(O)-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (105) represented by the general formula (I) in
15 which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-(CH_2)_4-G^2-S(O)_2-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which
20 they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

Process M



Step M1 is for producing compound (97) and implemented by reacting compound (96) with a base in an inert solvent to make a salt of compound (96) and then reacting it with compound (134) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

A compound having the hydroxyl group in 7-position of compound (96) oriented in β configuration is also known by being disclosed in, for example, J. Org. Che., 26, 2856-2859 (1961) and by using this compound in place of compound (96), one can obtain compounds having X^2 in compound (102), compound (104) and compound (105) oriented in β configuration.

Step M3 is for producing compound (100) and implemented by reacting compound (97) with compound (135) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the

aforementioned step A4 in process A.

Step M5 is for producing compound (101) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is
5 performed as in the aforementioned step A6 in process A.

Step M6 is for producing compound (102) and implemented by reacting compound (101) with a reducing agent in an optionally miscible inert solvent.

The inert solvent to be used is not limited in any
10 particular way as long as it does not interfere with the reaction; examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, alcoholic solvents such as methanol and ethanol, aromatic solvents such as benzene, toluene, xylene, quinoline and
15 chlorobenzene, and amines such as pyridine and triethylamine, and preferred examples are alcoholic solvents such as methanol and ethanol, with metanol being more preferred. The reducing agent to be used may be exemplified by: metal hydrogen complex compounds such as
20 aluminum lithium hydride, trimethoxyaluminum lithium hydride, tri-t-butoxyaluminum lithium hydride, aluminum lithium hydride-trichloroaluminum (alane), aluminum lithium hydride-boron trifluoride, aluminum hydride magnesium chloride, magnesium aluminum hydride, sodium aluminum
25 hydride, sodium triethoxyaluminum hydride, sodium bis(methoxyethoxy)aluminum hydride, sodium boron hydride, sodium boron hydride-palladium/carbon, sodium boron hydrogensulfide, sodium boron hydrogencyanide, sodium

trimethoxyboron hydride, lithium boron hydride, lithium boron hydrogencyanide, lithium triethylboron hydride, lithium tri-s-butylboron hydride, lithium tri-t-butylboron hydride, calcium boron hydride, potassium boron hydride, 5 potassium triisopropoxyboron hydride, potassium tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium boron hydride, and tetra-n-butylammonium cyanoboron hydride; metal hydrides such as diisobutylaluminum hydride, triphenyltin hydride, tri-n-butyltin hydride, diphenyltin 10 hydride, di-n-butyltin hydride, triethyltin hydride, trimethyltin hydride, trichlorosilane/tri-n-butylamine, trichlorosilane/tri-n-propylamine, triethylsilane, trimethylsilane, diphenylsilane, phenylsilane, polymethylhydrosiloxane, dimethylphenylsilane, di-n- 15 butylsilane, and methylphenylsilane; borane derivatives such as diborane, dimethylamine-borane, trimethylamine-borane, ethylenediamine-borane, pyridine-borane, dimethylsulfide-borane, 2,3-dimethyl-2-butylborane (thexylborane), bis-3-methyl-2-butylborane (disiamylborane), 20 diisopinocanepherylborane, dicyclohexylborane, and 9-borabicyclo[3,3,1]nonane (9-BBN); preferred examples are metal hydrogen complex compounds such as aluminum lithium hydride, trimethoxyaluminum lithium hydride, tri-t-butoxyaluminum lithium hydride, aluminum lithium hydride- 25 trichloroaluminum (alane), aluminum lithium hydride-boron trifluoride, aluminum hydride magnesium chloride, magnesium aluminum hydride, sodium aluminum hydride, sodium triethoxyaluminum hydride, sodium

bis(methoxyethoxy)aluminum hydride, sodium boron hydride, sodium boron hydride-palladium/carbon, sodium boron hydrogensulfide, sodium boron hydrogencyanide, sodium trimethoxyboron hydride, lithium boron hydride, lithium boron hydrogencyanide, lithium triethylboron hydride, lithium tri-s-butylboron hydride, lithium tri-t-butylboron hydride, calcium boron hydride, potassium boron hydride, potassium triisopropoxyboron hydride, potassium tri-s-butylboron hydride, zinc boron hydride, tetramethylammonium boron hydride, and tetra-n-butylammonium cyanoboron hydride, with sodium boron hydride being more preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of -30°C ~ 100°C , preferably 0°C ~ 70°C . The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 48 hours, preferably 30 minutes - 24 hours.

Step M8 is for producing compound (104) in the case where Q^2 in R^2 in compound (102) is -S- and implemented by reacting compound (102) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.


Step M9 is for producing compound (105) in the case where Q^2 in R^2 in compound (102) is -S- and implemented by reacting compound (102) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

Process N is for producing compound (112) represented

by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{R}^4$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a double bond; compound (113) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(\text{CH}_2)_3-\text{R}^4$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a double bond; compound (114) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(\text{CH}_2)_3-\text{R}^4$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (115) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(\text{CH}_2)_3-\text{G}^3-\text{S}(\text{O})-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which

they are bound, are $-(C=O)$, and the dashed line together with the solid line is a double bond; compound (116) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(CH_2)_3-G^3-S(O)_2-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a double bond; compound (117) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(CH_2)_3-G^3-S(O)-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; and compound (118) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-(CH_2)_3-G^3-S(O)_2-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond.

Process N



Step N3 is for producing compound (112) and implemented by reacting compound (108) with compound (257) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the aforementioned step A4 in process A.

Step N5 is for producing compound (113) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Step N6 is for producing compound (114) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent.

The solvent to be used may be exemplified by alcoholic solvents such as methanol, ethanol, n-propanol, i-propanol, n-butanol, s-butanol, t-butanol, pentanol, hexanol, cyclopropanol, cyclobutanol, cyclopentanol, cyclohexanol,

ethylene glycol, 1,3-propanediol, 1,4-butanediol, and 1,5-pentanediol, ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, 5 halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as cyclohexane, dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, ethyl acetate, acetonitrile and 10 nitromethane; preferred examples are methanol and ethanol.

The condition to be used in catalytic reduction is a homogeneous system such as hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-15 chlorotris(triparamethoxyphenylphosphine)rhodium(I), hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I), hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II) acetate, hydrogen-chlorohydridetris(triphenylphosphine)ruthenium(II), 20 hydrogen-carboxylatohydridetris(triphenylphosphine)ruthenium(II), hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I), hydrogen-platinum(II)-tin chloride complex, hydrogen-pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine 25 cobalt(II) complex, hydrogen-bis(dimethylglyoximato)cobalt(II) complex, hydrogen-methyl benzoate-tricarbonylchromium complex, hydrogen-bis(tricarbonylcyclopentadienylchromium), hydrogen-

pentacarbonyliron, hydrogen-
bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-
hydridecarbonylcobalt complex, hydrogen-
octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium,
5 hydrogen-chromium(III) acetylacetonato-triisobutylaluminum,
hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or
hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an
inhomogeneous system condition such as hydrogen-platinum
dioxide, hydrogen-platinum/carbon, hydrogen-
10 palladium/carbon, hydrogen-palladium/barium sulfate,
hydrogen-palladium/calcium carbonate, hydrogen-Raney nickel,
hydrogen-copper chromite, hydrogen-rhodium/carbon,
hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide, or
hydrogen-ruthenium/carbon; a preferred example is hydrogen-
15 palladium/carbon.

The reaction temperature is typically in the range of
0°C - 100°C, preferably 0°C - 60°C. The reaction time which
varies with the reaction temperature and the like is
typically in the range of 10 minutes - 24 hours, preferably
20 10 minutes - 6 hours.

Step N7 is for producing compound (115) in the case
where Q^4 in R^2 in compound (113) is -S- and implemented by
reacting compound (113) with an oxidizing agent in an inert
solvent. The reaction is performed as in the
25 aforementioned step A8 in process A.

Step N8 is for producing compound (116) in the case
where Q^4 in R^2 in compound (113) is -S- and implemented by
reacting compound (113) with an oxidizing agent in an inert

solvent. The reaction is performed as in the
aforementioned step A9 in process A.

Step N9 is for producing compound (117) in the case
where Q^4 in R^2 in compound (114) is -S- and implemented by
5 reacting compound (114) with an oxidizing agent in an inert
solvent. The reaction is performed as in the
aforementioned step A8 in process A.

Step N10 is for producing compound (118) in the case
where Q^4 in R^2 in compound (114) is -S- and implemented by
10 reacting compound (114) with an oxidizing agent in an inert
solvent. The reaction is performed as in the
aforementioned step A9 in process A.


Step N11 is an alternative method of producing
compound (114) and implemented by performing catalytic
15 reduction in an alcoholic solvent or an inert solvent. The
reaction is performed as in the aforementioned step N6 in
process N.

Process O is for producing compound (126) represented
by the general formula (I) in which X^1 is a hydrogen atom,
20 X^2 is a group of α configuration that is represented by the
general formula (II) in which Ar is an aromatic hydrocarbon
group (preferably a p-phenylene group), A is -O- and R^1 is
-CH₂-CH=CH-CH₂-R², R^a is a hydrogen atom, R^b and R^c, when
taken together with the carbon atom in 3-position to which
25 they are bound, are -(C=O)-, and the dashed line together
with the solid line is a double bond; compound (127)
represented by the general formula (I) in which X^1 is a
hydrogen atom, X^2 is a group of α configuration that is

represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-(CH_2)_4-R^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a double bond; compound (128) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-(CH_2)_4-R^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; compound (129) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-(CH_2)_4-G^2-S(O)-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a double bond; compound (130) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group, A is -O- and R^1 is $-(CH_2)_4-G^2-S(O)_2-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-

position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a double bond; compound (131) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-(CH_2)_4-G^2-S(O)-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond; and compound (132) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-(CH_2)_4-G^2-S(O)_2-Z$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond.

20 Process O



Step O1 is for producing compound (120) and implemented by reacting compound (143) with a metal (preferably magnesium) or an alkylolithium (preferably t-butylolithium) in an inert solvent to make a reactive derivative of compound (143) and reacting it with compound (119) in an inert solvent in the presence of an additive (preferably tetrakis[(tri-n-butylphosphine)copper(I)]

iodide])).

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ethers such as ether, tetrahydrofuran, dioxane and dimethoxyethane, with ether being further preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of $-78^{\circ}\text{C} \sim 80^{\circ}\text{C}$ (preferably $-78^{\circ}\text{C} \sim 50^{\circ}\text{C}$). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

As a by-product of the production of compound (12), there is formed a compound having the $-\text{C}_6\text{H}_4-\text{OR}^3$ in 7-position of compound (12) and by using this compound in place of compound (120), one can obtain compounds having X^2 in compound (126), compound (127), compound (128), compound (129), compound (130), compound (131) and compound (132) oriented in β configuration.

Step O2 is for producing compound (121) and implemented by reacting compound (120) with a deprotecting agent, namely by removing the substituted silyl group, in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not interfere with the reaction; examples include ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, as well as dimethylformamide and water, with tetrahydrofuran being preferred. The deprotecting agent to be used is not

limited in any particular way and may be exemplified by fluorides such as hydrogen fluoride, hydrogen fluoride-pyridine, sodium fluoride, potassium fluoride and tetra-n-butylammonium fluoride, inorganic acids such as

5 hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and phosphoric acid, and organic acids such as formic acid, acetic acid and p-toluenesulfonic acid, with tetra-n-butylammonium fluoride being preferred.

The reaction temperature which varies with the type of

10 solvent and the like is typically in the range of 0°C - 80°C (preferably 0°C - 50°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

15 Step 03 is for producing compound (122) and implemented by reacting compound (121) with a base in an inert solvent to make a salt of compound (121) and then reacting it with compound (134) in an inert solvent. The reaction is performed as in the aforementioned step A3 in

20 process A.

Step 05 is for producing compound (125) and implemented by reacting compound (122) with compound (135) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the

25 aforementioned step A4 in process A.

Step 06 is for producing compound (126) and implemented by hydrolyzing compound (125) in water or a water-soluble solvent in the presence of a base or an acid

(preferably a base).

The water-soluble solvent to be used is not limited in any particular way and may be exemplified by alcoholic solvents such as methanol, ethanol, n-propanol and i-propanol, ether solvents such as tetrahydrofuran and dioxane, as well as dimethylformamide, with methanol being preferred.

The base to be used is not limited in any particular way and may be exemplified by metal hydroxides such as lithium hydroxide, sodium hydroxide, potassium hydroxide, potassium hydroxide, calcium hydroxide, barium hydroxide and cesium hydroxide, and carbonates such as potassium carbonate and sodium carbonate, with sodium hydroxide being preferred.

The acid to be used is not limited in any particular way and may be exemplified by inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and phosphoric acid, with hydrochloric acid being preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 100°C (preferably 0°C - 80°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

Step O8 is for producing compound (127) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is

performed as in the aforementioned step A6 in process A.

Step O9 is for producing compound (128) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step N6 in process N.

Step O10 is for producing compound (129) in the case where Q^2 in R^2 in compound (127) is -S- and implemented by reacting compound (127) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.

Step O11 is for producing compound (130) in the case where Q^2 in R^2 in compound (127) is -S- and implemented by reacting compound (127) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.


Step O12 is for producing compound (131) in the case where Q^2 in R^2 in compound (128) is -S- and implemented by reacting compound (128) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.


Step O13 is for producing compound (132) in the case where Q^2 in R^2 in compound (128) is -S- and implemented by reacting compound (128) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

Process P is for producing compound (217) represented by the general formula (I) in which X^2 is a hydrogen atom, X^1 is a group of β configuration that is represented by the

general formula (II) in which Ar is single bond, A is a methylene group and R^1 is a group represented by the general formula (III) in which G is $-G^4-CH_2-$, E is a single bond, J is an optionally substituted aromatic hydrocarbon group (preferably a p-phenylene group), Y is a single bond, L is L^2 , Q is Q^{17} , with R^7 in Q^{17} being a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

Process P



Process P (continued)

Step P1 is for producing compound (205) and implemented by reacting compound (204) with an alkyllithium (preferably n-butyllithium) in an inert solvent to make a reactive derivative of compound (204) and reacting it with compound (164) in an inert solvent. The reaction is performed as in the aforementioned step F1 in process F.

Step P2 is for producing compound (206) and implemented by reacting compound (205) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step F2 in process F.

In this step, a compound having the substituent in the 11-position of compound (206) oriented in α configuration forms as a by-product and this may be used to give a compound having X^1 in compound (217) oriented in α configuration.

Step P3 is for producing compound (207) and implemented by reacting compound (206) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the
5 aforementioned step F3 in process F.

Step P4 is for producing compound (208) and implemented by reacting compound (207) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (207) with a halogenating agent in an
10 inert solvent. The reaction is performed as in the aforementioned step F4 in process F.

Step P5 is for producing compound (209) and implemented by reacting compound (218) with a metal (preferably magnesium) or an alkylolithium (preferably n-
15 butyllithium) in an inert solvent to make a reactive derivative of compound (218) and reacting it with compound (208) in an inert solvent. The reaction is performed as in the aforementioned step I1 in process I.

Step P6 is for producing compound (210) and
20 implemented by reacting compound (209) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the aforementioned step P3 in process P.

Step P7 is for producing compound (211) and
25 implemented by reacting compound (210) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (210) with a halogenating agent in an inert solvent. The reaction is performed as in the

aforementioned stepp P4 in process P.

Step P8 is for producing compound (212) and implemented by reacting compound (219) with a base in an inert solvent to make a reactive derivative of compound (219) and then reacting it with compound (211) in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, as well as dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, and N-methylpyrrolidone; preferred examples are ether solvents such as tetrahydrofuran, as well as dimethylformamide. The base to be used may be exemplified by metal alkoxides such as sodium alkoxide and potassium t-butoxide, metal hydrides such as sodium hydride, potassium hydride and calcium hydride, alkyl lithium compounds such as methyllithium, ethyllithium, n-butyllithium and t-butyllithium, metal amides such as sodium amide, potassium bistrimethylsilylamide, sodium bistrimethylsilylamide and lithium diisopropylamide, as well as carbonates such as cesium carbonate, potassium carbonate and sodium carbonate; preferred examples are metal hydrides such as sodium hydride, metal amides such as lithium diisopropylamide, and carbonates such as cesium carbonate.

The reaction temperature which varies with the type of solvent and the like is typically in the range of -78°C ~ 80°C , preferably 0°C ~ 30°C . The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 30 minutes - 15 hours.

Step P9 is for producing compound (213) and implemented by reacting compound (212) with a base in an inert solvent to make a reactive derivative of compound (212) and then reacting it with compound (220) in an inert solvent. The reaction is performed as in the aforementioned step P8 in process P.

If Z is a hydrogen atom in process P, step P9 may be omitted.

Step P10 is for producing compound (214) and implemented by reacting compound (213) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step L'5 in process L'.

Step P11 is for producing compound (215) and implemented by reacting compound (214) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step L'6 in process L'.

Step P12 is for producing compound (216) and implemented by reacting compound (215) with an acid, a base or a metal salt in an hydrous alcohol or an inert solvent.

The solvents to be used are not limited in any particular way as long as they do not interfere with the

reaction; examples are mixtures of water and alcoholic solvents such as methanol, ethanol, n-propanol, i-propanol, n-butanol, s-butanol and t-butanol, amine-containing solvents such as pyridine, as well as dimethyl sulfoxide, 5 dimethylacetamide, dimethylimidazolidinone and dimethylformamide; preferred examples are a mixture of water and an alcoholic solvent such as methanol or ethanol, and dimethyl sulfoxide.

The acid to be used may be exemplified by inorganic 10 acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and phosphoric acid, with hydrochloric acid and hydrobromic acid being preferred.

The base to be used may be exemplified by metal hydroxides such as lithium hydroxide, sodium hydroxide, 15 potassium hydroxide, potassium hydroxide, calcium hydroxide, barium hydroxide and cesium hydroxide, with sodium hydroxide and potassium hydroxide being preferred.

The metal salt to be used can be lithium chloride, sodium cyanide, etc., with lithium chloride being preferred.

20 The reaction temperature which varies with the type of solvent and the like is typically in the range of 25°C - 180°C, preferably 40°C - 150°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 25 30 minutes - 15 hours.

Step P13 is for producing compound (217) and implemented by performing catalytic reduction of compound (216) in an alcoholic solvent or an inert solvent. The

reaction is performed as in the aforementioned step F8 in process F.

As an ancillary to this reaction, conversion to a single bond may occasionally be effected if the dashed line
5 forms a double bond together with the solid line.

Steps P14 and P15 provide an alternative method of producing compound (209).

Step P14 is for producing compound (221) and implemented by reacting compound (223) with an alkyllithium
10 (preferably n-butyllithium) in an inert solvent to make a reactive derivative of compound (223) and reacting it with compound (164) in an inert solvent. The reaction is performed as in the aforementioned step P1 in process P.

Step P15 is for producing compound (209) and
15 implemented by reacting compound (209) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step P2 in process P.


In this step, a compound having the substituent in the
20 11-position of compound (209) oriented in α configuration forms as a by-product and this may be used to give a compound having X^1 in compound (217) oriented in α configuration.

Step P16 is for producing compound (222) and
25 implemented by performing catalytic reduction of compound (209) in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step P13 in process P.

By subjecting compound (222) to step P6 as in the case of compound (209), one can produce a compound having the dashed line in compound (217) forming a single bond together with the solid line.

- 5 Process Q is for producing compound (234) represented by the general formula (I) in which X^2 is a hydrogen atom, X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is single bond, A is a methylene group and R^1 is a group represented by the
- 10 general formula (III) in which G is $-(CH_2)_2-CH(OH)-G^3-$, E, J, Y and L are single bonds, and Q is Q^{63} , R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond
- 15 or a double bond; and compound (235) represented by the general formula (I) in which X^2 is a hydrogen atom, X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is single bond, A is a methylene group and R^1 is a group represented by the general formula
- 20 (III) in which G is $-(CH_2)_2-CH(OH)-G^3-$, E, J, Y and L are single bonds, and Q is Q^{64} , R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double
- 25 bond.

Process Q



Step Q1 is for producing compound (225) and implemented by reacting compound (224) with a metal

(preferably magnesium) or an alkyllithium (preferably t-butylolithium) in an inert solvent in the presence or absence (preferably the presence) of an additive {preferably mercury(II) chloride} to make a reactive
5 derivative of compound (224) and reacting it with compound (183) in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ether solvents such as
10 ether, tetrahydrofuran, dioxane and dimethoxyethane, with ether and tetrahydrofuran being more preferred.

The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 80°C (preferably 10°C - 50°C). The reaction time which varies
15 with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 15 hours).

Step Q2 is for producing compound (226) and implemented by reacting compound (225) with an alkyllithium
20 (preferably n-butylolithium) in an inert solvent to make a reactive derivative of compound (225) and reacting it with compound (164) in an inert solvent. The reaction is performed as in the aforementioned step F1 in process F.

Step Q3 is for producing compound (227) and
25 implemented by reacting compound (226) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step F2 in process F.

As a by-product of this step, there is formed a compound having the substituent in 11-position of compound (227) oriented in α configuration; by using this compound, one can obtain compounds having X^1 in compound (234) and
5 compound (235) oriented in α configuration.

Step Q4 is for producing compound (228) and implemented by reacting compound (227) with a deprotecting agent, namely, removing the substituted silyl group, in an inert solvent. The reaction is performed as in the
10 aforementioned step F3 in process F.

Step Q5 is for producing compound (229) and implemented by reacting compound (228) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (228) with a halogenating agent in an
15 inert solvent. The reaction is performed as in the aforementioned step F4 in process F.

Step Q6 is for producing compound (230) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive
20 derivative of compound (137) and reacting it with compound (229) in an alcoholic solvent. The reaction is performed as in the aforementioned step F5 in process F.

Step Q7 is for producing compound (231) and implemented by reacting compound (230) with an acid in an
25 aqueous solvent. The reaction is performed as in the aforementioned step P10 in process P.

Step Q8 is for producing compound (232) and implemented by reacting compound (231) with a reducing

agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step P11 in process P.

Step Q9 is for producing compound (233) and
5 implemented by reacting compound (232) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step F6 in process F.

Step Q10 is for producing compound (234) and
implemented by performing catalytic reduction of compound
10 (233) in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step F8 in process F.

As an ancillary to this reaction, conversion to a single bond may occasionally be effected if the dashed line
15 forms a double bond together with the solid line.

Step Q11 is for producing compound (235) and implemented by reacting compound (234) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step B7 in process B.


20 In process Q, step Q5, step Q6, step Q9 and step Q11 may be omitted and by so doing, one can produce a compound represented by the general formula (I) in which X^2 is a hydrogen atom, X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is
25 single bond, A is a methylene group and R^1 is a group represented by the general formula (III) in which G is $-(CH_2)_2-CH(OH)-G^3-$, E, J, Y, L and Q are single bonds, and Z is $-O-R^d$, R^a is a hydrogen atom, R^b and R^c , when taken

together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

In process Q, the hydroxyl group on G may optionally
5 be subjected to a protecting reaction and a deprotecting reaction in any desired steps.

Process R is a method for producing compound (242) represented by the general formula (I) in which X^2 is a hydrogen atom, X^1 is a group of β configuration that is
10 represented by the general formula (II) in which Ar is single bond, A is a methylene group and R^1 is a group represented by the general formula (III) in which G is $-(CH_2)_2-CH(OH)-G^3-$, E, J, Y, L and Q are single bonds, and Z is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when
15 taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond.

Process R



Step R1 is for producing compound (237) and implemented by reacting compound (224) with a metal (preferably magnesium) or an alkyllithium (preferably t-butylolithium) in an inert solvent in the presence or absence (preferably the presence) of an additive {preferably mercury(II) chloride} to make a reactive derivative of compound (224) and reacting it with compound (236) in an inert solvent. The reaction is performed as in the aforementioned step Q1 in process Q.

Step R2 is for producing compound (238) and implemented by reacting compound (237) with an alkyllithium (preferably n-butylolithium) in an inert solvent to make a reactive derivative of compound (237) and reacting it with

compound (164) in an inert solvent. The reaction is performed as in the aforementioned step Q2 in process Q.

Step R3 is for producing compound (239) and implemented by reacting compound (238) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step Q3 in process Q.

As a by-product of this step, there is formed a compound having the substituent in 11-position of compound (239) oriented in α configuration; by using this compound, one can obtain a compound having X^1 in compound (242) oriented in α configuration.

Step R4 is for producing compound (240) and implemented by reacting compound (239) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step Q7 in process Q.

Step R5 is for producing compound (241) and implemented by reacting compound (240) with a reducing agent in an optionally miscible inert solvent. The reaction is performed as in the aforementioned step Q8 in process Q.

Step R6 is for producing compound (242) and implemented by performing catalytic reduction of compound (241) in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step Q10 in process Q.


As an ancillary to this reaction, conversion to a single bond may occasionally be effected if the dashed line

forms a double bond together with the solid line.

In process R, the hydroxyl group on G may optionally be subjected to a protecting reaction and a deprotecting reaction in any desired steps.

- 5 Process S is for producing compound (244) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{COOH}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (245) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CON}(\text{R}^7)\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond.

Process S



Step S1 is for producing compound (243) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Step S2 is for producing compound (244) in the case where Q² in R² in compound (243) is Q¹⁷ and Z is a hydrogen atom and this is implemented by reacting compound (243) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.


Step S3 is for producing compound (245) and implemented by reacting compound (244) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

In process S, compound (245) can be produced whether the sequence of reaction steps is S2 → S3 → S1 or S2 → S1 → S3.

Process S' is for producing compound (251) represented

by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, 5 R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (254) represented by the general formula (I) in which X^1 is a group of β 10 configuration that is represented by the general formula (II) in which Ar is a single bond, A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}_2-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the 15 dashed line together with the solid line is a single bond or a double bond.

Process S'



Step S'1 is for producing compound (247) and implemented by reacting compound (4) with compound (246) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the aforementioned step A4 in process A.

Step S'2 is for producing compound (248) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and reacting it with compound (247) in an alcoholic solvent. The reaction is performed

as in the aforementioned step B4 in process B.

Step S'3 is for producing compound (249) and implemented by reacting compound (248) with an oxidizing agent in an inert solvent. The reaction is performed as in
5 the aforementioned step A8 in process A.

Step S'4 is for producing compound (250) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

10 Step S'5 is for producing compound (251) and implemented by reacting compound (250) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

Step S'6 is for producing compound (252) and
15 implemented by reacting compound (248) or compound (249) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

Step S'7 is for producing compound (253) and implemented by performing catalytic reduction in an
20 alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

This step can also be implemented by using compound (250) as the starting material.

Step S'8 is for producing compound (254) and
25 implemented by reacting compound (253) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

This step can also be implemented by using compound


(251) as the starting material.

In process S', compound (251) can be produced from compound (247) whether the sequence of reaction steps is S'4 \rightarrow S'2 \rightarrow S'5 \rightarrow S'3, or S'5 \rightarrow S'4 \rightarrow S'2 \rightarrow S'3, or S'4 \rightarrow S'5 \rightarrow S'2 \rightarrow S'3, or S'5 \rightarrow S'2 \rightarrow S'3 \rightarrow S'4, and compound (254) can be produced from compound (247) whether the sequence of reaction steps is S'4 \rightarrow S'2 \rightarrow S'5 \rightarrow S'6, or S'5 \rightarrow S'4 \rightarrow S'2 \rightarrow S'6, or S'4 \rightarrow S'5 \rightarrow S'2 \rightarrow S'6, or S'5 \rightarrow S'2 \rightarrow S'6 \rightarrow S'4.

Process T is for producing compound (261) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R¹ is -CH₂-CH₂-CH₂-G²-COOR⁷, X² is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; compound (262) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R¹ is -CH₂-CH₂-CH₂-G²-COOH, X² is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; and compound (263) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general

5

Process T



10

The inert solvent to be used is not limited in any particular way as long as it does not participate in the

reaction; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, with tetrahydrofuran being more preferred. Preferred examples of the base to be used are n-butyllithium and lithium diisopropylamide. The reaction temperature which varies with the type of solvent and the like is typically in the range of -100°C ~ 50°C , preferably -78°C ~ 30°C . The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 48 hours, preferably 30 minutes - 24 hours.

Step T2 is for producing compound (257) and implemented by reacting compound (256) with compound (326) in an inert solvent in the presence of a metal catalyst.

The inert solvent to be used is not limited in any particular way as long as it does not participate in the reaction; preferred examples are ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, with tetrahydrofuran being more preferred. The metal catalyst to be used is not limited in any particular way and may be exemplified by tetrakis(triphenylphosphine)palladium, palladium(II) acetate-triphenylphosphine, bis(triphenylphosphine)palladium(II) chloride, etc, with tetrakis(triphenylphosphine)palladium being preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 100°C , preferably 10°C - 80°C . The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 48 hours, preferably 30 minutes - 24

hours.

Step T3 is for producing compound (259) and implemented by reacting compound (257) with compound (258) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the
5 aforementioned step A4 in process A.

Step T4 is for producing compound (260) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent.

10 The solvent to be used may be exemplified by alcoholic solvents such as methanol, ethanol, n-propanol, i-propanol, n-butanol, s-butanol, t-butanol, pentanol, hexanol, cyclopropanol, cyclobutanol, cyclopentanol, cyclohexanol, ethylene glycol, 1,3-propanediol, 1,4-butanediol and 1,5-
15 pentanediol, ether solvents such as ether, tetrahydrofuran, dioxane and dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as cyclohexane,
20 dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, ethyl acetate, acetonitrile and nitromethane; preferred examples are ethanol, dioxane, benzene and ethyl acetate.

25 The condition to be used in catalytic reduction is a homogeneous system such as hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-

- chlorotris(triparamethoxyphenylphosphine)rhodium(I),
hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I),
hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II)
acetate, hydrogen-
- 5 chlorohydridetris(triphenylphosphine)ruthenium(II),
hydrogen-
carboxylatohydridetris(triphenylphosphine)ruthenium(II),
hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I),
hydrogen-platinum(II)-tin chloride complex, hydrogen-
- 10 pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine
cobalt(II) complex, hydrogen-
bis(dimethylglyoximato)cobalt(II) complex, hydrogen-methyl
benzoate-tricarbonylchromium complex, hydrogen-
bis(tricarbonylcyclopentadienylchromium), hydrogen-
- 15 pentacarbonyliron, hydrogen-
bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-
hydridecarbonylcobalt complex, hydrogen-
octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium,
hydrogen-chromium(III) acetylacetonato-triisobutylaluminum,
- 20 hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or
hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an
inhomogeneous system condition such as hydrogen-platinum
dioxide, hydrogen-platinum/carbon, hydrogen-
palladium/carbon, hydrogen-palladium hydroxide/carbon,
- 25 hydrogen-palladium/barium sulfate, hydrogen-
palladium/calcium carbonate, hydrogen-Raney nickel,
hydrogen-copper chromite, hydrogen-rhodium/carbon,
hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide,

hydrogen-ruthenium/carbon, or hydrogen-iridium black; a preferred example is hydrogen-iridium black.

The reaction temperature is typically in the range of 0°C - 100°C, preferably 0°C - 60°C. The reaction time which
5 varies with the reaction temperature and the like is typically in the range of 10 minutes - 100 hours, preferably 10 hours - 96 hours.

Step T5 is for producing compound (261) and implemented by reacting compound (260) with an acid in an
10 aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

In step T5, ester hydrolysis may occur and in that case, subsequent step T6 may be omitted.


Step T6 is for producing compound (262) and
15 implemented by hydrolyzing compound (261) in water or a water-soluble solvent in the presence of a base or an acid (preferably a base). The reaction is performed as in the aforementioned step O6 in process O.

Step T7 is for producing compound (263) and
20 implemented by reacting compound (262) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

25 Process T' is for producing compound (268) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group

and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CH}_2-\text{S}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond; compound (269) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CH}_2-\text{SO}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (270) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CH}_2-\text{SO}_2-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond or a double bond.

Process T'



Step T'1 is for producing compound (265) and implemented by reacting compound (260) with a reducing agent in an inert solvent and the reaction is performed as in the aforementioned step A2 in process A.

Step T'2 is for producing compound (266) and implemented by reacting compound (265) with a sulfonyl chloride compound in an amine-containing solvent or reacting compound (265) with a halogenating agent in an inert solvent. The reaction is performed as in the aforementioned step B3 in process B.

Step T'3 is for producing compound (267) and implemented by reacting compound (137) with a metal

alkoxide in an alcoholic solvent to make a reactive derivative of compound (137) and reacting it with compound (266) in an alcoholic solvent. The reaction is performed as in the aforementioned step B4 in process B.

5 Step T'4 is for producing compound (268) and implemented by reacting compound (267) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

10 Step T'5 is for producing compound (269) and implemented by reacting compound (268) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.

15 Step T'6 is for producing compound (270) and implemented by reacting compound (269) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

20 Step T'7 is an alternative method for producing compound (270) and implemented by reacting compound (268) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.


25 In process T', step T'3 and step T'4 may be interchanged in their sequence. If desired, step T'4 and step T'5 may also be interchanged in their sequence. Compound (270) can also be obtained from compound (267) if the sequence of reaction steps is T'6 → T'4.

Process U is for producing compound (276) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula

(II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R¹ is a methyl group, X² is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; compound (278) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R¹ is -CH₂-CH=CH₂, X² is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; compound (279) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R¹ is -CH₂-CH=CH-CH₂-R², X² is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; compound (280) represented by the general formula (I) in which X¹ is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R¹ is -CH₂-CH₂-CH₂-CH₂-R², X² is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-

position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; compound (281) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{COOH}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (282) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CON}(R^7)Z$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond

Process U



Step U1 is for producing compound (272) and implemented by reacting compound (271) with a metal (preferably magnesium) or an alkyl lithium (preferably n-butyl lithium) in an inert solvent to make a reactive derivative of compound (271) and reacting it with compound (2) in an inert solvent. The reaction is performed as in the aforementioned step I1 in process I.

Step U2 is for producing compound (273) and

implemented by reacting compound (272) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step E2 in process E.

- 5 Step U3 is for producing compound (274) and implemented by reacting compound (272) with an acid in an aqueous solvent.

10 The solvent to be used is not limited in any particular way as long as it does not interfere with the reaction; examples are mixtures of water and ether solvents such as ether, tetrahydrofuran and dioxane, alcoholic solvents such as methanol and ethanol, or ketonic solvents such as acetone, and hydrous acetone is preferred.

15 The acid to be used may be exemplified by inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and phosphoric acid, and organic acids such as acetic acid, p-toluenesulfonic acid and pyridinium-p-toluenesulfonate, with hydrochloric acid being preferred. The reaction temperature which varies
20 with the type of solvent and the like is typically in the range of 0°C - 50°C (preferably 10°C - 30°C). The reaction time which varies with the reaction temperature and the like is typically in the range of 15 minutes - 24 hours (preferably 30 minutes - 5 hours).

- 25 Step U4 is for producing compound (276) and implemented by reacting compound (273) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step U3 in process U.

Step U5 is for producing compound (275) and implemented by reacting compound (274) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step E2
5 in process E.

Step U6 is an alternative method for producing compound (276) and implemented by reacting compound (275) with an oxidizing agent in an inert solvent.

The inert solvent to be used is not limited in any
10 particular way as long as it does not interfere with the reaction; examples are halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, ethers such as tetrahydrofuran, dioxane and dimethoxyethane, and hydrocarbon solvents such as benzene, toluene, xylene,
15 quinoline and chlorobenzene, with dichloromethane and tetrahydrofuran being preferred. Water may optionally be added to these solvents. The oxidizing agent to be used is not limited in any particular way and examples can be manganese compounds such as potassium permanganate,
20 manganese dioxide, manganese(III) acetate, tris(acetonylacetonite)manganese(III) (MTA), manganese sulfate and manganese(III) pyrophosphate, chromates such as chromium(IV) oxide, Jones reagent, Sarett reagent, Collins reagent, chromic acid t-butyl ester, potassium bichromate,
25 Beckmann's mixture, sodium bichromate, Kiliani reagent, chromyl chloride, chromyl acetate, pyridinium chlorochromate (PCC), and pyridinium dichromate (PDC); ruthenium compounds such as ruthenium tetroxide,

- tris(triphenylphosphine)dichlororuthenium/iodosylbenzene,
tris(triphenylphosphine)dichlororuthenium/N-
methylmorpholin-N-oxide,
tris(triphenylphosphine)dichlororuthenium/t-butyl
- 5 hydroperoxide, tetrapropylammonium perruthenate (TPAP),
tetrapropylammonium perruthenate (TPAP)/N-methylmorpholin-
N-oxide, tetrabutylammonium perruthenate (TBAP), and
tetrabutylammonium perruthenate (TBAP)/N-methylmorpholin-N-
oxide; halogens such as hypochlorous acid, sodium
- 10 hypochlorite, potassium hypobromite, potassium hypiodite,
sodium chlorate, potassium chlorate, sodium bromate,
potassium bromate, sodium iodate, potassium iodate,
perchloryl fluoride, orthoperiodic acid, sodium
metaperiodate, potassium metaperiodate, N-bromoacetamide,
- 15 N-bromosuccinimide and N-bromophthalimide; as well as
dimethyl sulfoxide/oxalyl chloride; preferred examples are
chromates such as pyridinium chlorochromate (PCC) and
pyridinium dichromate (PDC), and ruthenium compounds such
tetrapropylammonium perruthenate (TPAP)/N-methylmorpholin-
- 20 N-oxide. The reaction temperature which varies with the
type of solvent and the like is typically in the range of
-30°C ~ 100°C, preferably 0°C - 30°C. The reaction time
which varies with the reaction temperature and the like is
typically in the range of 10 minutes - 48 hours, preferably
- 25 30 minutes - 24 hours.

Step U7 is an alternative method of producing compound (277) and implemented by reacting compound (276) with a deprotecting agent in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not interfere with the reaction; examples are dimethyl sulfoxide, dimethylacetamide, dimethylimidazolidinone, 5 dimethylformamide, N-methylpyrrolidone, etc., with dimethylformamide being preferred. The deprotecting agent to be used is not limited in any particular way and may be exemplified by sodium thiomethoxide, sodium cyanide, trimethylsilane iodide, boron tribromide, boron trichloride 10 and lithium chloride, with sodium thiomethoxide being preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of -80°C ~ 200°C, preferably 0°C ~ 180°C. The reaction time which varies with the reaction temperature and the like is 15 typically in the range of 10 minutes - 96 hours, preferably 30 minutes - 48 hours.

Step U8 is for producing compound (278) and implemented by reacting compound (277) with a base in an inert solvent to make a salt of compound (277) and then 20 reacting it with compound (134) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

Step U9 is for producing compound (279) and implemented by reacting compound (135) with compound (278) 25 in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the aforementioned step A4 in process A.

Step U10 is for producing compound (280) and

implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Step U11 is for producing compound (281) in the case
 5 where R^2 in compound (280) is G^2-COOR^7 and this is implemented by reacting compound (280) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.

Step U12 is for producing compound (282) and
 10 implemented by reacting compound (281) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

15 Process U' is for producing compound (283) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is an
 20 allyl group, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; compound (284) represented by the general
 25 formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-CH_2-CH=CH-G^3-R^5$, X^2 is a hydrogen

atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; compound (285)

5 represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{G}^3-\text{S}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom,

10 R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; compound (286) represented by the general formula (I) in which X^1 is a group of β configuration that


15 is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{G}^3-\text{SO}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they

20 are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond or a double bond; compound (287) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon

25 group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{G}^3-\text{SO}_2-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the

dashed line together with the solid line is a single bond or a double bond; compound (288) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond or a double bond; and compound (289) represented by the general formula (I) in which X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}_2-\text{Z}$, X^2 is a hydrogen atom, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond or a double bond.

20 Process U'



Step U'1 is for producing compound (283) and implemented by reacting compound (278) with an acid in an aqueous solvent. The reaction is performed as in the
 5 aforementioned step A5 in process A.

Step U'2 is for producing compound (284) and implemented by reacting compound (246) with compound (283) in an inert solvent in the presence of an organometallic

catalyst. The reaction is performed as in the
aforementioned step A4 in process A.

Step U'3 is for producing compound (285) and
implemented by reacting compound (137) with a metal
5 alkoxide in an alcoholic solvent to make a reactive
derivative of compound (137) and then reacting it with
compound (284) in an alcoholic solvent. The reaction is
performed as in the aforementioned step B4 in process B.

Step U'4 is for producing compound (286) and
10 implemented by reacting compound (285) with an oxidizing
agent in an inert solvent. The reaction is performed as in
the aforementioned step A8 in process A.

Step U'5 is for producing compound (287) and
implemented by reacting compound (285) with an oxidizing
15 agent in an inert solvent. The reaction is performed as in
the aforementioned step A9 in process A.

Step U'6 is an alternative method for producing
compound (287) and implemented by reacting compound (286)
with an oxidizing agent in an inert solvent. The reaction
20 is performed as in the aforementioned step A9 in process A.

Step U'7 is for producing compound (288) and
implemented by performing catalytic reduction in an
alcoholic solvent or an inert solvent. The reaction is
performed as in the aforementioned step A6 in process A.


25 Step U'8 is an alternative method for producing
compound (289) and implemented by reacting compound (288)
with an oxidizing agent in an inert solvent. The reaction
is performed as in the aforementioned step A9 in process A.

Step U'9 is an alternative method for producing compound (289) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in
5 process A.

In process U', compound (288) can be obtained from compound (284) if the sequence of reaction steps is U'7 → U'3 → U'4. Compound (289) can also be obtained from compound (284) if the sequence of reaction steps is U'7 →
10 U'3 → U'5.

Process U" is an alternative method to process U' for producing compound (273) and compound (275).

Process U"



Step U''1 is for producing compound (291) and implemented by reacting compound (256) with compound (290) in an optionally miscible inert solvent in the presence of a metal catalyst and a base.

The inert solvent to be used is not limited in any particular way as long as it does not interfere with the reaction; examples are ether solvents such as dioxane and tetrahydrofuran, aromatic hydrocarbon solvents such as toluene, alcoholic solvents such as ethanol, as well as dimethylformamide, dimethylacetamide and acetonitrile, with dioxane and ethanol-toluene being preferred. The metal

- catalyst to be used is not limited in any particular way and may be exemplified by tetrakis(triphenylphosphine)palladium, palladium(II) acetate-triphenylphosphine,
- 5 bis(triphenylphosphine)palladium(II) chloride, etc., with tetrakis(triphenylphosphine)palladium being preferred. The base to be used is not limited in any particular way and may be exemplified by potassium phosphate, sodium carbonate, etc., with sodium carbonate being preferred.
- 10 The reaction temperature which varies with the type of solvent and the like is typically in the range of 0°C - 180°C, preferably 10°C - 120°C. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 48 hours, preferably
- 15 30 minutes - 24 hours.

Step U² is for producing compound (273) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The solvent to be used may be exemplified by alcoholic solvents such as

20 methanol, ethanol, n-propanol, i-propanol, n-butanol, s-butanol, t-butanol, pentanol, hexanol, cyclopropanol, cyclobutanol, cyclopentanol, cyclohexanol, ethylene glycol, 1,3-propanediol, 1,4-butanediol and 1,5-pentanediol, ether solvents such as ether, tetrahydrofuran, dioxane and

25 dimethoxyethane, aromatic solvents such as benzene, toluene, xylene, quinoline and chlorobenzene, halogen-containing solvents such as dichloromethane, chloroform and carbon tetrachloride, as well as cyclohexane, dimethyl sulfoxide,

dimethylacetamide, dimethylimidazolidinone, dimethylformamide, N-methylpyrrolidone, ethyl acetate, acetonitrile and nitromethane; preferred examples are ethanol, dioxane, benzene, ethyl acetate and acetonitrile.

- 5 The condition to be used in catalytic reduction is a homogeneous system such as hydrogen-chlorotris(triphenylphosphine)rhodium(I), hydrogen-chlorotris(triparatolylphosphine)rhodium(I), hydrogen-chlorotris(triparamethoxyphenylphosphine)rhodium(I),
- 10 hydrogen-hydridecarbonyltris(triphenylphosphine)rhodium(I), hydrogen-rhodium(II) acetate, hydrogen-ruthenium(II) acetate, hydrogen-chlorohydridetris(triphenylphosphine)ruthenium(II), hydrogen-
- 15 carboxylatohydridetris(triphenylphosphine)ruthenium(II), hydrogen-hydridecarbonyltris(triphenylphosphine)iridium(I), hydrogen-platinum(II)-tin chloride complex, hydrogen-pentacyanocobalt(II) complex, hydrogen-tricyanobipyridine cobalt(II) complex, hydrogen-
- 20 bis(dimethylglyoximato)cobalt(II) complex, hydrogen-methyl benzoate-tricarbonylchromium complex, hydrogen-bis(tricarbonylcyclopentadienylchromium), hydrogen-pentacarbonyliron, hydrogen-bis(cyclopentadienyl)dicarbonyltitanium, hydrogen-
- 25 hydridecarbonylcobalt complex, hydrogen-octacarbonyldicobalt, hydrogen-hydridecarbonylrhodium, hydrogen-chromium(III) acetylacetonato-triisobutylaluminum, hydrogen-cobalt(II) acetylacetonato-triisobutylaluminum, or

hydrogen-nickel(II)-2-hexanoato-triethylaluminum, or an inhomogeneous system condition such as hydrogen-platinum dioxide, hydrogen-platinum/carbon, hydrogen-palladium/carbon, hydrogen-palladium hydroxide/carbon, 5 hydrogen-palladium/barium sulfate, hydrogen-palladium/calcium carbonate, hydrogen-Raney nickel, hydrogen-copper chromite, hydrogen-rhodium/carbon, hydrogen-rhodium/alumina, hydrogen-ruthenium dioxide, hydrogen-ruthenium/carbon, or hydrogen-iridium black; 10 preferred examples are hydrogen-palladium hydroxide/carbon, hydrogen-iridium black, etc.

The reaction temperature is typically in the range of 0°C - 100°C, preferably 0°C - 60°C. The reaction time which varies with the reaction temperature and the like is 15 typically in the range of 10 minutes - 100 hours, preferably 10 minutes - 96 hours.

Step U"3 is for producing compound (292) and implemented by reacting compound (291) with an acid in an aqueous solvent. The reaction is performed as in the 20 aforementioned step U3 in process U.

Step U"4 is for producing compound (275) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step U"2 in process U".

25 Step U"5 is for producing compound (293) and implemented by reacting compound (291) with an oxidizing agent in an inert solvent.

The inert solvent to be used is not limited in any

particular way as long as it does not interfere with the reaction and examples include ether solvents such as dioxane and tetrahydrofuran, aromatic hydrocarbon solvents such as toluene, halogen-containing solvents such as dichloromethane, as well as dimethylformamide, dimethyl acetamide and acetonitrile; a preferred example is dichloromethane. The oxidizing agent to be used is not limited in any particular way and can be perbenzoic acid, metachloroperbenzoic acid, p-nitroperbenzoic acid, monoperoxyphthalic acid, performic acid, peracetic acid, trifluoroperacetic acid, etc.; a preferred example is metachloroperbenzoic acid. The reaction temperature which varies with the type of solvent and the like is typically in the range of -10°C ~ 50°C , preferably 0°C ~ 30°C . The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 48 hours, preferably 30 minutes - 24 hours.

Step U"6 is for producing compound (294) and implemented by reacting compound (293) with a reducing agent in an inert solvent.

The inert solvent to be used is not limited in any particular way as long as it does not interfere with the reaction; examples are ether solvents such as tetrahydrofuran.

The reducing agent to be used can be sodium/liquid ammonia, lithium/liquid ammonia, lithium/methylamine, lithium/ethylamine, lithium/ethylenediamine, sodium/hexamethylphosphamide-t-butanol, sodium/ethanol,

sodium/t-butanol-tetrahydrofuran, sodium/toluene-t-amyl alcohol, etc.; sodium/liquid ammonia is preferred. The reaction temperature which varies with the type of solvent and the like is typically in the range of $-100^{\circ}\text{C} \sim 20^{\circ}\text{C}$,
5 preferably $-80^{\circ}\text{C} \sim 0^{\circ}\text{C}$. The reaction time which varies with the reaction temperature and the like is typically in the range of 10 minutes - 24 hours, preferably 30 minutes - 5 hours.


Step U"7 is an alternative method for producing
10 compound (273) and implemented by reacting compound (294) with a reducing agent in an inert solvent in the presence of an additive. The reaction is performed as in the aforementioned step E2 in process E.

Process V is for producing compound (296) represented
15 by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}-\text{CH}=\text{G}^3-\text{R}^5$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-
20 position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond; compound (297) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula
25 (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}-\text{CH}=\text{G}^3-\text{S}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line

together with the solid line is a single bond; compound (298) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}=\text{CH}-\text{G}^3-\text{SO}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (299) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}=\text{CH}-\text{G}^3-\text{SO}_2-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (300) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond; and compound (301) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group

and R^1 is $-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}_2-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond.

5 Process V



Step V1 is for producing compound (295) and implemented by reacting compound (108) with a reducing agent in an inert solvent. The reaction is performed as in the aforementioned step U"6 in process U".

Step V2 is for producing compound (296) and implemented by reacting compound (246) with compound (295) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the

aforementined step A4 in process A.

Step V3 is for producing compound (297) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive
5 derivative of compound (137) and reacting it with compound (296) in an alcoholic solvent. The reaction is performed as in the aforementioned step B4 in process B.

Step V4 is for producing compound (298) and implemented by reacting compound (297) with an oxidizing
10 agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.

Step V5 is for producing compound (299) and implemented by reacting compound (297) with an oxidizing agent in an inert solvent. The reaction is performed as in
15 the aforementioned step A9 in process A.

Step V6 is an alternative method for producing compound (299) and implemented by reacting compound (298) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

20 Step V7 is for producing compound (300) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Step V8 is an alternative method for producing
25 compound (301) and implemented by reacting compound (300) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

Step V9 is an alternative method of forming compound


(301) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

In process V, compound (300) can be obtained from
 5 compound (296) if the sequence of reaction steps is $V7 \rightarrow V3 \rightarrow V4$. Compound (301) can also be obtained from compound (300) if the sequence of reaction steps is $V7 \rightarrow V3 \rightarrow V5$.

Process V' is for producing compound (302) represented by the general formula (I) in which X^1 is a hydrogen atom,
 10 X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}=\text{CH}-\text{CH}_2-\text{R}^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the
 15 dashed line together with the solid line is a single bond; compound (303) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group
 20 and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{R}^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (304) represented by the general formula (I) in which X^1 is
 25 a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{COOH}$, R^a is a hydrogen atom, R^b and R^c , when taken

together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; and compound (305) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is a single bond, A is a methylene group and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CON}(\text{R}^7)\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond.

Process V'



Step V'1 is for producing compound (302) and implemented by reacting compound (135) with compound (295) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the aforementioned step A4 in process A.

Step V'2 is for producing compound (303) and implemented by performing catalytic reduction in an

alcoholic solvent or an inert solvent. The reaction is performed as in the aforementioned step A6 in process A.

Step V'3 is for producing compound (304) in the case where R^2 in compound (303) is G^2-COOR^7 and this is
5 implemented by hydrolyzing compound (303) in water or a water-soluble solvent in the presence of a base or an acid (preferably a base). The reaction is performed as in the aforementioned step O6 in process O.

Step V'4 is for producing compound (305) and
10 implemented by reacting compound (304) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

15 Process W is for producing compound (308) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is
20 a methyl group, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a double bond; compound (309) represented by the general formula (I) in which X^1 is a hydrogen atom,
25 X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is a methyl group, R^a is a hydrogen atom, R^b and R^c , when taken

together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; compound (311) represented by the general formula (I) in which X^1 is a hydrogen atom,

5 X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is an allyl group, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they

10 are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; compound (312) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group

15 (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{R}^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; compound (313) represented


20 by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group, A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{R}^2$, R^a is a hydrogen atom, R^b and R^c , when taken together with the

25 carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond; compound (314) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of

α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{SO}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (315) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{SO}_2-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (316) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{COOH}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; and compound (317) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^2-\text{CON}(\text{R}^7)\text{Z}$, R^a is a hydrogen atom, R^b and R^c ,

when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond.

Process W



5

Step W1 is for producing compound (308) and implemented by reacting compound (307) with a metal (preferably magnesium) or an alkylolithium (preferably t-

butyllithium) in an inert solvent to make a reactive derivative of compound (307) and reacting it with compound (306) in an inert solvent in the presence of an additive (preferably tetrakis[(tri-n-butylphosphine)copper(I) iodide])). The reaction is performed as in the
5 aforementioned step O1 in process O.

Step W2 is for producing compound (309) and implemented by reacting compound (308) with a reducing agent in an inert solvent. The reaction is performed as in
10 the aforementioned step U"6 in process U".

Step W3 is for producing compound (310) and implemented by reacting compound (309) with a deprotecting agent in an inert solvent. The reaction is performed as in the aforementioned step U7 in process U.

15 Step W4 is for producing compound (311) and implemented by reacting compound (310) with a base in an inert solvent to make a salt of compound (310) and then reacting it with compound (134) in an inert solvent. The reaction is performed as in the aforementioned step A3 in
20 process A.

Step W5 is for producing compound (312) and implemented by reacting compound (135) with compound (311) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the
25 aforementioned step A4 in process A.

Step W6 is for producing compound (313) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is

performed as in the aforementioned step A6 in process A.

Step W7 is for producing compound (314) in the case where R^2 in compound (313) is G^2-S-Z and this is implemented by reacting compound (313) with an oxidizing agent in an inert solvent. The reaction is performed as in the
5 aforementioned step A8 in process A.

Step W8 is for producing compound (315) in the case where R^2 in compound (313) is G^2-S-Z and this is implemented by reacting compound (313) with an oxidizing agent in an inert solvent. The reaction is performed as in the
10 aforementioned step A9 in process A.


Step W10 is for producing compound (316) in the case where R^2 in compound (313) is G^2-COOR^7 and this is implemented by hydrolyzing compound (313) with a base or an acid (preferably a base) in water or a water-soluble
15 solvent. The reaction is performed as in the aforementioned step O6 in process O.

Step W11 is for producing compound (317) and implemented by reacting compound (316) or reactive derivatives thereof (acid halides, mixed acid anhydrides or
20 active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

Process W' is for producing compound (319) represented
25 by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is

-G⁴-COOR⁷, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O), and the dashed line together with the solid line is a single bond; compound (320) represented
 5 by the general formula (I) in which X¹ is a hydrogen atom, X² is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R¹ is -G⁴-COOH, R^a is a hydrogen atom, R^b and R^c, when taken
 10 together with the carbon atom in 3-position to which they are bound, are -(C=O), and the dashed line together with the solid line is a single bond; and compound (321) represented by the general formula (I) in which X¹ is a
 hydrogen atom and X² is a group of α configuration that is
 15 represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R¹ is -G⁴-CON(R⁷)Z, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-
 position to which they are bound, are -(C=O)-, and the
 20 dashed line together with the solid line is a single bond.

Process W'



Step W'1 is for producing compound (319) and implemented by reacting compound (310) with a base in an inert solvent to make a salt of compound (310) and then reacting it with compound (318) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.


Step W'2 is for producing compound (320) and implemented by hydrolyzing compound (319) with a base or an acid (preferably a base) in water or a water-soluble solvent. The reaction is performed as in the aforementioned step O6 in process O.

Step W'3 is for producing compound (321) and implemented by reacting compound (320) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

Process W" is for producing compound (323) represented by the general formula (I) in which X² is a hydrogen atom,

X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -G⁴-COOR⁷, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; compound (324) represented by the general formula (I) in which X^2 is a hydrogen atom, X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -G⁴-COOR⁷, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond; and compound (325) represented by the general formula (I) in which X^2 is a hydrogen atom and X^1 is a group of β configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is -G⁴-CON(R⁷)Z, R^a is a hydrogen atom, R^b and R^c, when taken together with the carbon atom in 3-position to which they are bound, are -(C=O)-, and the dashed line together with the solid line is a single bond or a double bond.

25 Process W"



Step W''1 is for producing compound (323) and implemented by reacting compound (277) with a base in an inert solvent to make a salt of compound (277) and then reacting it with compound (318) in an inert solvent. The reaction is performed as in the aforementioned step A3 in process A.

Step W''2 is for producing compound (324) and implemented by reacting compound (323) with an acid in an aqueous solvent. The reaction is performed as in the aforementioned step A5 in process A.


Step W''3 is for producing compound (325) and implemented by reacting compound (324) or reactive derivatives thereof (acid halides, mixed acid anhydrides or active esters) with compound (322) or acid addition salts thereof in an inert solvent. The reaction is performed as in the aforementioned step C3 in process C.

Process W' is for producing compound (327) represented by the general formula (I) in which X¹ is a

hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{G}^3-\text{R}^5$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (328) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group, A is -O- and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{G}^3-\text{S}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})-$, and the dashed line together with the solid line is a single bond; compound (329) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}=\text{CH}-\text{G}^3-\text{SO}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(\text{C}=\text{O})$, and the dashed line together with the solid line is a single bond; compound (330) represented by the general formula (I) in which X^1 is a hydrogen atom, X^2 is a group of α configuration represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is -O- and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken

together with the carbon atom in 3-position to which they are bound, are $-(C=O)$, and the dashed line together with the solid line is a single bond; and compound (331) represented by the general formula (I) in which X^1 is a hydrogen atom and X^2 is a group of α configuration that is represented by the general formula (II) in which Ar is an aromatic hydrocarbon group (preferably a p-phenylene group), A is $-O-$ and R^1 is $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{G}^3-\text{SO}_2-\text{Z}$, R^a is a hydrogen atom, R^b and R^c , when taken together with the carbon atom in 3-position to which they are bound, are $-(C=O)-$, and the dashed line together with the solid line is a single bond.

Process W' "



Step W'"1 is for producing compound (327) and implemented by reacting compound (246) with compound (311) in an inert solvent in the presence of an organometallic catalyst. The reaction is performed as in the

aforementioned step A4 in process A.

Step W'"2 is for producing compound (328) and implemented by reacting compound (137) with a metal alkoxide in an alcoholic solvent to make a reactive
 5 derivative of compound (137) and then reacting it with compound (327) in an alcoholic solvent. The reaction is performed as in the aforementioned step B4 in process B.

Step W'"3 is for producing compound (329) and implemented by reacting compound (328) with an oxidizing
 10 agent in an inert solvent. The reaction is performed as in the aforementioned step A8 in process A.

Step W'"4 is for producing compound (330) and implemented by performing catalytic reduction in an alcoholic solvent or an inert solvent. The reaction is
 15 performed as in the aforementioned step A6 in process A.

Step W'"5 is an alternative method of producing compound (331) and implemented by reacting compound (330) with an oxidizing agent in an inert solvent. The reaction is performed as in the aforementioned step A9 in process A.

20 In process W'", compound (330) can also be obtained from compound if the sequence of reaction steps is W'"4 → W'"2 → W'"3. Compound (331) can also be obtained from compound (327) if the sequence of reaction steps is W'"4 → W'"2 → W'"5. Compound (331) can also be obtained from
 25 compound (329) if the sequence of reaction steps is W'"5 → W'"4.

In the above-described processes A - W, B' - L', S' - W', U", W" and W'", if G and/or J and/or Q² is a group

containing a carboxyl group protected by a straight-chained or branched lower alkyl group having 1 - 6 carbon atoms, deprotection can easily be achieved by any known methods of hydrolysis to effect conversion to a carboxyl-containing
5 group.

If any of the steps in the above-described processes A - W, B' - L', S' - W', U", W" and W'" involves groups that need be protected and deprotected, each of them can be protected and deprotected by methods well known to the
10 skilled artisan. For the purposes of protecting and deprotecting, reference can be had, for example, to "Protective Groups in Organic Synthesis", 2nd edition, Theodora W. Green, John Wiley & Sons, Inc., 1991.

Starting material compound (1) is either known or can
15 be easily prepared by known methods or similar methods. [See, for example, J. Med. Chem. 35(11), 2113-2129 (1992); Synth. Commun. 24(16), 2325-2340 (1994); Steroids, 60(5), 414-422 (1995).]

Starting material compound (108) is either known or
20 can be easily prepared by known methods or similar methods. [See, for example, Tetrahedron Letters, 29(13), 1533-1536 (1988).]

Starting material compound (96) is either readily available as a commercial product or can be easily prepared
25 by known methods or similar methods. [See, for example, J. Chem. Res. Miniprint, 2, 0650-0669 (1986).]

Starting material compounds (119) and (144) are readily available as commercial products.

Starting material compounds (133) - (143), compound (183), compound (204), compounds (218) - (220), compound (224), compound (236) and compound (257) are either readily available as commercial products or can be easily prepared
5 by known methods or similar methods.

Starting material compounds (148) and (164) are either known or can be easily prepared by known methods or similar methods. [See, for example, Steroids, 59, 190-195 (1994).]

Starting material compound (223) is either known or
10 can be easily prepared by known methods or similar methods. [See, for example, Synth. Commun. 27(23), 4035-4040 (1997).]

The compounds of the invention which are represented by the general formula (I) and the substances which act as
15 antagonist against but not as agonist for the androgen receptor (which are hereunder also referred to as the test substance) have antiandrogenic activity and other effects and these effects can be measured by the androgen receptor reporter gene assay which has been used in defining the
20 expression "acting as antagonist" and/or the expression "not acting as agonist" for the purposes of the invention as it is optionally combined where appropriate with the following methods of measurement A - F.

Method A: Measurement by in vivo experiment with rats

25 Method A-1: Measuring the antagonist action

If a castrated rat is administered testosterone or dihydrotestosterone, its prostate gland and seminal vesicles increase in weight. By checking to see if the

test substance suppresses the action of testosterone or dihydrotestosterone for increasing the weights of prostate gland and seminal vesicles, one can evaluate the antagonist action of the test substance. For this measurement, reference can be had, for example, to J. Med. Chem., 41:623-639, 1998, and Kiso to Rinsho, 29(4):877-885, 1995.

Method A-2: Measuring the agonist action

A castrated rat is continuously administered the test substance. By checking to see if the weights of the prostate gland and seminal vesicles which are androgen-responsive organs increase after the administration, one can evaluate the agonist action of the test substance. For this measurement, reference can be had, for example, Folia endocrinol., 66:597-606, 1990.

Method B: Measurement based on dimer formation of the androgen receptor

Method B-1: Measuring the action for inhibiting dimer formation

Dihydrotestosterone helps the androgen receptor form a dimer. By applying a gel shift assay to determine if the test substance inhibits the dimer formation of the androgen receptor, one can evaluate the antagonist action of the test substance. For this measurement, reference can be had, for example, to J. Biol. Chem., 268:19004-19012, 1993 and J. Biol. Chem., 270:19998-20003, 1995.

Method B-2: Measuring the action for promoting dimer formation of the androgen receptor

By applying a gel shift assay to determine if the test

substance promotes the dimer formation of the androgen receptor, one can evaluate the agonist action of the test substance. For this measurement, reference can be had, for example, to J. Biol. Chem., 268:19004-19012, 1993 and J. Biol. Chem., 270:19998-20003, 1995.

Method C: Measurement based on ornithine decarboxylase (ODC) activity

By determining whether the test substance elevates or lowers the ODC activity which is believed to reflect androgen-dependent activity, one can evaluate the agonist and antagonist actions of the test substance. For this measurement, reference can be had, for example, to Anal. Biochem., 113:352-355, 1981 and Folia endocrinol., 66:597-606, 1990.

Method D: Measurement based on androgen receptor binding activity

By applying a binding assay to determine whether the test substance inhibits the binding of the androgen receptor to androgen, one can evaluate the antagonist action of the test substance. For this measurement, reference can be had, for example, to Urology, 48:157-163, 1996, J. Biol. Chem., 270:19998-20003, 1995 and Kiso to Rinsho, 29(4):877-885, 1995.

Method E: Measurement based on the increase or decrease in the amount of the androgen receptor

Cells expressing the androgen receptor are treated with the test substance in both the presence and the absence of androgen. By measuring the change in the amount

of the androgen receptor in the cells, one can evaluate the action of the test substance in working as agonist for or antagonist against the androgen receptor. For this measurement, reference can be had, for example, to

5 Endocrinology, 129:2000-2010, 1991.

Method F: Measurement based on nuclear migration of the androgen receptor

Cells exprerssing the androgen receptor are treated with the test substance in the presence or absence of
10 androgen. By applying immunohistostaining to determine the localization of the androgen receptor in the cells, one can check for the nuclear migration of the androgen receptor and determine the action of the test substance for inhibiting the nuclear migration of the androgen receptor,
15 thereby evaluating the action of the test substance as agonist and/or antagonist. For these measurements, reference can be had, for example, to J. Biol. Chem., 267:968-974, 1992.

The compounds of the invention which are represented
20 by the general formula (I) and the substances of the invention which act as antagonist against but not as agonist for the androgen receptor are potential antiandrogenic agents that do exhibit any side effects such as the development of androgen tolerance due to long-term
25 administration and/or hepatotoxicity and, hence, are expected to be useful as pharmaceutical compositions, say, therapeutics for diseases such as prostate cancer, prostatomegaly, male pattern alopecia, sexual prematurity,

acne vulgaris, seborrhea and hirsutism. If the compounds of the invention which are represented by the general formula (I) and the substances of the invention which act as antagonist against but not as agonist for the androgen receptor are preliminarily administered, the onset of diseases such as prostate cancer, prostatomegaly, male pattern alopecia, sexual prematurity, acne vulgaris, seborrhea and hirsutism can hopefully be prevented or retarded, so they are also potential preventives of these diseases.

Pharmaceutical compositions containing as an active ingredient the compounds of the invention which are represented by the general formula (I) and pharmaceutical compositions containing as an active ingredient the substances of the invention which act as antagonist against but not as agonist for the androgen receptor can be administered either orally or parenterally and oral administration is desirable. Prior to administration, such pharmaceutical compositions can be formulated as preparations suitable for the specific method of administration.

Pharmaceutical compositions containing as an active ingredient the compounds of the invention which are represented by the general formula (I) and pharmaceutical compositions containing as an active ingredient the substances of the invention which act as antagonist against but not as agonist for the androgen receptor can be formulated by customary pharmaceutical formulation

techniques and, depending on their use, can be applied as solid and liquid preparations including tablets, capsules, granules, powder, syrup, injection and ointment. Carriers and excipients for such preparations include solid or
5 liquid substances. These may be exemplified by lactose, magnesium stearate, starch, talc, gelatin, agar, pectin, gum arabic, olive oil, sesame oil, ethylene glycol and others in common use.

In these preparations, the pharmaceutical compositions
10 containing as an active ingredient the compounds of the invention which are represented by the general formula (I) and the pharmaceutical compositions containing as an active ingredient the substances of the invention which act as antagonist against but not as agonist for the androgen
15 receptor are incorporated in amounts that vary with their dosage form but it is generally desirable that they be contained at concentrations of 5 - 100 wt%. The pharmaceutical compositions containing as an active ingredient the compounds of the invention which are
20 represented by the general formula (I) and pharmaceutical compositions containing as an active ingredient the substances of the invention which act as antagonist against but not as agonist for the androgen receptor can be adjusted over a broad range depending on the kind of warm-
25 blooded animals including human to be treated, the severity of the disease, doctor's diagnosis, etc. In terms of the active ingredient, the range is from 1 μ g to 500 mg/kg per day, preferably from 20 μ g to 100 mg/kg per day. This dose

can be administered once or several times in one or divided portions per day to month and is variable as appropriate according to the severity of the disease and at doctor's discretion.

5 Examples

Example 1: Evaluating the Agonist Action of Flutamide and Bicaltamide

Twenty-four hours before transfection, 1.0×10^5 HeLa cells were cultured in phenol red free DMEM/5% DCC-FBS on 12-well microplates. Five hundred nanograms/well of MMTV-Luc vector, 100 ng/well of pSG5-hAR and 5 ng/well of Renilla Luc vector were transfected into the HeLa cells. The transfection was performed in a liquid culture of the phenol red free DMEM using 3 mL/well of lipofectoamine. Nine hours after the transfection, the liquid culture was replaced by phenol red free DMEM/3% DCC-FBS containing 10 mmol/L of hydroxyflutamide or bicaltamide. The transcriptional activity value was measured 48 hours after the replacement of the liquid culture. Transcriptional activity was measured with a dual-luciferase reporter assay system. The transcriptional activity value was calculated as the value for firefly luciferase divided by the value for sea pansy luciferase. Hydroxyflutamide and bicaltamide exhibited more than five times the value for the case of non-addition and, hence, the agonist action of hydroxyflutamide and bicaltamide was verified (Table 1).

<Table 1>

<u>Luciferase Activity (fold induction)¹⁾</u>	
Not added	1.00
10 μ mol/L of hydroxyflutamide	7.84 (> 5.0)
5 <u>10 μmol/L of bicaltamide</u>	<u>7.62 (> 5.0)</u>

1) The value with the luciferase activity value for
"not added" being taken as 1.00.

10 Example 2: Evaluating the Antagonist Action of Flutamide
and Bicaltamide

Twenty-four hours before transfection, 1.0×10^5 HeLa cells were cultured in phenol red free DMEM/5% DCC-FBS on 12-well microplates. Five hundred nanograms/well of MMTV-Luc vector, 100 ng/well of pSG5/hAR and 5 ng/well of Renilla Luc vector were transfected into the HeLa cells. The transfection was performed in a liquid culture of the phenol red free DMEM using 3 mL/well of lipofectoamine. Nine hours after the transfection, the liquid culture was replaced by phenol red free DMEM/3% DCC-FBS containing 0.1 nmol/L of DHT and 1.0 mmol/L of hydroxyflutamide or bicaltamide. The transcriptional activity value was measured 48 hours after the replacement of the liquid culture. Transcriptional activity was measured with a dual-luciferase reporter assay system. The transcriptional activity value was calculated as the value for firefly luciferase divided by the value for sea pansy luciferase. Hydroxyflutamide and bicaltamide lowered the transcriptional activity value of DHT to less than 50% and,

hence, the antagonist action of hydroxyflutamide and bicaltamide was verified (Table 2).

<Table 2>

5	<u>Luciferase Activity (relative activity)²⁾</u>
	0.1 nmol/L of DHT 100
	1.0 μ mol/L of hydroxyflutamide 29.0 (< 50.0)
	<u>1.0 μmol/L of bicaltamide</u> 32.0 (< 50.0)
10	2) The value with the luciferase activity value of 0.1 nmol/L of DHT being taken as 100.

Example 3: Synthesis of 17 β -hydroxy-7 α -(7-carboxyheptyl)-5 α -androstan-3-one

(Step 1)

15 17 β -t-butyldimethylsilyloxy-7 α -(2-propen-1-yl)-5 α -androstan-3-one

20 Metallic lithium (220 mg) was added to liquid ammonia (150 ml) at -78 °C. After 5-minute stirring, 17 β -t-butyldimethylsilyloxy-7 α -(2-propen-1-yl)-4-androsten-3-one (1.261 g) and a tetrahydrofuran solution (20 ml) of t-butanol (0.41 ml) were added and the mixture was stirred for 20 minutes. After adding 1,2-dibromoethane (3 ml) and ammonium chloride (30 g), the mixture was stirred at 25°C for 30 minutes. After adding water, extraction with ethyl acetate was conducted. The organic layer was dried with magnesium sulfate and, after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl

acetate/n-hexane = 1/10) gave the end compound in 810.7 mg (yield, 64%).

¹H-NMR(270MHz, CDCl₃)δ: 0.01(6H, s), 0.73(3H, s), 0.88(9H, s), 1.04(3H, s), 0.92-2.45(23H, m), 3.55(1H, t, J=8.3Hz),
5 4.93(1H, d, J=3.8Hz), 4.99(1H, s), 5.58-5.72(1H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/10): 0.54

(Step 2)

17β-t-butyldimethylsilyloxy-7α-(7-methoxycarbonyl-2-hepten-
10 1-yl)-5α-androstan-3-one

17β-t-Butyldimethylsilyloxy-7α-(2-propen-1-yl)-5α-androstan-3-one (596.1 mg) was dissolved in dichloromethane (5 ml) and, after adding methyl 6-heptenoate (384.4 mg) and benzylidenebis(tricyclohexylphosphine)-dichlororuthenium
15 (57.0 mg), the mixture was heated under reflux for 5 hours in an argon atmosphere. After standing to cool, purification was effected by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/10) to give the end compound in 527.6 mg (yield,
20 70%).

¹H-NMR(270MHz, CDCl₃)δ: 0.01(6H, s), 0.71(3H, s), 0.88(9H, s), 1.03(3H, s), 0.90-2.10(26H, m), 2.18-2.43(5H, m), 3.51(1H, t, J=8.4Hz), 3.67(3H, s), 5.18-5.40(2H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.43
25

(Step 3)

17β-hydroxy-7α-(7-carboxyheptyl)-5α-androstan-3-one

17β-t-Butyldimethylsilyloxy-7α-(7-methoxycarbonyl-2-


hepten-1-yl)-5 α -androstan-3-one (505.5 mg) was dissolved in ethyl acetate (30 ml) and, after adding 10% palladium/carbon (148 mg), the mixture was stirred for 4 hours at 25°C in a hydrogen atmosphere. The reaction mixture was filtered and the solvent was distilled off at reduced pressure; the resulting residue was dissolved in acetone (10 ml) and, after adding 1 N-HCl (1 ml), the mixture was heated under reflux for 26 hours. After standing to cool, water was added and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2 ~ 1/1) gave the end compound in 362.6 mg (yield, 93%).

¹H-NMR(270MHz, CDCl₃) δ : 0.76(3H, s), 1.04(3H, s), 1.00-1.82(27H, m), 1.98-2.15(3H, m), 2.23-2.48(5H, m), 3.65(1H, t, J=8.7Hz).

Mass(FAB): 433(M+1).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.27

The following compounds were synthesized by similar methods to Example 3.



Example	n	Z	MW (Molecular weight)	Mass
4	4	H	404	405 (FAB)
5	8	H	460	461 (FAB)
6	10	H	488	489 (FAB)
7	12	H	516	517 (FAB)
8	8	$-(\text{CH}_2)_3\text{CF}_2\text{CF}_3$	620	621 (ESI)

[Example 9]

5 Synthesis of 17 β -hydroxy-7 α -(7-(N,N-dimethylaminocarbonyl)-heptyl)-5 α -androstan-3-one

The 17 β -hydroxy-7 α -(7-carboxyheptyl)-5 α -androstan-3-one (9.9 mg) obtained in Example 3 was dissolved in tetrahydrofuran (0.5 ml) and, after adding 1-(N,N-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (13.0 mg), 1-hydroxybenzotriazole monohydrate (10.5 mg) and a solution (68.7 μ l) of 2.0 M-dimethylamine in tetrahydrofuran, the mixture was stirred for 15 hours at 25 °C. After adding ethyl acetate (2.0 ml), the mixture was washed with 1 N-hydrochloric acid, a saturated aqueous solution of sodium hydrogencarbonate and a saturated aqueous solution of sodium chloride. After drying with magnesium sulfate, the mixture was filtered through NH silica gel (Pro. No. DM1020; product of Fuji Silicia Chemical Co., Ltd.) and the solvent was distilled off at reduced pressure to give the end compound in 10.5 mg (99.7%).

¹H-NMR(270MHz, CDCl₃) δ : 0.76(3H, s), 1.04(3H, s), 1.00-1.83(27H, m), 1.95-2.16(3H, m), 2.23-2.47(5H, m), 2.94(3H,


s), 3.01(3H, s), 3.65(1H, t, J=8.7Hz).

Mass(ESI): 460(M+1).

Rf value (on silica gel plate, developing solvents:
methanol/chloroform = 1/1): 0.28

5


The following compounds were synthesized by similar methods to Example 9.



Example	n	RM	RN	MW	Mass
10	7	H	Et	459	460(FAB)
11	7	H	cyclohexylmethyl	527	528(FAB)
12	7	H	cyclopropylmethyl	485	486(FAB)
13	7	H	n-Bu	487	488(ESI)
14	7	H	i-Pr	473	474(FAB)
15	7	H	t-Bu	487	488(FAB)
16	7	H	c-hexyl	513	514(ESI)
17	7	H	-(CH ₂) ₃ OH	489	490(ESI)
18	7	Me	n-Bu	501	502(ESI)
19	7	Et	Et	487	488(ESI)
20	7		-(CH ₂) ₅ -	499	500(ESI)
21	7	H	4-t-butylbenzyl	577	578(ESI)
22	7	H	-CH ₂ CHPh ₂	611	612(ESI)
23	7	H	2-furylmethyl	511	512(ESI)
24	7	H	Me	445	446(ESI)
25	7	Me	Et	473	474(ESI)
26	7	Me	n-Pr	487	488(ESI)
27	7	Me	i-Pr	487	488(FAB)
28	7	Me	Bn	535	536(ESI)

29	7	-(CH ₂) ₄ -		485	486(ESI)
30	7	-CH ₂ CH ₂ OCH ₂ CH ₂ -		501	502(ESI)
31	7	Me	t-Bu	501	502(ESI)
32	7	H	cyclopropyl	471	472(ESI)
33	6	Me	Me	445	446(FAB)
34	6	Et	Et	473	474(FAB)
35	6	-(CH ₂) ₅ -		485	486(FAB)
36	8	Me	Me	473	474(ESI)
37	8	Et	Et	501	524(ESI)
38	8	Me	n-Bu	515	538(ESI)
39	8	H	Bn	535	558(ESI)
40	8	H	-(CH ₂) ₂ OH	489	512(ESI)
41	8	-(CH ₂) ₅ -		513	514(ESI)
42	9	Me	Me	487	488(ESI)
43	9	Et	Et	515	516(ESI)
44	9	-(CH ₂) ₄ -		513	514(ESI)
45	9	Me	Et	501	502(ESI)
46	9	Me	n-Bu	529	530(ESI)
47	9	H	Bn	549	550(ESI)
48	9	-(CH ₂) ₅ -		527	528(ESI)
49	9	H	-(CH ₂) ₂ OH	503	504(ESI)
50	9	Me	n-Pr	515	516(ESI)
51	9	-CH ₂ CH ₂ OCH ₂ CH ₂ -		529	530(ESI)
52	10	Me	Me	501	502(FAB)
53	10	Et	Et	529	530(FAB)
54	10	Me	Et	515	516(FAB)
55	10	Me	n-Pr	529	530(FAB)
56	10	Me	n-Bu	543	544(FAB)
57	10	-CH ₂ CH ₂ OCH ₂ CH ₂ -		543	544(FAB)
58	11	Me	Me	515	516(FAB)
59	11	Et	Et	543	544(FAB)
60	11	-(CH ₂) ₅ -		555	556(FAB)
61	11	H	Bn	577	578(FAB)
62	11	Me	n-Bu	557	558(FAB)
63	11	H	-(CH ₂) ₂ OH	531	532(FAB)

[Example 64]



Synthesis of 17β-hydroxy-7α-[7-{N-(2-hydroxyethyl)-aminocarbonyl}heptyl]-5α-androstan-3-one

5 The 17β-hydroxy-7α-(7-carboxyheptyl)-5α-androstan-3-one (10.3 mg) obtained in Example 3 and O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluromium hexafluorophosphate (30 mg) were dissolved in tetrahydrofuran (1 ml) and, after adding N,N-

10 diisopropylethylamine (25 μl) and 2-aminoethanol (4.4 μl), the mixture was stirred for 2 hours at 25°C. After adding ethyl acetate, the reaction mixture was washed with a saturated aqueous solution of sodium hydrogencarbonate, 1 N-hydrochloric acid and a saturated aqueous solution of

15 sodium chloride. To the organic layer, NH silica gel (Pro. No. DM1020; product of Fuji Silicia Chemical Co., Ltd.) was added and the mixture was stirred for 5 minutes; after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel

20 column chromatography (developing solvents: methanol/chloroform = 1/10) to give the end compound in 7.5 mg (yield, 66%).


¹H-NMR(270MHz, CDCl₃)δ: 0.76(3H, s), 1.04(3H, s), 0.95-1.82(28H, m), 1.95-2.10(3H, m), 2.20(2H, t, J=7.4Hz), 2.28-

2.45(2H, m), 3.43(2H, q, J=5.2Hz), 3.60-3.78(3H, m),
5.98(1H, brs).

Mass(ESI): 476(M+1).

Rf value (on silica gel plate, developing solvents:
5 methanol/chloroform = 1/10): 0.12


The following compounds were synthesized by similar
methods to Example 64.



Example	n	R ^M	R ^N	MW	Mass
65	7	H	n-Pr	473	474(FAB)
66	7	H	n-hexyl	515	516(ESI)
67	7	H	i-pentyl	501	502(FAB)
68	7	H	i-Bu	487	488(FAB)
69	7	H	neopentyl	501	502(ESI)
70	7	H	3-pentyl	501	502(ESI)
71	7	n-hexyl	n-hexyl	599	600(ESI)
72	7	H	Ph	507	508(ESI)
73	7	H	Bn	521	522(ESI)
74	7	H	-CH ₂ CH ₂ Ph	535	536(ESI)

10 [Example 75]

Synthesis of 17 β -hydroxy-11 β -(9-carboxynonyl)-5 α -androstan-
3-one
(Step 1)




3,3-ethylenedioxy-17β-(methoxymethoxy)-5α-androstan-11-one

3,3-Ethylenedioxy-17β-hydroxy-5α-androstan-11-one
(1.84 g) was dissolved in dichloromethane (30 ml) and,
5 after adding N,N-diisopropylethylamine (2.7 ml) and
chloromethylmethyl ether (1.2 ml) dropwise, the mixture was
stirred for 8 hours at 25°C. The reaction mixture was
poured into a saturated aqueous solution of ammonium
chloride and subjected to extraction with dichloromethane.
10 The organic layer was washed with a saturated aqueous
solution of sodium chloride and dried with magnesium
sulfate; after filtering, the solvent was distilled off at
reduced pressure. The resulting residue was purified by
silica gel column chromatography (developing solvents:
15 ethyl acetate/n-hexane = 1/1) to give the end compound in
1.98 g (yield, 95%).

¹H-NMR(270MHz, CDCl₃)δ: 0.72(3H, s), 1.03(3H, s), 1.03-
1.38(7H, m), 1.52-1.80(9H, m), 2.05-2.23(2H, m) 2.36-
2.48(2H, m), 3.33(3H, s), 3.70(1H, t, J=8.4Hz), 3.92(4H, s),
20 4.57(1H, d, J=14.2Hz), 4.60(1H, d, J=14.2Hz).

R_f value (on silica gel plate, developing solvents: ethyl
acetate/n-hexane = 1/1): 0.61

(Step 2)



3,3-ethylenedioxy-17β-methoxymethoxy-11-
[[(1,1,2,2,3,3,4,4,4-nonafluorobutyl)sulfonyl]oxy]-5α-
androst-11-ene

5 To a solution of lithium diisopropylamide (as prepared from diisopropylamine (0.15 ml) and n-butyllithium (1.5 M hexane solution) (0.69 ml)) in tetrahydrofuran (1.3 ml), a solution of 3,3-ethylenedioxy-17β-methoxymethoxy-5α-androstan-11-one (100 mg) in tetrahydrofuran (1.3 ml) was
10 added dropwise over 5 minutes. After stirring for 30 minutes at -78°C, perfluorobutanesulfonyl fluoride (0.13 ml) was added dropwise over 5 minutes. After stirring for 5 minutes at -78°C, the mixture was stirred for 2 hours at room temperature. A saturated aqueous solution of ammonium
15 chloride was added to the reaction mixture and extraction with ethyl acetate was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure.


20 Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/6) gave the end compound in 98.8 mg (yield, 57%).

¹H-NMR(270MHz, CDCl₃)δ: 0.93(3H, s), 0.96(3H, s), 0.83-2.23(18H, m), 3.34(3H, s), 3.70(1H, t, J=8.2Hz), 3.93(4H, s), 4.58(1H, d, J=6.6Hz), 4.63(1H, d, J=6.6Hz), 6.20(1H, s).

25

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/2): 0.67

(Step 3)



5 3,3-ethylenedioxy-17β-(methoxymethoxy)-11-(2-propen-1-yl)-
5α-androst-11-ene


3,3-Ethylenedioxy-17β-methoxymethoxy-11-
 [{{(1,1,2,2,3,3,4,4,4-nonafluorobutyl)sulfonyl}oxy]-5α-
 androst-11-ene) (454.5 mg) was dissolved in tetrahydrofuran
 10 (6 ml) and, after adding allyltributyltin (299.1 mg),
 lithium chloride (90.0 mg) and
 tetrakis(triphenylphosphine)palladium (47.7 mg), the
 mixture was heated under reflux for 22 hours in an argon
 atmosphere. After standing to cool, an aqueous solution of
 15 potassium fluoride was added and extraction with ethyl
 acetate was effected. The organic layer was washed with
 water and a saturated aqueous solution of sodium chloride
 and dried with magnesium sulfate; after filtering, the
 solvent was distilled off at reduced pressure. The
 20 resulting residue was purified by silica gel column
 chromatography (developing solvents: ethyl acetate/n-
 hexane = 1/9) to give the end compound in 275.8 mg (yield,
 98%).

¹H-NMR(270MHz, CDCl₃)δ: 0.83(3H, s), 0.89(3H, s), 0.93-

1.88(16H, m), 1.98-2.12(2H, m), 2.72-2.92(2H, m), 3.37(3H, s), 3.59(1H, t, J=8.7Hz), 3.94(4H, s), 4.62(1H, d, J=10.1Hz), 4.65(1H, d, J=10.1Hz), 4.91-5.02(2H, m), 5.65-5.82(1H, m), 5.88(1H, s).

5 Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.53

(Step 4)



10 3,3-ethylenedioxy-17β-(methoxymethoxy)-11-(9-methoxycarbonyl-2-nonen-1-yl)-5α-androst-11-ene


3,3-Ethylenedioxy-17β-(methoxymethoxy)-11-(2-propen-1-yl)-5α-androst-11-ene (248.5 mg) was dissolved in dichloromethane (3 ml) and, after adding methyl 8-nonenoate (202.9 mg) and benzylidenebis(tricyclohexylphosphine)-
15 dichlororuthenium (27.3 mg), the mixture was heated under reflux for 5 hours in an argon atmosphere. After standing to cool, purification was effected by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/6) to give the end compound in 227.8 mg (yield,
20 68%).

¹H-NMR(270MHz, CDCl₃)δ: 0.83(3H, s), 0.88(3H, s), 0.90-1.40(13H, m), 1.40-1.83(11H, m), 1.90-2.12(4H, m), 2.30(2H, t, J=7.6Hz), 2.65-2.84(2H, m), 3.36(3H, s), 3.60(1H, t, J=8.4Hz), 3.66(3H, s), 3.93(4H, s), 4.63(1H, d, J=11.2Hz),

4.65(1H, d, J=11.2Hz), 5.23-5.40(2H, m), 5.87(1H, s).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/6): 0.23

(Step 5)



5


3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonylnonyl)-5α-androstane

3,3-Ethylenedioxy-17β-(methoxymethoxy)-11-(9-methoxycarbonyl-2-nonen-1-yl)-5α-androst-11-ene (226.3 mg) was dissolved in ethyl acetate (5 ml) and, after adding iridium black (55.7 mg), the mixture was stirred for 5 days at 25°C in a hydrogen atmosphere. The reaction mixture was filtered through Celite and concentrated at reduced pressure; the resulting residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/10 - 1/6) to give the end compound in 165.9 mg (yield, 73%).

¹H-NMR(270MHz, CDCl₃)δ: 0.83(3H, s), 0.93(3H, s), 0.95-1.05(3H, m), 1.10-1.41(20H, m), 1.45-1.80(11H, m), 1.90-2.07(2H, m), 2.16(1H, d, J=12.5Hz), 2.30(2H, t, J=7.6Hz), 3.35(3H, s), 3.43(1H, t, J=8.7Hz), 3.66(3H, s), 3.93(4H, s), 4.62(2H, s).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.47

(Step 6) 32-54



17β-hydroxy-11β-(9-carboxynonyl)-5α-androstan-3-one


3,3-Ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonylnonyl)-5α-androstane (91.0 mg) was dissolved in acetone (3 ml) and, after adding 1 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux for 24 hours. After standing to cool, water was added and extraction with ethyl acetate was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/chloroform = 1/6 - 1/2) to give the end compound in 70.0 mg (yield, 94%).

¹H-NMR(270MHz, CDCl₃)δ: 0.85(3H, s), 0.90-1.03(4H, m), 1.15(3H, s), 1.16-1.88(25H, m), 1.98-2.48(10H, m), 3.58(1H, t, J=8.7Hz).

Mass(FAB): 461(M+1).


R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/2): 0.086

The following compounds were synthesized by similar methods to Example 75.




Example	n	Z	MW	Mass
76	6	H	432	433 (FAB)
77	7	H	446	447 (FAB)
78	10	H	488	489 (FAB)
79	8	$-(CH_2)_3CF_2CF_3$	620	621 (ESI)

[Example 80]



5 MW 634, Mass(ESI): 635(M+1).

[Example 81]



10 Synthesis of 17 β -hydroxy-11 β -(9-aminocarbonylnonyl)-5 α -androstan-3-one

The 17 β -hydroxy-11 β -(9-carboxynonyl)-5 α -androstan-3-one (16.6 mg) obtained in Example 75 was dissolved in tetrahydrofuran (1 ml) and, after adding triethylamine (6.0 μ l) and ethyl chlorocarbonate (4.0 μ l) at -10°C, the mixture was stirred for 10 minutes. Ammonia gas was blown

into the reaction mixture for 5 minutes and the mixture was stirred for 20 minutes at -10°C . A saturated aqueous solution of ammonium chloride was added to the reaction mixture, which was then reverted to room temperature.


- 5 After extraction with ethyl acetate, the organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography (developing solvents: methanol/chloroform = 1/20) to give the end compound in 15.5 mg (yield, 94%).

$^1\text{H-NMR}$ (270MHz, CDCl_3) δ : 0.85(3H, s), 0.88-1.05(4H, m), 1.15(3H, s), 1.10-1.88(25H, m), 1.97-2.50(10H, m), 3.58(1H, t, $J=8.7\text{Hz}$), 5.34(2H, brs).

- 15 Mass(FAB): 460(M+1)


Rf value (on silica gel plate, developing solvents: methanol/chloroform = 1/10): 0.33

- 20 The following compounds were synthesized by similar methods to Example 81.




Example	n	Z	MW	Mass
82	9	n-phenyl	529	530(FAB)
83	11	H	487	488(FAB)
84	11	n-phenyl	557	558(FAB)

The following compounds were synthesized by similar methods to Example 9.



Example	n	R ^M	R ^N	MW	Mass
85	7	Me	Me	459	460 (FAB)
86	7	H	Me	445	446 (FAB)
87	7	Me	Et	473	474 (FAB)
88	7	Me	n-Pr	487	488 (FAB)
89	7	-CH ₂ CH ₂ OCH ₂ CH ₂ -		501	502 (FAB)
90	8	Me	Me	473	474 (FAB)
91	8	H	Me	459	460 (FAB)
92	8	Me	Et	487	488 (ESI)
93	8	Me	n-Pr	501	502 (FAB)
94	8	-CH ₂ CH ₂ OCH ₂ CH ₂ -		515	516 (FAB)
95	9	Me	Me	487	488 (ESI)
96	9	Et	Et	515	516 (ESI)
97	9	-(CH ₂) ₅ -		527	528 (ESI)
98	9	H	Bn	549	550 (ESI)
99	9	Me	n-Bu	529	530 (ESI)
100	9	H	-CH ₂ CH ₂ OH	503	504 (ESI)


[Example 101]



5

Synthesis of 17β-hydroxy-11β-(9-carboxynonyloxy)-5α-androstan-3-one

(Step 1)




3,3-ethylenedioxy-11β-hydroxy-17β-(methoxymethoxy)-5α-androstane

3,3-Ethylenedioxy-17β-(methoxymethoxy)-5α-androstan-
 5 11-one (2.84 g) was dissolved in diethyl ether (500 ml) and, after adding lithium aluminum hydride (548 mg), the mixture was heated under reflux for 2 hours in an argon atmosphere. The reaction mixture was cooled to 0°C and, after adding water, filtered through Celite. After extraction with
 10 ethyl acetate, the organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography
 15 (developing solvents: ethyl acetate/n-hexane = 1/2) to give the end compound in 2.68 g (yield, 94%).

¹H-NMR(270MHz, CDCl₃)δ: 0.75-0.99(3H, m), 1.01(3H, s), 1.06(3H, s), 1.20-1.92(16H, m), 1.94-2.07(2H, m), 3.35(3H, s), 3.49(1H, t, J=8.3Hz), 3.94(4H, s), 4.29-4.36(1H, m),
 20 4.61(2H, s).

R_f value (on silica gel plate, developing solvents: methanol/chloroform = 1/50): 0.31

(Step 2)




3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(2-propen-1-yloxy)-5α-androstane

In an argon atmosphere, 3,3-ethylenedioxy-11β-hydroxy-
 5 17β-(methoxymethoxy)-5α-androstane (953.9 mg) was dissolved
 in N,N-dimethylformamide (10 ml) and, after adding sodium
 hydride (60% in oil) (486.7 mg), the mixture was stirred
 for 3 hours at 50°C. After reversion to 25°C, allyl bromide
 (2.20 ml) and tetra-n-butylammonium iodide (208.5 mg) were
 10 added and the mixture was stirred for 3 hours at 50°C. The
 reaction mixture was cooled to 0°C and water was added.
 After extraction with ethyl acetate, the organic layer was
 washed with a saturated aqueous solution of sodium chloride
 and dried with magnesium sulfate; after filtering, the
 15 solvent was distilled off at reduced pressure. The
 resulting residue was purified by silica gel column
 chromatography (developing solvents: ethyl acetate/n-
 hexane = 1/5) to give the end compound in 684.5 mg (yield,
 65%).

20 ¹H-NMR(270MHz, CDCl₃)δ: 0.74-0.93(3H, m), 0.95(3H, s),
 1.03(3H, s), 1.18-2.09(16H, m), 2.32(1H, dd, J=2.9, 14.4Hz),
 3.36(3H, s), 3.47(1H, t, J=8.3Hz), 3.70(1H, dd, J=7.2,
 16.2Hz), 3.77-3.83(1H, m), 3.93(4H, s), 4.10(1H, dd, J=7.2,
 16.2Hz), 4.63(2H, AB-q), 5.08(1H, split-d, J=10.6Hz),
 25 5.25(1H, dd, J=1.7, 17.2Hz), 5.83-6.00(1H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/2): 0.59

(Step 3)




5 3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonyl-2-nonen-1-yloxy)-5α-androstane

3,3-Ethylenedioxy-17β-(methoxymethoxy)-11β-(2-propen-1-yloxy)-5α-androstane (18.9 mg) was dissolved in dichloromethane (0.5 ml) and, after adding methyl 8-
 10 nonenoate (14.8 mg) and benzylidenebis(tricyclohexylphosphine)-dichlororuthenium (2.0 mg), the mixture was heated under reflux for 4 hours in an argon atmosphere. After standing to cool and concentrating at reduced pressure, purification was
 15 effected by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/5) to give the end compound in 15.5 mg (yield, 62%).

¹H-NMR(270MHz, CDCl₃)δ: 0.70-0.95(3H, m), 0.95(3H, s), 1.02(3H, s), 1.15-1.86(23H, m), 1.93-2.08(3H, m), 2.30(3H, t, J=7.6Hz), 3.36(3H, s), 3.46(1H, t, J=8.7Hz), 3.64(1H, dd, J=5.0, 11.3Hz), 3.67(3H, s), 3.74-3.80(1H, m), 3.93(4H, s), 4.02(1H, dd, J=5.0, 11.3Hz), 4.63(2H, AB-q), 5.44-5.69(1H, m).

Rf value (on silica gel plate, developing solvents: ethyl
 25 acetate/n-hexane = 1/4): 0.26

(Step 4)



3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonylnonyloxy)-5α-androstane


5 3,3-Ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonyl-2-nonen-1-yloxy)-5α-androstane (17.2 mg) was dissolved in ethyl acetate (3 ml) and, after adding 10%-palladium/carbon (6.5 mg), the mixture was stirred for 1 hour at 25 °C in a hydrogen atmosphere. The reaction mixture was filtered and the solvent was distilled off at reduced pressure to give the residue in 16.7 mg. In a separate run, 3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonyl-2-nonen-1-yloxy)-5α-androstane (32.1 mg) was dissolved in ethyl acetate (6 ml) and, after adding 10%-palladium/carbon (6.5 mg), the solution was stirred for 1 hour at 25°C in a hydrogen atmosphere. The reaction mixture was filtered and the solvent was distilled off at reduced pressure to give the residue in 30.1 mg. The two residues were combined and purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2) to give the end compound in 44.7 mg (yield, 90%).

1H-NMR(270MHz, CDCl₃)δ: 0.71-0.94(3H, m), 0.94(3H, s), 1.02(3H, s), 1.20-2.07(30H, m), 2.30(3H, t, J=7.6Hz), 3.07(3H, dt, J=5.7, 8.4Hz), 3.36(3H, s), 3.43-3.58(2H, m),

3.67(3H, s), 3.68-3.74(1H, m), 3.93(4H, s), 4.63(2H, AB-q).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/2): 0.55

(Step 5)



5

17β-hydroxy-11β-(9-carboxynonyloxy)-5α-androstan-3-one

3,3-Ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonylnonyloxy)-5α-androstane (21.0 mg) was dissolved in acetone (2 ml) and, after adding 1 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux for 10 hours. After adding water to the reaction mixture, extraction with dichloromethane was effected and the organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure to give the residue in 20.6 mg. In a separate run, 3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(9-methoxycarbonylnonyloxy)-5α-androstane (22.0 mg) was dissolved in acetone (2 ml) and, after adding 1 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux for 10 hours. After adding water to the reaction mixture, extraction with dichloromethane was effected and the organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at


reduced pressure to give the residue in 21.6 mg. The two residues were combined and purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/1) to give the end compound in 29.3 mg (yield, 70%).

$^1\text{H-NMR}$ (270MHz, CDCl_3) δ : 0.74(1H, dd, $J=3.3, 10.9\text{Hz}$), 0.85-0.98(3H, m), 0.94(3H, s), 1.24(3H, s), 1.26-1.72(22H, m), 1.79-2.14(5H, m), 2.25-2.57(6H, m), 3.10(1H, dt, $J=6.1, 8.8\text{Hz}$), 3.53-3.63(2H, m), 3.71-3.78(1H, m).

Mass(FAB): 477(M+1).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.12

The following compounds were synthesized by similar methods to Example 101.




Example	n	Z	MW	Mass(FAB)
102	4	H	420	421
103	6	H	448	449
104	9	H	490	491
105	10	H	504	505
106	12	H	532	533
107	22	H	672	673
108	8	$-(\text{CH}_2)_3\text{CF}_2\text{CF}_3$	636	637

[Example 109]

Synthesis of 17 β -hydroxy-11 β -(10-(4,4,5,5,5-

pentafluoropentylsulfanyl)decyl)-5 α -androstan-3-one



The 3,3-diethylenedioxy-17 β -(methoxymethoxy)-11 β -(9-methoxycarbonylnonyl)-5 α -androstane (64.8 mg) obtained in
 5 step 5 of Example 75 was dissolved in tetrahydrofuran (3 ml) and, after adding lithium borohydride (11 mg), the mixture was stirred for 4 hours at 25°C. To the reaction mixture, lithium triethylborohydride (1.0 M-tetrahydrofuran solution, 100 μ l) was added and the mixture was stirred for
 10 4 hours at 25°C. To the reaction mixture, lithium borohydride (20 mg) was added and the mixture was stirred for 15 hours at 25°C. After adding water, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and
 15 dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4 - 1/2) to give an oil in 63.8 mg. This oil was dissolved in
 20 dichloromethane (2 ml) and, after adding triethylamine (30 μ l) and methanesulfonyl chloride (15 μ l) at 0°C, the mixture was stirred for 4 hours at 25°C. The reaction mixture was poured into a saturated aqueous solution of sodium chloride and extraction with dichloromethane was
 25 conducted. The organic layer was dried with magnesium

sulfate and after filtering, the solvent was distilled off at reduced pressure. The resulting residue was dissolved in acetone (3 ml) and, after adding sodium iodide (300 mg), the mixture was stirred for 15 hours at 25°C. A saturated aqueous solution of sodium sulfite was added to the reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) to give an oil in 53.9 mg. This oil and 1-(acetylsulfanyl)-4,4,5,5,5-pentafluoropentane (45 mg) were dissolved in methanol (1 ml) and tetrahydrofuran (0.5 ml) and, after adding a methanol solution (0.18 ml) of 1 N-sodium methoxide, the mixture was stirred for 15 hours at 25°C. After adding water to the reaction mixture, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) to give an oil in 66.3 mg. This oil was dissolved in acetone (2 ml) and, after adding 1 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux for 36 hours. After standing to cool, water was added and extraction with ethyl acetate was

conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The residue was
 5 purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) to give the end compound in 52.9 mg (yield, 74%).


¹H-NMR(270MHz, CDCl₃)δ: 0.85(3H, s), 0.89-1.09(3H, m), 1.15(3H, s), 1.18-1.74(28H, m), 1.80-2.45(13H, m), 2.51(2H, t, J=7.4Hz), 2.59(2H, t, J=6.9Hz), 3.51-3.62(1H, m).

Mass(FAB): 623(M+1).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.19

15 [Example 110]

Synthesis of 17β-hydroxy-11β-(10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl)-5α-androstan-3-one



The 17β-hydroxy-11β-(10-(4,4,5,5,5-pentafluoropentylsulfinyl)decyl)-5α-androstan-3-one (20.5 mg) obtained
 20 in Example 109 was dissolved in tetrahydrofuran (0.8 ml) and, after adding OXONE (registered trademark, 12.4 mg) and water (0.4 ml), the mixture was stirred for 2 hours at 0°C. A saturated aqueous solution of sodium hydrogencarbonate

was added to the reaction mixture and, after reverting it to room temperature, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2 ~ 1/1 ~ 2/1) to give the end compound in 19.5 mg (yield, 93%).

¹H-NMR(270MHz, CDCl₃)δ: 0.85(3H, s), 0.86-1.05(3H, m), 1.15(3H, s), 1.10-1.90(27H, m), 1.95-2.50(13H, m), 2.60-2.82(4H, m), 3.57(1H, t, J=8.2Hz).


Mass(FAB): 639(M+1).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.056

[Example 111]

Synthesis of 17β-hydroxy-7α-(7-hydroxyheptyl)-5α-androstan-3-one

(Step 1)



17β-t-butyldimethylsilyloxy-7α-(7-hydroxy-2-hepten-1-yl)-5α-androstan-3-one


17β-t-Butyldimethylsilyloxy-7α-(2-propen-1-yl)-5α-androstan-3-one (33.4 mg) was dissolved in dichloromethane

(0.5 ml) and, after adding 5-hexen-1-ol (20.0 mg) and benzylidenebis(tricyclohexylphosphine)-dichlororuthenium (5.2 mg), the mixture was heated under reflux for 2 hours in an argon atmosphere. After standing to cool, purification was effected by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) to give the end compound in 19.7 mg (yield, 51%).

¹H-NMR(270MHz, CDCl₃)δ: 0.01(6H, s), 0.72(3H, s), 0.88(9H, s), 1.04(3H, s), 0.90-2.10(27H, m), 2.20-2.47(3H, m), 3.54(1H, t, J=8.6Hz), 3.63(2H, t, J=6.8Hz), 5.19-5.42(2H, m).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.13

(Step 2)



17β-hydroxy-7α-(7-hydroxyheptyl)-5α-androstan-3-one

17β-t-Butyldimethylsilyloxy-7α-(7-hydroxy-2-hepten-1-yl)-5α-androstan-3-one (19.6 mg) was dissolved in ethyl acetate (10 ml) and, after adding 10%-palladium/carbon (6.3 mg), the mixture was stirred for 1 hour at 25°C in a hydrogen atmosphere. The reaction mixture was filtered and the solvent was distilled off at reduced pressure; the resulting residue was dissolved in acetone (2 ml) and,

after adding 2 N-hydrochloric acid (0.5 ml), the mixture was stirred for 2 hours at 25°C. After adding water to the reaction mixture, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2 - 1/1) to give the end compound in 15.0 mg (yield, 98%).

¹H-NMR(270MHz, CDCl₃)δ: 0.76(3H, s), 1.04(3H, s), 0.80-1.83(29H, m), 1.95-2.15(3H, m), 2.22-2.47(3H, m), 3.60-3.70(3H, m).


Mass(ESI): 405(M+1).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.24

The following compounds were synthesized by similar methods to Example 111.

Example	n	MW	Mass (ESI)
112	8	418	419
113	9	432	433

[Example 114]



Synthesis of 17 β -hydroxy-7 α -(7-carbamoylheptyl)-5 α -androstan-3-one


The 17 β -hydroxy-7 α -(7-carboxyheptyl)-5 α -androstan-3-one (12.6 mg) obtained in Example 3 was dissolved in tetrahydrofuran (0.5 ml) and then triethylamine (8.1 μ l) and ethyl chloroformate (4.2 μ l) were added dropwise at 0°C. After stirring for 5 minutes, ammonia gas was bubbled for 30 seconds. After stirring for 30 minutes, water was added to the reaction mixture and extraction with dichloromethane was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: dichloromethane/methanol = 20/1) gave the end compound in 11.6 mg (92%).

¹H-NMR(270MHz, CDCl₃) δ : 0.76(3H, s), 1.04(3H, s), 1.00-1.83(27H, m), 1.94-2.16(3H, m), 2.20-2.50(5H, m), 3.65(1H, t, J=8.4Hz), 5.34-5.54(2H, m).

Mass(FAB): 432(M+1).


Rf value (on silica gel plate, developing solvents: dichloromethane/methanol = 20/1): 0.48

The following compounds were synthesized by similar methods to Example 114.




Example	n	R ^N	MW	Mass (FAB)
115	5	H	403	404
116	5	n-pentyl	473	474
117	7	n-pentyl	501	502
118	9	H	459	460
119	9	n-pentyl	529	529
120	11	H	487	487
121	11	n-pentyl	557	557
122	13	H	515	515
123	13	n-pentyl	585	585

[Example 124]



5 17β-hydroxy-7α-{13-(4,4,5,5,5-pentafluoropentylsulfinyl)-tridecyl}-5α-androstan-3-one
(Step 1)




Synthesis of 17β-hydroxy-7α-(2-propen-1-yl)-5α-androstan-3-one

10 17β-t-Butyldimethylsilyloxy-7α-(2-propen-1-yl)-5α-androstan-3-one (170 mg) was dissolved in acetone (2 ml) and then 2 N-hydrochloric acid (0.5 ml) was added dropwise. After stirring for 4 hours at 25°C, water was added to the

reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was
 5 distilled off at reduced pressure to give the end compound in 126 mg (100%).

¹H-NMR(270MHz, CDCl₃)δ: 0.77(3H, s), 1.04(3H, s), 0.96-2.44(23H, m), 3.60-3.70(1H, m), 4.94(1H, d, J=3.5Hz), 5.00(1H, s), 5.58-5.72(1H, m).

10 R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.54
 (Step 2)



15 Synthesis of 3,3-ethylenedioxy-17β-hydroxy-7α-(2-propen-1-yl)-5α-androstane


To a benzene solution (5 ml) of 17β-hydroxy-7α-(2-propen-1-yl)-5α-androstan-3-one (126 mg), ethylene glycol (2 ml) and p-toluenesulfonic acid (13.2 mg) were added and the mixture was heated under reflux with water being
 20 continuously removed with a Dean-Stark trap. After two hours, a saturated aqueous solution of sodium hydrogencarbonate was added under cooling with ice and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of

sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure to give the end compound in 141 mg (yield, 99%).

¹H-NMR(270MHz, CDCl₃)δ: 0.74(3H, s), 0.85(3H, s), 0.92-
5 2.20(23H, m), 3.56-3.70(1H, m), 3.93(4H, s), 4.92-5.04(2H, m), 5.62-5.80(1H, m).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.60

(Step 3)



Synthesis of 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(2-propen-1-yl)-5α-androstane


To a dichloromethane solution (4 ml) of 3,3-ethylenedioxy-17β-hydroxy-7α-(2-propen-2-yl)-5α-androstane
15 (141 mg), diisopropylethylamine (0.227 ml) and chloromethyl methyl ether (0.087 ml) were added dropwise under cooling with ice. After stirring for 14 hours at 25°C, a saturated aqueous solution of sodium hydrogencarbonate was added to the reaction mixture and extraction with dichloromethane
20 was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl

acetate/n-hexane = 1/4) gave the end compound in 131 mg (yield, 83%).

¹H-NMR(270MHz, CDCl₃)δ: 0.77(3H, s), 0.84(3H, s), 0.92-2.20(23H, m), 3.35(3H, s), 3.53(1H, t, J=8.3Hz), 3.92(4H, s), 4.62(2H, d, J=1.8Hz), 4.92-5.04(2H, m), 5.62-5.80(1H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.50

(Step 4)



Synthesis of 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(13-bromo-2-tridecen-1-yl)-5α-androstane


3,3-Ethylenedioxy-17β-methoxymethoxy-7α-(2-propen-1-yl)-5α-androstane (42.6 mg) was dissolved in dichloromethane (1.5 ml) and, after adding 12-bromododecene (50.4 mg) and benzyldienebis(tricyclohexylphosphine)-dichlororuthenium (8.4 mg), the mixture was heated under reflux for 5 hours in an argon atmosphere. After standing to cool, purification was performed by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/10) to give the end compound in 56.0 mg (yield, 86%).

¹H-NMR(270MHz, CDCl₃)δ: 0.76(3H, s), 0.83(3H, s), 0.94-

2.14(4H, m), 3.34(3H, s), 3.41(2H, t, $J=6.9\text{Hz}$), 3.52(1H, t, $J=8.3\text{Hz}$), 3.92(4H, s), 4.62(2H, d, $J=1.8\text{Hz}$), 5.22-5.46(2H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.56

(Step 5)




Synthesis of 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(13-bromotridecyl)-5α-androstane

3,3-Ethylenedioxy-17β-methoxymethoxy-7α-(13-bromo-2-tridecen-1-yl)-5α-androstane (55.3 mg) was dissolved in ethyl acetate (2 ml) and, after adding 10%-palladium/carbon (10 mg), the mixture was stirred for 13 hours at 25 °C in a hydrogen atmosphere. After filtering the reaction mixture, the solvent was distilled off at reduced pressure to give the end compound in 47.4 mg (86%).

$^1\text{H-NMR}$ (270MHz, CDCl_3) δ : 0.76(3H, s), 0.84(3H, s), 0.94-2.10(45H, m), 3.34(3H, s), 3.41(2H, t, $J=6.9\text{Hz}$), 3.53(1H, t, $J=8.3\text{Hz}$), 3.93(4H, s), 4.62(2H, d, $J=1.8\text{Hz}$).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.56

(Step 6)



Synthesis of 17β-hydroxy-7α-((13-(4,4,5,5,5-pentafluoro-
pentylsulfanyl)tridecyl))-5α-androstan-3-one


5 4,4,5,5,5-Pentafluoropentanethioacetate (35.0 mg) was dissolved in methanol (1 ml) and then 1 M sodium methylate/methanol solution (0.12 ml) was added dropwise. After stirring for 30 minutes, a solution of 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(13-bromotridecyl)-5α-
 10 androstane (47.4 mg) in tetrahydrofuran (1 ml) was added to the reaction mixture. After stirring for 18 hours, water was added to the reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried
 15 with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Crude purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/10) gave 3,3-ethylenedioxy-17β-methoxymethoxy-7α-((13-(4,4,5,5,5-
 20 pentafluoropentylsulfanyl)-tridecyl)androstane (42.6 mg), which was then dissolved in acetone (2 ml); after adding 2 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux for 3 hours at 60°C. After standing to cool down to 0°C, water was added and extraction with chloroform was
 25 conducted. The organic layer was washed with a saturated

aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound in 40.2 mg (82%).

¹H-NMR(270MHz, CDCl₃)δ: 0.76(3H, s), 1.04(3H, s), 0.88-2.40(49H, m), 2.50(2H, t, J=7.3Hz), 2.59(2H, t, J=7.9Hz), 3.58-3.70(1H, m).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.32
(Step 7)

Synthesis of 17β-hydroxy-7α-{13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyl}-5α-androstan-3-one



15

17β-Hydroxy-7α-{13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyl}-5α-androstan-3-one (26.0 mg) was dissolved in tetrahydrofuran (1 ml) and then OXONE (14.4 mg) and water (0.2 ml) were added at 0°C. After stirring for 30 minutes, a saturated aqueous solution of sodium hydrogencarbonate was added to the reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after

filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 2/1) gave the end compound in 19.5 mg (73%).


5 $^1\text{H-NMR}$ (270MHz, CDCl_3) δ : 0.76(3H, s), 1.04(3H, s), 0.98-2.40(49H, m), 2.58-2.82(4H, m), 3.60-3.70(1H, m).

Mass(FAB): 681(M+1).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.10

10


The following compounds were synthesized by similar methods to Example 124.



Example	n	MW	Mass (FAB)
125	5	568	569
126	7	596	597
127	9	624	625
128	11	652	653

15 [Example 129]

Synthesis of 17 β -hydroxy-7 α -(13-(4,4,5,5,5-pentafluoropentylsulfonyl)tridecyl)-5 α -androstan-3-one




17 β -Hydroxy-7 α -(13-(4,4,5,5,5-pentafluoropentyl-sulfanyl)tridecyl)-5 α -androstan-3-one (15.0 mg) was dissolved in tetrahydrofuran (1 ml) and then OXONE (27.8 mg) and water (0.2 ml) were added at 25°C. After stirring
 5 for 1 hour, a saturated aqueous solution of sodium hydrogencarbonate was added to the reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after
 10 filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2) gave the end compound in 15.0 mg (95%).

1H-NMR(270MHz, CDCl₃) δ : 0.76(3H, s), 1.04(3H, s), 0.98-
 15 2.40(49H, m), 2.92-3.08(4H, m), 3.60-3.70(1H, m).

Mass (FAB): 697(M+1).


Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.32

The following compounds were synthesized by similar
 20 methods to Example 129.




Example	N	MW	Mass (FAB)
130	7	612	613
131	9	640	641
132	11	668	669

[Example 133]



Synthesis of 17β-hydroxy-7α-(4-(3-carboxypropoxy)phenyl)-5α-androstan-3-one

5 (Step 1)



Synthesis of 17β-hydroxy-7-(4-methoxyphenyl)-5α-androst-4-en-3-one

In an argon atmosphere, copper(I) iodide (1.14 g) was
 10 dissolved in anhydrous tetrahydrofuran (5 ml) and then 0.5
 M 4-methoxyphenylmagnesium bromide/tetrahydrofuran solution
 (11.9 ml) was added dropwise at -50 °C. After stirring
 for 10 minutes, 17β-t-butyldimethylsilyloxyandrost-4,6-
 dien-3-one (600 mg), chlorotrimethylsilane (0.376 ml) and a
 15 tetrahydrofuran solution (6 ml) of hexamethylphosphoric
 triamide (0.518 ml) were added dropwise at -78°C. The
 temperature of the mixture was elevated to -40°C over 1
 hour. To the reaction mixture, 2 N-hydrochloric acid was
 added and, after stirring for 1 hour at 25°C, extraction
 20 with ethyl acetate was conducted. The organic layer was


dried with magnesium sulfate and, after filtering, the solvent was distilled off at reduced pressure.

Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2) gave the end compound in 67.9 mg (yield, 12%) as a diastereomeric mixture.

¹H-NMR(270MHz, CDCl₃)δ: 0.55-0.99(4/3H, m), 0.76(2H, s), 0.81(1H, s), 1.00-2.54(47/3H, m), 1.32(2H, s), 1.34(1H, s), 2.82-3.00(2/3H, m), 3.00-3.08(1/3H, m), 3.40-3.56(1H, m), 3.78(1H, s), 3.80(2H, s), 5.71(1H, s), 6.74-6.86(2H, m), 7.04-7.18(2H, m).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.38

(Step 2)



Synthesis of 17β-hydroxy-7-(4-methoxyphenyl)-5α-androstan-3-one

Metallic lithium (11.9 mg) was added to liquid ammonia (15 ml) at -78°C. After stirring for 5 minutes, 17β-hydroxy-7-(4-methoxyphenyl)-5α-androst-4-en-3-one (67.8 mg) and a solution of t-butanol (25.3 μl) in tetrahydrofuran (3 ml) were added and the mixture was stirred for 5 minutes. After adding 1,2-dibromoethane (0.1 ml) and ammonium chloride (1 g), the mixture was stirred


for 30 minutes at 25°C. After adding water, extraction with ethyl acetate was conducted. The organic layer was dried with magnesium sulfate and, after filtering, the solvent was distilled off at reduced pressure.

- 5 Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2) gave the end compound in 49.8 mg (yield, 73%) as a diastereomeric mixture.

1H-NMR(270MHz, CDCl₃)δ: 0.50-0.62(2/3H, m), 0.73(2H, s),
 10 0.78(1H, s), 0.84-1.00(2/3H, m), 1.11(1H, m), 1.15(2H, m),
 1.04-2.44(58/3H, m), 2.90-3.00(1/3H, m), 3.42-3.58(1H, m),
 3.78(2H, s), 3.79(1H, s), 6.70-7.30(4H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.48

- 15 (Step 3)



Synthesis of 3,3-ethylenedioxy-17β-hydroxy-7-(4-methoxyphenyl)-5α-androstane


- 17β-Hydroxy-7-(4-methoxyphenyl)-5α-androstan-3-one
 20 (49.7 mg) was dissolved in benzene (2 ml) and, after adding ethylene glycol (0.5 ml) and p-toluenesulfonic acid (2.2 mg), the mixture was heated under reflux as water was continuously removed by means of a Dean-Stark trap. After one hour, the reaction mixture was cooled to 0°C and, after

adding a saturated aqueous solution of sodium hydrogencarbonate, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure to give the end compound in 55.0 mg (yield, 100%) as a diastereomeric mixture.

¹H-NMR(270MHz, CDCl₃)δ: 0.54-0.60(2/3H, m), 0.70(2H, s), 0.75(1H, s), 0.91(1H, s), 0.95(2H, s), 0.90-1.98(58/3H, m), 2.22-2.38(2/3H, m), 2.86-2.94(1/3H, m), 3.44-3.58(1H, m), 3.72-3.98(4H, m), 3.78(2H, s), 3.80(1H, s), 6.70-7.30(4H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.52

(Step 4)



Synthesis of 3,3-ethylenedioxy-17β-methoxymethoxy-7-(4-methoxyphenyl)-5α-androstane


To a solution of 3,3-ethylenedioxy-17β-hydroxy-7-(4-methoxyphenyl)-5α-androstane (55.0 mg) in dichloromethane (2 ml), diisopropylethylamine (0.128 ml) and chloromethyl methyl ether (0.047 ml) were added dropwise at 0°C. After stirring for 12 hours at 25°C, a saturated aqueous solution of sodium hydrogencarbonate was added to the reaction

mixture and extraction with dichloromethane was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound in 56.2 mg (yield, 93%) as a diastereomeric mixture.

¹H-NMR(270MHz, CDCl₃)δ: 0.42-0.60(2/3H, m), 0.74(2H, s), 0.78(1H, s), 0.80-0.90(2/3H, m), 0.91(1H, s), 0.94(2H, s), 1.00-1.96(56/3H, m), 2.22-2.36(2/3H, m), 2.84-2.92(1/3H, m), 3.30(3H, s), 3.34-3.44(1H, m), 3.78(2H, s), 3.80(1H, s), 3.82-3.98(4H, m), 4.56(2H, s), 6.70-7.30(4H, m).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.68

(Step 5)



Synthesis of 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(4-hydroxyphenyl)-5α-androstane and 3,3-ethylenedioxy-17β-methoxymethoxy-7β-(4-hydroxyphenyl)-5α-androstane

To a solution of 3,3-ethylenedioxy-17β-methoxymethoxy-7-(4-methoxyphenyl)-5α-androstane (151 mg) in N,N-dimethylacetamide (3 ml), sodium thiomethylate (109 mg) was added and the mixture was heated under reflux. After 3

hours, the reaction mixture was cooled to 0°C and, after adding water, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound as both 7 α form in 42.3 mg (yield, 29%) and 7 β form in 88.0 mg (yield, 60%).

3,3-ethylenedioxy-17 β -methoxymethoxy-7 α -(4-hydroxyphenyl)-5 α -androstane;

¹H-NMR(270MHz, CDCl₃) δ : 0.78(3H, s), 0.90(3H, s), 1.00-2.08(20H, m), 2.84-2.92(1H, m), 3.30(3H, s), 3.38(1H, t, J=8.6Hz), 3.80-3.94(4H, m), 4.56(2H, s), 4.64(1H, s), 6.72(2H, d, J=8.4Hz), 7.23(2H, d, J=8.4Hz).


Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.20

3,3-ethylenedioxy-17 β -methoxymethoxy-7 β -(4-hydroxyphenyl)-5 α -androstane;

¹H-NMR(270MHz, CDCl₃) δ : 0.46-0.60(1H, m), 0.73(3H, s), 0.94(3H, s), 0.82-1.96(19H, m), 2.22-2.34(1H, m), 3.30(3H, s), 3.38(1H, t, J=8.6Hz), 3.93(4H, brs), 4.56(2H, s), 4.68(1H, s), 6.66(1H, dd, J=2.3, 8.2Hz), 6.74(1H, dd, J=2.3, 8.2Hz), 6.92(1H, dd, J=1.8, 8.2Hz), 7.07(1H, dd, J=1.8, 8.2Hz).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.28

(Step 6)



Synthesis of 3,3-ethylenedioxy-17 β -methoxymethoxy-7 α -{4-(3-t-butoxycarboxypropoxy)phenyl}-5 α -androstane

5 To an N,N-dimethylacetamide solution (0.5 ml) of 3,3-ethylenedioxy-17 β -methoxymethoxy-7 α -(4-hydroxyphenyl)-5 α -androstane (22.0 mg), t-butyl 4-bromobutyrate (31.3 mg), potassium carbonate (64.5 mg) and 18-crown-6 (123 mg) were added. After one hour, water was added to the reaction


10 mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column

15 chromatography (developing solvents: ethyl acetate/n-hexane = 1/8) gave the end compound in 27.8 mg (yield, 99%).

¹H-NMR(270MHz, CDCl₃) δ : 0.78(3H, s), 0.90(3H, s), 1.04-2.12(22H, m), 1.46(9H, s), 2.43(2H, t, J=7.3Hz), 2.82-2.92(1H, m), 3.30(3H, s), 3.38(1H, t, J=8.4Hz), 3.80-3.94(4H, m), 3.97(2H, t, J=6.1Hz), 4.56(2H, s), 6.78(2H, d, J=8.6Hz), 7.26(2H, d, J=8.6Hz).

R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/4): 0.40

(Step 7)



Synthesis of 17β-hydroxy-7α-{4-(3-carboxypropoxy)phenyl}-5α-androstan-3-one


5 3,3-Ethylenedioxy-17β-methoxymethoxy-7α-(4-(3-methoxycarbonylpropoxy)phenyl)-5α-androstane (27.6 mg) was dissolved in acetone (2 ml) and, after adding 2 N-hydrochloric acid (0.5 ml), the mixture was heated at 60°C. After two hours, water was added to the reaction mixture and extraction with dichloromethane was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: dichloromethane/methanol = 10/1) gave the end compound in 19.2 mg (yield, 89%).

1H-NMR(270MHz, CDCl₃)δ: 0.78(3H, s), 1.10(3H, s), 1.00-2.44(22H, m), 2.60(2H, t, J=7.3Hz), 2.88-2.96(1H, m), 3.50(1H, t, J=8.2Hz), 4.00(2H, t, J=5.9Hz), 6.77(2H, d, J=8.6Hz), 7.20(2H, d, J=8.6Hz).

Mass (FAB): 469(M+1).


R_f value (on silica gel plate, developing solvent: ethyl acetate): 0.54

The following compounds were synthesized by similar methods to Example 133.



Example	n	MW	Mass (FAB)
134	1	440	441
135	7	524	525

[Example 136]



Synthesis of 17β-hydroxy-7α-{4-(3-carbamoylpropoxy)phenyl}-5α-androstan-3-one

The 17β-hydroxy-7α-{4-(3-carboxypropoxy)phenyl}-5α-androstan-3-one obtained in Example 133 was dissolved in tetrahydrofuran (0.5 ml) and then triethylamine (3.2 μl) and ethyl chloroformate (1.8 μl) were added dropwise under cooling with ice. After stirring for 5 minutes, ammonia gas was bubbled for 1 minute. After stirring for 15 minutes, water was added to the reaction mixture and extraction with dichloromethane was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced


pressure. Purification by preparative chromatography (developing solvents: dichloromethane/methanol = 20/1) gave the end compound in 6.6 mg (yield, 90%).

¹H-NMR(270MHz, CDCl₃)δ: 0.78(3H, s), 1.11(3H, s), 1.00-2.44(22H, m), 2.45(2H, t, J=7.1Hz), 2.88-2.98(1H, m), 3.50(1H, t, J=8.2Hz), 4.01(2H, t, J=5.7Hz), 5.30-5.60(2H, m), 6.77(2H, d, J=8.6Hz), 7.20(2H, d, J=8.6Hz).

Mass (FAB): 468(M+1).


Rf value (on silica gel plate, developing solvents: dichloromethane/methanol = 20/1): 0.14

The following compounds were synthesized by similar methods to Example 136.



Example	n	MW	Mass (FAB)
137	1	439	440
138	7	523	524


15 [Example 139]



Synthesis of 17β-hydroxy-7α-4-{3-(N-pentylcarbamoyl)propoxy}phenyl)-5α-androstan-3-one


- The 17 β -hydroxy-7 α -(4-(3-carboxypropoxy)phenyl)-5 α -androstan-3-one (7.0 mg) obtained in Example 133 was dissolved in tetrahydrofuran (0.5 ml) and then 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (8.6 mg),
- 5 1-hydroxybenzotriazole monohydrate (6.8 mg) and pentylamine (10.4 ml) were added at 25°C. After stirring for 4 hours, a saturated aqueous solution of sodium hydrogencarbonate was added to the reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a
- 10 saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by preparative chromatography (developing solvent: ethyl acetate) gave the end compound in 5.8 mg (yield, 72%).
- 15 ¹H-NMR(270MHz, CDCl₃) δ : 0.78(3H, s), 0.80-0.94(3H, m), 1.11(3H, s), 1.00-2.44(30H, m), 2.88-2.98(1H, m), 3.24(2H, dt, J=6.1, 7.1Hz), 3.50(1H, t, J=8.3Hz), 3.99(2H, t, J=5.8Hz), 5.50(1H, brs), 6.77(2H, d, J=8.6Hz), 7.20(2H, d, J=8.6Hz).
- 20 Mass (FAB): 538(M+1).
- Rf value (on silica gel plate, developing solvent: ethyl acetate): 0.62

The following compound was synthesized by a similar method to Example 139.



Example	n	MW	Mass (FAB)
140	7	593	594


[Example 141]



Synthesis of 17β-hydroxy-7α-(4-methoxyphenyl)-5α-androstan-

5 3-one

(Step 1)




Synthesis of 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(4-
methoxyphenyl)-5α-androstane

10 The 3,3-ethylenedioxy-17β-methoxymethoxy-7α-(4-
hydroxyphenyl)-5α-androstane (10.0 mg) obtained in step 5
of Example 136 was dissolved in N,N-dimethylformamide (1
ml) and then 60% sodium hydride (2.5 mg) and iodomethane
(13.2 μl) were added at 0°C. After stirring for 13 hours
15 at 25°C, a saturated aqueous solution of NH₄Cl was added to
the reaction mixture and extraction with ethyl acetate was
conducted. The organic layer was washed with a saturated
aqueous solution of sodium chloride and dried with
magnesium sulfate; after filtering, the solvent was

distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound in 10.2 mg (yield, 99%).

5 $^1\text{H-NMR}$ (270MHz, CDCl_3) δ : 0.78(3H, s), 0.91(3H, s), 1.00-1.96(20H, m), 2.84-2.92(1H, m), 3.30(3H, s), 3.38(1H, t, $J=8.6\text{Hz}$), 3.80(3H, s), 3.82-3.92(4H, m), 4.56(2H, s), 6.80(2H, d, $J=8.6\text{Hz}$), 7.27(2H, d, $J=8.6\text{Hz}$).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1:1): 0.68
10 (Step 2)



Synthesis of 17β-hydroxy-7α-(4-methoxyphenyl)androstane-3-one

15 3,3-Ethylenedioxy-17β-methoxymethoxy-7α-(4-methoxyphenyl)androstane (10.2 mg) was dissolved in acetone (2 ml) and, after adding 2 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux. After 2 hours, water was added to the reaction mixture and extraction with
20 dichloromethane was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography

(developing solvents: ethyl acetate/n-hexane = 1/1) gave the end compound in 7.2 mg (yield, 85%).


¹H-NMR(270MHz, CDCl₃)δ: 0.78(3H, s), 1.11(3H, s), 1.00-2.50(20H, m), 2.88-2.96(1H, m), 3.50(1H, t, J=8.6Hz),
5 3.79(3H, s), 6.79(2H, d, J=8.7Hz), 7.22(2H, d, J=8.7Hz).

Mass (FAB): 397(M+1).

Rf value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 1/1): 0.48


[Example 142]

10 Synthesis of 17β-hydroxy-11β-(13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyloxy)-5α-androstan-3-one



(Step 1)

15 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(13-bromo-2-tridecenyloxy)-5α-androstane



In an argon atmosphere, the 3,3-ethylenedioxy-17β-(methoxymethoxy)-11β-(2-propen-1-yloxy)-5α-androstane (100.8 mg) obtained in step 2 of Example 104, 12-bromododecan-1-ene (114.7 mg) and


benzylidenebistricyclohexylphosphine dichlororuthenium (19.1 mg) were dissolved in toluene (2.3 ml) and the mixture was stirred for 26 hours at 110 °C. The solvent was distilled off at reduced pressure and purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/5) gave the end compound in 98.6 mg (yield, 65%).

¹H-NMR(270MHz, CDCl₃)δ: 0.95(3H, s), 1.02(3H, s), 0.70-2.38(38H, m), 3.36(3H, s), 3.93(4H, s), 3.31-4.12(6H, m), 4.56-4.67(2H, m), 5.38-5.69(2H, m).

R_f value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/4): 0.49

(Step 2)

3,3-ethylenedioxy-17β-methoxymethoxy-11β-(13-(4,4,5,5,5-pentafluoropentylthio)-2-tridecenyloxy)-5α-androstane



In a nitrogen atmosphere, 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(13-bromo-2-tridecenyloxy)-5α-androstane (98.6 mg), 4,4,5,5,5-pentafluoropentane-1-thiol acetate (71.2 mg) and sodium methylate (1.0 M methanol solution) (0.30 ml) were dissolved in methanol (1.5 ml) and tetrahydrofuran (0.8 ml) and the mixture was stirred for 11 hours at room temperature. After adding water to the reaction mixture, extraction with ethyl acetate was


conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; then, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound in 97.3 mg (yield, 84%).

1H-NMR(270MHz, CDCl₃) δ : 0.95(3H, s), 1.02(3H, s), 0.69-2.38(42H, m), 2.50(2H, t, J=7.3Hz), 2.59(2H, t, J=6.9Hz), 3.36(3H, s), 3.47(1H, t, J=8.2Hz), 3.59-4.17(3H, m), 3.93(4H, s), 4.57-4.68(2H, m), 5.39-5.70(2H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/4): 0.42

(Step 3)

3,3-ethylenedioxy-17 β -methoxymethoxy-11 β -{13-(4,4,5,5,5-pentafluoropentylsulfinyl)-2-tridecenyloxy}-5 α -androstande



3,3-Ethylenedioxy-17 β -methoxymethoxy-11 β -{13-(4,4,5,5,5-pentafluoropentylthio)-2-tridecenyloxy}-5 α -androstande (50.1 mg) was dissolved in tetrahydrofuran (0.6 ml) and, after adding oxone (20.1 mg) and water (0.3 ml) under cooling with ice, the mixture was stirred for 1 hour. After adding a saturated aqueous solution of sodium hydrogencarbonate to the reaction mixture, extraction with ethyl acetate was conducted. The organic layer was washed


with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; then, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 3/2) gave the end compound in 29.4 mg (yield, 57%).

¹H-NMR(270MHz, CDCl₃)δ: 0.95(3H, s), 1.02(3H, s), 0.70-2.37(42H, m), 2.58-2.82(4H, m), 3.36(3H, s), 3.46(1H, t, J=8.2Hz), 3.93(4H, s), 3.59-4.17(3H, m), 4.57-4.68(2H, m), 5.35-5.70(2H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/2): 0.12

(Step 4)

17β-hydroxy-11β-{13-(4,4,5,5,5-pentafluoropentylsulfinyl)tridecyloxy}-5α-androstan-3-one




3,3-Ethylenedioxy-17β-methoxymethoxy-11β-{13-(4,4,5,5,5-pentafluoropentylsulfinyl)-2-tridecenyloxy}-5α-androstane (29.4 mg) was dissolved in ethyl acetate (1 ml) and, after adding 10% palladium/carbon (10 mg), the mixture was stirred for 3 hours at room temperature in a hydrogen atmosphere. After filtering the reaction mixture, the solvent was distilled off at reduced pressure and the resulting residue was dissolved in acetone (2 ml); after

adding 1 N-hydrochloric acid (1 ml), the mixture was heated under reflux for 3 hours. After standing to cool, a saturated aqueous solution of sodium hydrogencarbonate was added to the reaction mixture and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 2/1) gave the end compound in 20.5 mg (yield, 78%).

¹H-NMR(270MHz, CDCl₃)δ: 0.94(3H, s), 1.24(3H, s), 0.68-2.54(47H, m), 2.58-2.83(4H, m), 3.04-3.16(1H, m), 3.51-3.63(2H, m), 3.72-3.79(1H, m).


Mass (FAB): 697(M+1).

The following compounds were synthesized by similar methods to Example 142.




Example	n	MW	Mass (FAB)
143	5	584	585
144	7	612	613
145	9	640	641
146	11	668	669

[Example 147]



Synthesis of 17β-hydroxy-11β-(13-(4,4,5,5,5-
pentafluoropentylsulfonyl)tridecyloxy)-5α-androstan-3-one

5 (Step 1)



3,3-ethylenedioxy-17β-methoxymethoxy-11β-(13-(4,4,5,5,5-
pentafluoropentylsulfonyl)-2-tridecenyloxy)-5α-androstane


3,3-Ethylenedioxy-17β-methoxymethoxy-11β-(13-
 10 (4,4,5,5,5-pentafluoropentylthio)-2-tridecenyloxy)-5α-
 androstane (47.2 mg) was dissolved in tetrahydrofuran (0.6
 ml) and, after adding OXONE (75.7 mg) and water (0.3 ml) at
 room temperature, the mixture was stirred for 1 hour.
 After adding a saturated aqueous solution of sodium
 15 hydrogencarbonate to the reaction mixture, extraction with
 ethyl acetate was conducted. The organic layer was washed
 with a saturated aqueous solution of sodium chloride and
 dried with magnesium sulfate; then, the solvent was
 distilled off at reduced pressure. Purification by silica
 20 gel column chromatography (developing solvents: ethyl
 acetate/n-hexane = 1/3) gave the end compound in 33.6 mg

(yield, 68%).

¹H-NMR(270MHz, CDCl₃)δ: 0.95(3H, s), 1.02(3H, s), 0.70-2.39(42H, m), 2.92-3.12(4H, m), 3.36(3H, s), 3.46(1H, t, J=8.1Hz), 3.93(4H, s), 3.59-4.18(3H, m), 4.57-4.68(2H, m),
5 5.37-5.70(2H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/2): 0.47

(Step 2)



10 17β-hydroxy-11β-{13-(4,4,5,5,5-
pentafluoropentylsulfonyl)-2-tridecenyloxy}-5α-androstan-3-one
3,3-Ethylenedioxy-17β-methoxymethoxy-11β-{13-
(4,4,5,5,5-pentafluoropentylsulfonyl)-2-tridecenyloxy}-5α-
androstande (33.6 mg) was dissolved in ethyl acetate (1 ml)
15 and, after adding 10% palladium/carbon (10 mg), the mixture
was stirred for 4 hours at room temperature in a hydrogen
atmosphere. After filtering the reaction mixture, the
solvent was distilled off at reduced pressure and the
resulting residue was dissolved in acetone (2 ml); after
20 adding 1 N-hydrochloric acid (1 ml), the mixture was heated
under reflux for 4 hours. After standing to cool, a
saturated aqueous solution of sodium hydrogencarbonate was
added and extraction with ethyl acetate was effected. The
organic layer was washed with a saturated aqueous solution


of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 2/3) gave the end compound in 22.9 mg (yield, 76%).

$^1\text{H-NMR}$ (270MHz, CDCl_3) δ : 0.94(3H, s), 1.24(3H, s), 0.68-2.54(47H, m), 2.92-3.15(5H, m), 3.51-3.64(2H, m), 3.72-3.78(1H, m).

Mass (FAB): 713(M+1).

10


The following compounds were synthesized by similar methods to Example 147.



Example	n	MW	Mass (FAB)
148	7	628	629
149	9	656	657
150	11	684	685


15

[Example 151]



17 β -hydroxy-11 β -[4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyloxy}phenyl]-5 α -androstan-3-one

(Step 1)



5


3,3-ethylenedioxy-17 β -methoxymethoxy-11-(4-methoxyphenyl)-5 α -androst-11-ene

A mixture of 3,3-ethylenedioxy-17 β -methoxymethoxy-11-
 10 [(1,1,2,2,3,3,4,4,4-nonafluorobutyl)sulfonyl}oxy]-5 α -
 androst-11-ene (98.8 mg), 4-methoxyphenylboronic acid (223
 mg), tetrakis(triphenylphosphine) palladium (6.8 mg), lithium
 chloride (12.4 mg), 2 M aqueous solution of sodium
 carbonate (0.5 ml), toluene (2 ml) and ethanol (1 ml) was
 heated under reflux for 13 hours in an argon atmosphere.
 15 After adding a saturated aqueous solution of sodium
 hydrogencarbonate to the reaction mixture, extraction with
 ethyl acetate was conducted. The organic layer was washed
 with a saturated aqueous solution of sodium chloride and
 dried with magnesium sulfate; after filtering, the solvent
 20 was distilled off at reduced pressure. Purification by
 silica gel column chromatography (developing solvents:
 ethyl acetate/n-hexane = 1/6) gave the end compound in 65.4
 mg (yield, 93%).

¹H-NMR(270MHz, CDCl₃) δ : 0.59-2.28(18H, m), 0.85(3H, s),

0.94(3H, s), 3.31(3H, s), 3.63(1H, t, $J=8.0\text{Hz}$), 3.78(3H, s), 3.80-3.96(4H, m), 4.57(1H, d, $J=6.6\text{Hz}$), 4.61(1H, d, $J=6.6\text{Hz}$), 5.86(1H, d, $J=1.7\text{Hz}$), 6.68-6.83(2H, m), 6.95-7.08(2H, m).

- 5 Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/4): 0.40
(Step 2)




10 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-methoxyphenyl)-5α-androstane

3,3-Ethylenedioxy-17β-methoxymethoxy-11-(4-methoxyphenyl)-5α-androst-11-ene (29.9 mg) was dissolved in ethyl acetate (2 ml) and, after adding acetic acid (0.2 ml) and 10%-palladium/carbon (30 mg), the mixture was stirred for 3 hours at 25°C under hydrogen pressure (25 atm). After filtering the reaction mixture, the solvent was distilled off at reduced pressure and the reaction mixture was purified by silica gel column chromatography (developing solvents: ethyl acetate/dichloromethane = 1/20) to give the end compound in 20.1 mg (yield, 67%).

¹H-NMR(300MHz, CDCl₃)δ: 7.40-7.25(2H, m), 6.75(2H, d, $J=8.2\text{Hz}$), 4.55(3H, s), 3.92(4H, s), 3.78(3H, s), 3.42(1H, dd, $J=6.8, 6.6\text{Hz}$), 3.38-3.28(1H, m), 3.28(3H, s), 2.40-0.80(20H, m), 0.76(3H, s), 0.65(3H, s).

Mass (EI): 484(M+).

(Step 3)




3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-hydroxyphenyl)-
5 5α-androstane

3,3-Ethylenedioxy-17β-methoxymethoxy-11β-(4-methoxyphenyl)-5α-androstane (114.8 mg) and a solution of sodium methanethiolate (69.9 mg) in dimethylformamide (3 mm1) were heated under reflux for 1 hour in a nitrogen atmosphere. After standing to cool, a saturated aqueous solution of ammonium chloride was added to the reaction mixture and extraction with ethyl acetate was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2) gave the end compound in 105.2 mg (yield, 94%).

20 ¹H-NMR(270MHz, CDCl₃)δ: 0.65(3H, s), 0.75(3H, s), 0.90-2.19(20H, m), 3.28(3H, s), 3.24-3.34(1H, m), 3.43(1H, t, J=8.1Hz), 3.91(4H, s), 4.54(2H, s), 4.64(1H, s), 6.64(2H, d, J=8.7Hz), 7.13-7.32(2H, m).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/2): 0.29

(Step 4)



5 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-(2-propen-1-yloxy)phenyl)-5α-androstane


To a solution of 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-hydroxyphenyl)-5α-androstane (56.2 mg) in dimethylformamide (2 ml), sodium hydride (9.6 mg) was added under cooling with ice and the mixture was stirred for 5 minutes. After adding allyl bromide (28.9 mg), the mixture was stirred for 1 hour under cooling with ice. After adding water to the reaction mixture, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/2) gave the end compound in 51.0 mg (yield, 84%).

¹H-NMR(270MHz, CDCl₃)δ: 0.65(3H, s), 0.76(3H, s), 0.82-2.18(20H, m), 3.28(3H, s), 3.24-3.36(1H, m), 3.43(1H, t, J=8.0Hz), 3.90(4H, s), 4.49(2H, d, J=5.3Hz), 4.53(2H, s), 5.26(1H, dd, J=10.5, 1.2Hz), 5.40(1H, dd, J=17.3, 1.5Hz),

5.98-6.13(1H, m), 6.72(2H, d, $J=8.7\text{Hz}$), 7.26(2H, brs).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/2): 0.59

(Step 5)



5


3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-(5-bromo-2-penten-1-yloxy)phenyl)-5α-androstane

In an argon atmosphere, 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-(2-propen-1-yloxy)phenyl)-5α-androstane (16.0 mg), 4-bromo-1-butene (8.5 mg) and benzylidenebistricyclohexylphosphine dichlororuthenium (2.6 mg) were dissolved in dichloromethane (0.3 ml) and the mixture was stirred for 16.5 hours at room temperature. The solvent was distilled off at reduced pressure and purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound in 14.4 mg (yield, 74%).

¹H-NMR(270MHz, CDCl₃)δ: 0.65(3H, s), 0.76(3H, s), 0.81-2.18(20H, m), 2.60-2.78(2H, m), 3.28(3H, s), 3.41(2H, t, $J=6.9\text{Hz}$), 3.25-3.34(1H, m), 3.37-3.48(1H, m), 3.90(4H, s), 4.43-4.51(2H, m), 4.54(2H, s), 5.77-5.88(2H, m), 6.71(2H, d, $J=8.7\text{Hz}$), 7.26(2H, brs).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/4): 0.48

(Step 6)



5 3,3-ethylenedioxy-17β-methoxymethoxy-11β-[4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)-2-penten-1-yloxy}phenyl]-5α-androstane


In a nitrogen atmosphere, 3-ethylenedioxy-17β-methoxymethoxy-11β-{4-(5-bromo-2-penten-1-yloxy)phenyl}-5α-androstane (14.4 mg), 4,4,5,5,5-pentafluoropentane-1-thiol acetate (11.0 mg) and sodium methylate (1.0 M methanol solution) (0.05 ml) were dissolved in methanol (0.2 ml) and tetrahydrofuran (0.2 ml) and the solution was stirred for 17 hours at room temperature. After adding water to the reaction mixture, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; then, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/4) gave the end compound in 14.8 mg (yield, 87%).

1H-NMR(270MHz, CDCl₃)δ: 0.65(3H, s), 0.75(3H, s), 0.81-2.49(26H, m), 2.54-2.67(4H, m), 3.28(3H, s), 3.43(1H, t, J=7.9Hz), 3.24-3.37(1H, m), 3.90(4H, s), 4.45(2H, d, J=4.8Hz), 4.53(2H, s), 5.66-5.94(2H, m), 6.71(2H, d,

$J=8.6\text{Hz}$), 7.26(2H, brs).

Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/4): 0.48

(Step 7)



5

3,3-ethylenedioxy-17β-methoxymethoxy-11β-[4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)-2-pentenyl}oxy}phenyl]-5α-androstane


3,3-Ethylenedioxy-17β-methoxymethoxy-11β-[4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)-2-pentenyl}oxy}phenyl]-5α-androstane (14.8 mg) was dissolved in tetrahydrofuran (0.5 ml) and, after adding OXONE (6.2 mg) and water (0.3 ml) under cooling with ice, the mixture was stirred for 50 minutes. After adding a saturated aqueous solution of sodium hydrogencarbonate to the reaction mixture, extraction with ethyl acetate was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; then, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 4/1) gave the end compound in 14.2 mg (yield, 94%).

$^1\text{H-NMR}(270\text{MHz}, \text{CDCl}_3)\delta$: 0.65(3H, s), 0.75(3H, s), 0.82-

2.37(24H, m), 2.55-2.90(6H, m), 3.28(3H, s), 3.43(1H, t, J=7.8Hz), 3.24-3.37(1H, m), 3.90(4H, s), 4.38-4.51(2H, m), 4.53(2H, s), 5.68-5.95(2H, m), 6.71(2H, d, J=8.6Hz), 7.26(2H, brs).

5 Rf value (on silica gel plate, developing solvents: ethyl acetate/hexane = 1/2): 0.07

(Step 8)



10 17β-hydroxy-11β-[4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)pentenyloxy}phenyl]-5α-androstan-3-one


3,3-Ethylenedioxy-17β-methoxymethoxy-11β-[4-{5-(4,4,5,5,5-pentafluoropentylsulfinyl)-2-pentenyl}oxy}phenyl]-5α-androstane (14.2 mg) was dissolved in ethyl acetate (1 ml) and, after adding 10% palladium/carbon (10 mg), the mixture was stirred for 2 hours at room temperature in a hydrogen atmosphere. After filtering the reaction mixture, the solvent was distilled off at reduced pressure and the resulting residue was dissolved in acetone (2 ml); after adding 1 N-hydrochloric acid (1 ml), the reaction mixture was heated under reflux for 1.5 hours. After standing to cool, a saturated aqueous solution of sodium hydrogencarbonate was added and

extraction with ethyl acetate was effected. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 4/1) gave the end compound in 11.5 mg (yield, 92%).

¹H-NMR(270MHz, CDCl₃)δ: 0.75(3H, s), 0.86(3H, s), 0.82-2.37(31H, m), 2.62-2.87(4H, m), 3.28-3.40(1H, m), 3.48-3.58(1H, m), 3.96(2H, t, J=6.0Hz), 6.72(2H, d, J=8.7Hz), 7.26(2H, brs).


Mass (FAB): 661(M+1).

The following compound was synthesized by a similar method to Example 151.



Example	n	MW	Mass(ESI)
152	7	688	689

[Example 153]



Synthesis of 17 β -hydroxy-11 β -[4-{5-(4,4,5,5,5-pentafluoropentylsulfonyl)pentyloxy}phenyl]-5 α -androstan-3-one


The 17 β -hydroxy-11 β -[4-{7-(4,4,5,5,5-pentafluoropentylsulfinyl)pentyloxy}phenyl]-5 α -androstan-3-one (3.0 mg) obtained in Example 151 was dissolved in tetrahydrofuran (1.0 ml) and, after adding OXONE (2.8 mg) and water (0.5 ml) at room temperature, the mixture was stirred for 1.5 hours. After adding a saturated aqueous solution of sodium hydrogencarbonate to the reaction mixture, extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; then, the solvent was distilled off at reduced pressure.

Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 2/1) gave the end compound in 2.6 mg (yield, 85%).

¹H-NMR(270MHz, CDCl₃) δ : 0.75(3H, s), 0.86(3H, s), 0.78-2.39(31H, m), 2.98-3.09(4H, m), 3.29-3.38(1H, m), 3.48-3.59(1H, m), 3.96(2H, t, J=5.9Hz), 6.72(2H, d, J=8.4Hz), 7.26(2H, brs).


Mass (EI): 676(M⁺).

The following compound was synthesized by a similar method to Example 153.



Example	n	MW	Mass (FAB)
154	7	704	705

[Example 155]



5

Synthesis of 17β-hydroxy-11β-(4-methoxyphenyl)-5α-androstan-3-one


To a solution of 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-methoxyphenyl)-5α-androstane (5.8 mg) in acetone (2 ml), 1 N-hydrochloric acid (1 ml) was added and the mixture was heated under reflux for 1 hour. After standing to cool, a saturated aqueous solution of sodium hydrogencarbonate was added and extraction with ethyl acetate was conducted. The organic layer was washed with a saturated aqueous solution of sodium chloride and dried with magnesium sulfate; after filtering, the solvent was distilled off at reduced pressure. Purification by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/1) gave the end compound in 4.6 mg (yield, 97%).

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ : 0.75 (3H, s), 0.86 (3H, s), 0.82-2.30 (21H, m), 3.31-3.39 (1H, m), 3.48-3.59 (1H, m), 3.79 (3H, s), 6.74 (2H, d, $J=8.7\text{ Hz}$), 7.28 (2H, brs).

Mass (EI): 396 (M^+).


5

[Example 156]



Synthesis of 17β-hydroxy-11β-[4-(3-carboxypropyloxy)phenyl]-5α-androstan-3-one

10 (Step 1)



3,3-ethylenedioxy-17β-methoxymethoxy-11β-[4-(3-tert-butoxycarbonylpropyloxy)phenyl]-5α-androstane


In a nitrogen atmosphere, 3,3-ethylenedioxy-17β-methoxymethoxy-11β-(4-hydroxyphenyl)-5α-androstane (30.2 mg), potassium carbonate (89 mg), 3-bromobutanoic acid t-butyl ester (0.029 ml), potassium iodide (21.3 mg) and 18-crown-6 (200 mg) were dissolved in N,N-dimethylacetamide (0.5 ml) and the mixture was stirred for 10 minutes at 60°C. After adding water to the reaction mixture, extraction was

effected with a solvent system consisting of a mixture of ethyl acetate and hexane. The organic layer was dried with sodium sulfate and, after filtering, the solvents were distilled off at reduced pressure. The resulting residue
 5 was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/5) to give the end compound in 38.1 mg (yield, 97%).

¹H-NMR(270MHz, CDCl₃)δ: 7.40-7.20(2H, m), 6.70(2H, d, J=8.4Hz), 4.54(2H, s), 3.96(2H, t, J=6.1Hz), 3.91(4H, s),
 10 3.46(1H, dd, J=7.9, 8.1Hz), 3.38-3.27(1H, m), 3.28(3H, s), 2.42(2H, t, J=7.3Hz), 2.20-0.82(22H, m), 1.45(9H, s), 0.76(3H, s), 0.65(3H, s).

Mass (EI): 612(M⁺).

(Step 2)



17β-hydroxy-11β-[4-(3-carboxypropyloxy)phenyl]-5α-androstan-3-one

3,3-Ethylenedioxy-17β-methoxymethoxy-11β-[4-(3-t-butoxycarbonylpropyloxy)phenyl]-5α-androstane (38.1 mg) was
 20 dissolved in acetone (1 ml) and, after adding 6 N-hydrochloric acid (0.5 ml), the mixture was heated under reflux for 20 minutes. After adding dichloromethane, the reaction mixture was dried and filtered and the solvent was distilled off at reduced pressure; purification by silica


gel column chromatography (developing solvents: ethyl acetate/n-hexane: 1/1) gave the end compound in 29.0 mg (yield, 100%).

1H-NMR(300MHz, CDCl₃)δ: 7.40-7.10(2H, m), 6.72(2H, d, J=8.8Hz), 3.84(1H, brs), 3.99(2H, t, J=6.0Hz), 3.54(1H, dd, J=7.1, 8.5Hz), 3.33(1H, dd, J=5.8, 6.0Hz), 2.57(2H, t, J=7.1Hz), 2.28-1.86(11H, m), 1.74-1.16(9H, m), 1.08-0.90(2H, m), 0.85(3H, s), 0.74(3H, s).

Mass (ESI): 469(M+1).


10

The following compound was synthesized by a similar method to Example 156.




Example	n	MW	Mass(FAB)
157	7	524	525


15 [Example 158]



17β-hydroxy-11β-[4-(3-aminocarbonylpropyloxy)phenyl]-5α-androstan-3-one




methods to Example 158.




Example	n	R ^N	MW	Mass
159	3	n-pentyl	537	537(EI)
160	7	H	523	524(ESI)
161	7	n-pentyl	593	593(EI)

[Example 162]



Synthesis of 17β-hydroxy-7α-[3-(3-
 5 carboxypropyloxyphenyl)propyl]-5α-androstan-3-one
 (Step 1)



17β-(t-butyldimethylsilyloxy)-7α-[3-(3-benzyloxy)phenyl]-2-
propenyl]-5α-androstan-3-one


10 The 17β-(t-butyldimethylsilyloxy)-7α-(2-propen-1-yl)androstan-3-one (110 mg) obtained in step 1 of Example 3 was dissolved in dichloromethane (0.5 ml) and, after adding 3-benzyloxystyrene (156 mg) and
 15 benzyldienebis(tricyclohexylphosphine)-dichlororuthenium (10.0 mg), the mixture was heated under reflux for 24 hours in an argon atmosphere. After standing to cool, the reaction mixture was concentrated at reduced pressure and

purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/3) to give the end compound in 104.1 mg (yield, 67%).

1H-NMR(300MHz, CDCl₃) δ : 7.49-7.28(5H, m), 7.21(1H, dd, J=7.9, 8.0Hz), 6.98-6.90(2H, m), 6.82(1H, dd, J=1.4, 8.0Hz), 6.30(1H, d, J=5.7Hz), 6.14-6.00(1H, m), 5.07(2H, s), 3.57(1H, dd, J=8.0, 8.5Hz), 2.46-0.92(22H, m), 1.05(3H, s), 0.88(9H, s), 0.74(3H, s), 0.01(6H, s).

Mass (EI): 626(M⁺).

10 (Step 2)



17 β -(t-butyldimethylsilyloxy)-7 α -[3-(3-hydroxyphenyl)propyl]-5 α -androstan-3-one


15 17 β -(t-Butyldimethylsilyloxy)-7 α -[3-(3-benzyloxy)phenyl-2-propenyl]-5 α -androstan-3-one (104.1 mg) was dissolved in ethyl acetate (20 ml) and, after adding acetic acid (0.2 ml) and 10%-palladium/carbon (20 mg), the mixture was stirred for 4 hours at 25 ° in a hydrogen atmosphere. After filtering the reaction mixture, the solvent was distilled off at reduced pressure to give the end compound in 79.8 mg (yield, 89%).

1H-NMR(300MHz, CDCl₃) δ : 7.18-7.09(1H, m), 6.73(1H, d, J=7.7Hz), 6.69-6.62(2H, m), 5.07(1H, s), 3.54(1H, dd, J=8.0, 8.8Hz), 2.66-2.20(5H, m), 2.07-0.85(21H, m), 1.02(3H, s),

0.88(9H, s), 0.70(3H, s), 0.010(3H, s), 0.008(3H, s).

Mass (EI): 538(M⁺).

(Step 3)



5 17β-(t-butyldimethylsilyloxy)-7α-[3-(3-(3-t-butoxycarbonylpropoxy)phenyl)propyl]-5α-androstan-3-one


In a nitrogen atmosphere, 17β-(t-butyldimethylsilyloxy)-7α-[3-(3-hydroxyphenyl)propyl]-5α-androstan-3-one (30.2 mg), potassium carbonate (62 mg), 3-bromobutanoic acid t-butyl ester (0.020 ml) and 18-crown-6 (100 mg) were dissolved in N,N-dimethylacetamide (0.2 ml) and the solution was stirred for 10 minutes at 60°C. After adding water to the reaction mixture, extraction was effected using a solvent system consisting of a mixture of ethyl acetate and hexane. The organic layer was dried with sodium sulfate and, after filtering, the solvents were distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 1/5) to give the end compound in 24.8 mg (yield, 80%).

¹H-NMR(300MHz, CDCl₃)δ: 7.22-7.13(1H, m), 6.79-6.67(3H, m), 3.98(2H, t, J=6.1Hz), 3.54(1H, dd, J=8.0, 8.5Hz), 2.65-2.14(5H, m), 2.43(2H, t, J=7.4Hz), 2.12-0.90(23H, m), 1.45(9H, s), 1.02(3H, s), 0.88(9H, s), 0.71(3H, s),

0.009(3H, s), 0.005(3H, s).

Mass (EI): 567(M+).

(Step 4)




5 17β-hydroxy-7α-[3-{3-(3-carboxypropoxy)phenyl}propyl]-5α-androstan-3-one

17β-(t-Butyldimethylsilyloxy)-7α-[3-{3-(3-t-butoxycarbonylpropoxy)phenyl}propyl]-5α-androstan-3-one (24.8 mg) was dissolved in acetone (4 ml) and, after adding
 10 6 N-hydrochloric acid (1 ml), the mixture was heated under reflux for 2 hours. After distilling off the solvent at reduced pressure, purification was effected by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 2/1) to give the end compound in 19.0 mg
 15 (yield, 100%).

¹H-NMR(300MHz, CDCl₃)δ: 7.16(1H, dd, J=7.1, 8.0Hz), 6.80-6.68(3H, m), 4.02(2H, dt, J=1.4, 6.3Hz), 3.63(1H, dd, J=8.2, 8.8Hz), 3.56(1H, br), 3.13-2.98(4H, m), 2.45-1.95(8H, m), 1.82-0.94(18, m), 1.03(3H, s), 0.74(3H, s).


20 Mass (ESI): 511(M+1).

The following compound was synthesized by a similar method to Example 162.



Example	n	MW	Mass (FAB)
163	4	524	525

[Example 164]



5 Synthesis of 17β-hydroxy-7α-[3-{3-[3-(N-methylaminocarbonyl)propoxy]phenyl}propyl]-5α-androstan-3-one

The 17β-hydroxy-7α-[3-{3-[3-(3-carboxypropoxy)phenyl}propyl]-5α-androstan-3-one (6.6 mg)
 10 obtained in Example 162 was dissolved in tetrahydrofuran (0.5 ml) and, after adding 1-(N,N-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (8.1 mg), 1-hydroxybenzotriazole monohydrate (6.4 mg) and methylamine 40% methanol solution (60 μl), the mixture was stirred for
 15 18 hours at 25°C. After adding ethyl acetate (2.0 ml), the reaction mixture was washed with 1 N-hydrochloric acid, 1 N aqueous solution of sodium hydroxide, and a saturated aqueous solution of sodium chloride. The organic layer was


dried with magnesium sulfate and, after filtering, the solvent was distilled off at reduced pressure. The resulting residue was purified by silica gel column chromatography (developing solvents: ethyl acetate/n-hexane = 5/1) to give the end compound in 0.6 mg (yield, 8.8%).

¹H-NMR(270MHz, CDCl₃)δ: 0.75(3H, s), 1.03(3H, s), 1.05-1.95(22H, m), 1.95-2.21(5H, m), 2.26-2.42(4H, m), 2.51-2.65(2H, m), 2.81(3H, d), 3.63(1H, t, J=8.1Hz), 3.99(2H, t, J=5.9Hz), 6.70(3H, m), 7.17(1H, dd).

Mass(ESI): 524(M+1).


R_f value (on silica gel plate, developing solvents: ethyl acetate/n-hexane = 4/1): 0.21.

The following compounds were synthesized by similar methods to Example 164.



Example	n	R ^M	R ^N	MW	Mass(ESI)
165	4	H	Me	537	538
166	4	Me	Me	551	552
167	4	-(CH ₂) ₄ -		577	578
168	3	Me	Me	537	538
169	3	-(CH ₂) ₄ -		563	564

The following compounds were synthesized by similar methods to Example 114.



Example	n	R ^N	MW	Mass (FAB)
170	5	H	419	420
171	5	n-pentyl	489	490
172	7	H	447	548
173	7	n-pentyl	517	518
174	9	H	475	476
175	9	n-pentyl	545	546
176	11	H	503	504
177	11	n-pentyl	573	574
178	13	H	531	532
179	13	n-pentyl	601	602

[Example 180] Evaluating the agonist and antagonist actions

5 The compound of Example 4 was evaluated for its agonist and antagonist actions in relation to the androgen receptor mediated transcriptional activity.

The agonist action was measured by the same method as described in Example 1; the agonist activity was computed
 10 by the following formula and the determined agonist activity was used to compute the FI₅ value (the concentration for a compound treated group at which it shows a transcriptional activity five times the
 transcriptional activity for the case where the compound is
 15 not added). The compound was added at concentrations of 1, 10, 100, 1000 and 10000 nmol/L.

Agonist activity = Transcriptional activity when
the compound was added/Transcriptional activity
when the compound was not added

The antagonist action was measured by the same method
5 as described in Example 2; the antagonist activity was
computed by the following formula and the determined
antagonist activity was used to compute the IC_{50} value (the
concentration for a compound treated group at which it
shows a 50% decrease in the transcriptional activity of DHT
10 0.1 nmol/L when the compound was not added).. The compound
was added at concentrations of 1, 10, 100, 1000 and 10000
nmol/L and at each of these concentrations, measurement was
done in the presence of DHT (0.1 nmol/L).

15 Antagonist activity = Transcriptional activity when
the compound was added/Transcriptional activity
when the compound was not added x 100

Compound	IC_{50} value (nM)	FI_5 value (nM)
Compound of Example 9	451	ND*
10	937	ND
14	1984	ND
18	342	ND
19	295	ND
20	37	ND
23	1302	ND
24	477	ND
25	415	ND
26	1128	ND
27	421	ND
28	1614	ND
29	304	ND
30	733	ND

42	342	ND
43	1299	ND
46	1751	ND
50	737	ND
51	474	ND
52	277	ND
64	809	ND
65	1831	ND
73	1099	ND
74	2036	ND
96	1601	ND
164	291	ND
167	475	ND
168	540	ND
EM-101	2619	ND
Hydroxyflutamide	31	1000
Bicaltamide	136	767

* ND* in the table signifies that even when the compound was added at a concentration of 10000 nM, the transcriptional activity of the compound-treated group was less than 5 times the transcriptional activity of the control group, making it impossible to compute the FI₅ value.

The above test results verify that existing anti-androgenic agents, hydroxyflutamide and bicaltamide, also exhibit agonist action for the androgen receptor mediated transcriptional activity whereas the compounds of the invention are substantially free of such agonist action for the androgen receptor mediated transcriptional activity. Thus, it is suggested that the compounds of the invention can potentially reduce the development of androgen

tolerance which has been a problem with the conventionally used antiandrogenic agents.

It has also been verified that the compounds of the invention have better agonist action than EM-101. Thus, it is suggested that the compounds of the invention have a sufficient antiandrogenic action to be used as pharmaceuticals and that they can advantageously be used as antiandrogenic agents.

10 INDUSTRIAL APPLICABILITY

The compounds of the invention which are represented by the general formula (I) and the substances of the invention which act as antagonist against but not as agonist for the androgen receptor are potential antiandrogenic agents that do exhibit any side effects such as the development of androgen tolerance due to long-term administration and/or hepatotoxicity and, hence, are expected to be useful as pharmaceutical compositions, say, therapeutics for diseases such as prostate cancer, prostatomegaly, male pattern alopecia, sexual prematurity, acne vulgaris, seborrhea and hirsutism. If the compounds of the invention which are represented by the general formula (I) and the substances of the invention which act as antagonist against but not as agonist for the androgen receptor are preliminarily administered, the onset of diseases such as prostate cancer, prostatomegaly, male pattern alopecia, sexual prematurity, acne vulgaris, seborrhea and hirsutism can hopefully be prevented or

retarded, so they are also potential preventives of these diseases. Further, the compounds of the invention which are represented by the general formula (I) and the substances of the invention which act as antagonist against
5 but not as agonist for the androgen receptor have toxicity such as cytotoxicity sufficiently reduced that they are expected to find advantageous use as therapeutics and/or preventives of the diseases mentioned above.